

Hyperbaric Oxygen Therapy Reduces the Risk of QTc Interval Prolongation in Patients with Diabetes and Hard-to-Heal Foot Ulcers

Katarina Fagher, Per Katzman, Magnus Löndahl

PII: S1056-8727(15)00295-0
DOI: doi: 10.1016/j.jdiacomp.2015.07.023
Reference: JDC 6511

To appear in: *Journal of Diabetes and Its Complications*

Received date: 28 April 2015
Revised date: 23 July 2015
Accepted date: 24 July 2015

Please cite this article as: Fagher, K., Katzman, P. & Löndahl, M., Hyperbaric Oxygen Therapy Reduces the Risk of QTc Interval Prolongation in Patients with Diabetes and Hard-to-Heal Foot Ulcers, *Journal of Diabetes and Its Complications* (2015), doi: 10.1016/j.jdiacomp.2015.07.023

Hyperbaric Oxygen Therapy Reduces the Risk of QTc Interval Prolongation in Patients with Diabetes and Hard-to-Heal Foot Ulcers.

Katarina Fagher MD[1,2], Per Katzman MD, PhD[1,2], Magnus Löndahl MD, PhD[1,2]

[1]Clinical Sciences in Lund, Lund University, Lund, Sweden,

[2] Department of Endocrinology, Skåne University Hospital, Sweden

Corresponding author:

Dr Katarina Fagher

Department of Endocrinology

Skåne University Hospital

S-22185 Lund

Sweden

Phone: +46 46 176749

Fax: +46 46 176024

Email: Katarina.fagher@med.lu.se

Number of words: 2459 (200 in abstract), 2 tables, 2 figures