




Contents lists available at ScienceDirect

## Molecular & Biochemical Parasitology



Short technical report

# A vanillic acid inducible expression system for *Trypanosoma brucei*

Jack D. Sunter\*

Sir William Dunn School of Pathology, University of Oxford, South Parks Road, OX1 3RE, UK

### ARTICLE INFO

#### Article history:

Received 24 February 2016

Received in revised form 31 March 2016

Accepted 1 April 2016

Available online xxx

#### Keywords:

*Trypanosoma brucei*

Inducible expression

Gene regulation

Vanillic acid

### ABSTRACT

Reverse genetics in *Trypanosoma brucei* is dependent on the tetracycline inducible system for the precise control over the expression of both genes and dsRNA. Another independent inducible system for trypanosomes would enable the control of the activities of two different genes in the same cell, providing greater experimental sophistication. Here, I describe the development of the vanillic acid based inducible expression system for *T. brucei*, which operates independently of, and can be used in parallel with the tetracycline inducible system.

© 2016 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

*Trypanosoma brucei* has a well annotated genome and well developed reverse genetic tools, making it a highly tractable experimental organism used for the study of a large range of biological processes such as antigenic variation and the eukaryotic flagellum [1,2]. Precise control over the expression of exogenous genes, mutants and dsRNA is critical for reverse genetics. In *T. brucei* inducible expression is controlled by the tetracycline inducible system in the tetON configuration (Fig. 1A), where tet operator (TetO) sites are located downstream of a strong promoter and the binding of tet repressor (TetR) to the TetO sites inhibits transcription from the promoter [3,4]. Addition of tetracycline releases this inhibition by binding to the TetR, which is now unable to bind to the TetO sites, allowing transcription to occur. The successful operation of the tetracycline inducible system requires the expression of TetR; there are now several *T. brucei* cell lines expressing these proteins and a few examples are referenced here [4–6].

Over the last 20 years the tetracycline inducible system in *T. brucei* has been extremely effective in enabling functional genomic research. Another inducible system that operated simultaneously with the tetracycline based system would greatly increase experimental flexibility, allowing the independent control of two different gene activities in the same cell, hence creating higher-order control networks. An IPTG inducible system has previously been described in trypanosomes [7]; however this was only used for the expression of a tRNA. Here, I describe the development of

an inducible expression system for expression of both protein and dsRNA in procyclic form (PCF) *T. brucei* using the components that control vanillic acid metabolism in *Caulobacter crescentus*, which has successfully been used for inducible gene expression in mammalian cells [8].

Inducible gene expression using the vanillic acid system requires the expression of the vanillic acid repressor (VanR) [8]. To achieve this I modified the pSMOX plasmid, which encodes both TetR and T7 RNAP, by inserting a trypanosome codon-optimised VanR gene with a C-terminal nuclear localisation signal to create pJ1173 (Fig. 1B) [6,9]. This single plasmid now encodes the T7 RNAP, TetR and VanR proteins and was integrated into the tubulin locus in the TREU927 PCF cell line [6]. The other essential component of the vanillic acid system is a strong promoter with vanillic acid operator (VanO) sites downstream of it (Fig. 1C). I used the pDEX877 plasmid, a derivative of the pDEX777 that encodes the blasticidin resistance gene as the starting point to create the vanillic acid inducible plasmid, pJ1271 [6]. The tetO site in pDEX877 was replaced with five VanO sites and as the VanO site contains a BamHI site, the BamHI site that is located at the 3' end of the eGFP-Ty construct was replaced with a BglII site ensuring that the plasmid remains compatible with other plasmid systems (Fig. 1A) [5,6,10]. The plasmid, pJ1271 targets the 177 bp repeats like other pDEX plasmid derivatives.

pJ1271 was integrated into the 927 pJ1173 cells to give a cell line with the expression of eGFP under vanillic acid control. To determine whether this cell line (927 pJ1173 pJ1271) was able to express eGFP a range of vanillic acid concentrations were added to the cells before analysing the eGFP expression level by flow cytometry (Fig. 1D). Without vanillic acid the fluorescent signal from the


Abbreviations: TetO, tet operator; RNAP, RNA polymerase; TetR, tetracycline repressor; VanR, vanillic acid repressor; VanO, vanillic acid operator.

\* Corresponding author.

E-mail address: [jack.sunter@path.ox.ac.uk](mailto:jack.sunter@path.ox.ac.uk)

<http://dx.doi.org/10.1016/j.molbiopara.2016.04.001>

0166-6851/© 2016 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).



**Fig. 1.** Vanillic acid inducible system is functional and independent of the tetracycline inducible system. A) Cartoon showing how the tetON and vanON inducible systems operate in *T. brucei* with the plasmids pDEX877 and pJ1271. B) Map of the modified pSMOX2 plasmid, pJ1173 that encodes the vanillic acid repressor protein (VanR), tetracycline repressor protein (TetR), T7 RNA polymerase (RNAP) and puromycin N-acetyl-transferase (pac). C) The sequence of the 5 VanO sites introduced into pJ1271. D) Titration of vanillic acid concentration shows that eGFP expression was induced after addition of vanillic acid and was tunable. Cells were incubated for 24 h with a range of different vanillic acid concentrations, and eGFP expression was measured by flow cytometry with 50,000 events captured per concentration. Vanillic acid was purchased from Sigma-Aldrich and dissolved in DMSO. E) and F) Comparison of eGFP expression induction by vanillic acid and doxycycline (final concentration of 1 μg/ml was used for all experiments) from pJ1271 and pDEX877. eGFP expression was induced by incubating the cells for 24 h with vanillic acid, doxycycline or both, and was measured by flow cytometry with 50,000 events captured per induction. A histogram (E) and graph of normalised eGFP fluorescence intensity (FI) (F) is shown. This experiment was repeated three times for pJ1271, error bars are ±S.D. G) Growth curves for the 927 pJ1173 pJ1271 and 927 pJ1173 pDEX877 cell lines induced with vanillic acid, doxycycline or both over a 96 h period. Every 24 h the cells were split to 1 × 10<sup>6</sup> cells/ml and fresh vanillic acid and doxycycline added.

927 pJ1173 pJ1271 cells was indistinguishable from the 927 pJ1173 cell line; no protein expression was therefore detectable under non-induced conditions, demonstrating that the vanillic acid system was able to repress expression. Vanillic acid addition led to the inducible expression of eGFP with increasing vanillic acid giving higher levels of eGFP expression up to ~125 μM vanillic acid after

which eGFP expression plateaued. Based on this titration 250 μM vanillic acid was used for all subsequent inductions to ensure maximal expression.

To compare the relative expression levels between the vanillic acid and tetracycline inducible systems the pDEX877 plasmid was integrated into the 927 pJ1173 cell line to create 927 pJ1173

Download English Version:

<https://daneshyari.com/en/article/5915309>

Download Persian Version:

<https://daneshyari.com/article/5915309>

[Daneshyari.com](https://daneshyari.com)