Severe hypoxia during incremental exercise to exhaustion provokes negative post-exercise affects

Michail E. Keramidas, Nektarios A.M. Stavrou, Stylianos N. Kounalakis, Ola Eiken, Igor B. Mekjavic

PII: S0031-9384(16)30020-8
DOI: doi: 10.1016/j.physbeh.2016.01.021
Reference: PHB 11170

To appear in: *Physiology & Behavior*

Received date: 20 March 2015
Revised date: 9 November 2015
Accepted date: 19 January 2016

Please cite this article as: Keramidas Michail E., Stavrou Nektarios A.M., Kounalakis Stylianos N., Eiken Ola, Mekjavic Igor B., Severe hypoxia during incremental exercise to exhaustion provokes negative post-exercise affects, *Physiology & Behavior* (2016), doi: 10.1016/j.physbeh.2016.01.021

Severe hypoxia during incremental exercise to exhaustion provokes negative post-exercise affects

Authors

Michail E. Keramidas[a,b], Nektarios A. M. Stavrou[c,d], Stylianos N. Kounalakis[b], Ola Eiken[a], Igor B. Mekjavic[b]

[a] Department of Environmental Physiology, Swedish Aerospace Physiology Center, School of Technology and Health, Royal Institute of Technology, Stockholm, Sweden

[b] Department of Automation, Biocybernetics and Robotics, Jozef Stefan Institute, Ljubljana, Slovenia

[c] Exercise and Sport Science Department, ASPETAR Orthopaedic and Sports Medicine Hospital, Doha, Quatar

[d] Faculty of Physical Education and Sport Science, University of Athens, Athens, Greece

Correspondence

Michail E. Keramidas

Department of Environmental Physiology

Swedish Aerospace Physiology Center

School of Technology and Health, Royal Institute of Technology

Berzelius väg 13, SE-171 65, Solna, Sweden

Phone: +46 8 524 839 69

Fax: +46 8 330923

Email: michail.keramidas@sth.kth.se