

Emerging Role of Endoscopic Ultrasound in the Diagnostic Evaluation of Idiopathic Pancreatitis

Ioana Smith, MD, Jayapal Ramesh, MD, Kondal R. Kyanam Kabir Baig, MD, Klaus Mönkemüller, MD, PhD and C. Mel Wilcox, MD

Abstract: *Background:* “Idiopathic pancreatitis” is diagnosed when clinical, laboratory and conventional radiologic methods do not provide a clear etiology for the episode. Given its associated morbidity and mortality, it is important to determine the cause of pancreatitis to provide early treatment and prevent recurrence. *Methods:* The aim of this systematic review was to evaluate the utility of endoscopic ultrasound (EUS) in determining an etiology in patients classified as having idiopathic pancreatitis and to assess how EUS performed compared with other modalities. A PubMed search for relevant articles (January 2000–November 2014) was performed using the search terms “(pancreatitis or idiopathic pancreatitis or unexplained pancreatitis) and (EUS or endoscopic ultrasound).” *Results:* The search yielded a total of 963 articles, and 13 studies were included in the final review. In some studies, the yield of EUS was higher than magnetic resonance cholangiopancreatography in idiopathic pancreatitis. EUS more accurately detected biliary stones, whereas magnetic resonance cholangiopancreatography more often identified pancreatic duct abnormalities. The yield of EUS was lower in patients postcholecystectomy but was not influenced by gender, severity of pancreatitis, or recurrent disease. The most frequent diagnoses by EUS for those with idiopathic pancreatitis were biliary tract disease (41%). Overall, EUS identified additional diagnostic information in 61% of patients with idiopathic pancreatitis. *Conclusions:* Given the high incidence of microlithiasis and/or biliary sludge as a cause of idiopathic pancreatitis as well as the safety and high accuracy, EUS should be considered 1st for evaluation of idiopathic pancreatitis if conventional cross-sectional radiography fails to reveal a cause.

Key Indexing Terms: Endoscopic; Ultrasound; Idiopathic pancreatitis; Pancreatitis; Endoscopic ultrasonography. [*Am J Med Sci* 2015;350(3):229–234.]

Alcohol and gallstone disease are responsible for most cases of acute pancreatitis. Patients are classified as having idiopathic pancreatitis after clinical (history), laboratory studies (triglyceride and calcium level) and conventional radiological methods (transabdominal ultrasound and computed tomography [CT]) do not reveal an etiology of the pancreatitis. Idiopathic pancreatitis is diagnosed in 10% to 30%^{1–4} of acute pancreatitis episodes. Recent studies have suggested that microlithiasis is a cause of unexplained pancreatitis in up to 75% of patients with an intact gallbladder.⁵ Microlithiasis, defined as a stone less than 3 mm in diameter,¹ may be undetected by transabdominal ultrasound. Sphincter of Oddi dysfunction (SOD) can be identified in up to 30% of patients with idiopathic pancreatitis.⁶ Given the associated morbidity and mortality, it is important to

determine the cause of pancreatitis to provide therapy early and prevent further recurrence. Pancreatitis with or without a known cause may recur in 30% of patients with a bout of pancreatitis, and up to 30% of these cases remain undiagnosed.³

Endoscopic ultrasonography (EUS) has a high diagnostic accuracy for biliary pathology including cholelithiasis, biliary sludge and choledocholithiasis,⁷ pancreatic disease and pancreatic divisum⁸ but lacks in diagnosing SOD.³ Magnetic resonance cholangiopancreatography (MRCP) with secretin administration is a promising and suitable alternative diagnostic modality but may not be widely available.⁹ Endoscopic retrograde cholangiopancreatography (ERCP) can be used in combination with bile crystal analysis with or without sphincter of Oddi manometry and such an approach has led to identifying the underlying etiology in 30% to 60% of cases, but ERCP carries a risk of pancreatitis.⁴

To better understand the diagnostic role of EUS in identifying an etiology of idiopathic pancreatitis, the authors reviewed studies that included patients with idiopathic pancreatitis and used EUS with or without a comparator such as MRCP, ERCP, and/or bile crystal analysis to establish a cause of the pancreatitis. Second, we sought to identify clinical factors (previous cholecystectomy, single versus recurrent pancreatitis) that may affect the diagnostic yield of EUS.

METHODS

Review Protocol and Study Eligibility

The Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) checklist¹⁰ served as a guideline and was used by 2 authors (I.S. and C.W.) for the execution of this systematic review.

Eligibility Criteria

Definitions

Idiopathic pancreatitis was defined as acute pancreatitis after clinical, laboratory, and conventional radiological methods failed to reveal an etiology.

Identification of Relevant Studies

A systematic search was conducted on PubMed to identify articles published between January 1, 2000, and November 1, 2014. The following search terms were used: (“pancreatitis” [MESH] OR “idiopathic pancreatitis” [MESH] OR “unexplained pancreatitis” [All Fields]) AND (“EUS” [MESH] OR “endoscopic ultrasound” [All Fields]). Two authors (I.S. and C.W.) independently searched and abstracted the data according to standard methods.¹¹ Additionally, the references in each of the selected articles were searched to identify any missed references.

Study Selection

Eligible studies could be either retrospective or prospective and had to have enrolled at least 5 patients with

From the Division of Gastroenterology and Hepatology, University of Alabama School of Medicine, Birmingham, Alabama.

Submitted March 11, 2015; accepted in revised form April 21, 2015.

The authors have no financial or other conflicts of interest to disclose.

Correspondence: C. Mel Wilcox, MD, Division of Gastroenterology and Hepatology, University of Alabama School of Medicine, 1720 2nd Avenue South, BDB 380, Birmingham, AL 35294-0113 (E-mail: melw@uab.edu).




FIGURE 1. Flowchart of the search strategy and selection of studies eligible for inclusion in review.

idiopathic pancreatitis who underwent EUS to further investigate the etiology of the acute or recurrent attack when CT, ERCP, MRCP, ultrasound, and/or repeated ultrasound had failed to reveal a cause of pancreatitis.

Data Extraction and Analysis

Two investigators (I.S. and C.W.) performed the searches and reviewed the articles for appropriateness of inclusion into the study. The data extracted from the studies included authors, year of publication, number of patients included in the study, imaging before EUS, mean age, gender, follow-up duration in months, and EUS-quantified determined causes. Total number of patients was defined as those with idiopathic pancreatitis that were included in the study of which a defined number had to undergo EUS for further investigation. For each study, we calculated the overall yield of EUS in attaining a diagnosis for patients with idiopathic pancreatitis.

RESULTS

A total of 963 records were initially identified in the literature search (Figure 1). Four additional records were

identified by means of hand searching of reference lists. All 967 records were screened based on title and abstract. After careful review and using the aforementioned inclusion and exclusion criteria, 13 studies were included in the final review.^{1,4,7,8,12-20} Table 1 outlines studies evaluating EUS in idiopathic pancreatitis and Table 2 shows the EUS-identified causes of the pancreatitis. Representative images are demonstrated in Figures 2–6. The most frequent diagnosis after EUS in patients with idiopathic pancreatitis was biliary tract disease (biliary stones, microlithiasis and sludge) (41.1% ± 23.7%). EUS detected pancreatic disease (chronic pancreatitis, pancreatic divisum, pancreatic mass, pancreatic cancer, pancreatic parenchymal change and/or pancreatic ductal change) in 22.1% ± 26.6% of patients with idiopathic pancreatitis. Overall, EUS identified additional diagnostic information in 61.0% ± 18.0% of patients with idiopathic pancreatitis, with 41% having biliary tract disease. In one series, EUS demonstrated a high yield for occult gallbladder disease (6%) as well as the importance of chronic pancreatitis (45%).⁴ Refer to Figure 2 for a proposed approach to the evaluation of idiopathic pancreatitis.

Single Versus Multiple Attacks

Overall, the yield of EUS was not influenced by recurrent disease. In the study by Yusoff et al,⁸ chronic pancreatitis was the most commonly identified abnormality found (16.4%–42.0%) and was approximately twice as frequent in patients with recurrent episodes versus a single episode of idiopathic pancreatitis (no-cholecystectomy: 42.0% versus 21.6%, *P* = 0.0008; post-cholecystectomy: 38.6% versus 16.4%, *P* = 0.008). In one study, biliary pancreatitis (cholelithiasis, choledocholithiasis and microlithiasis) was the most common etiology of recurrent acute pancreatitis (37%).²¹ In another study, the probability of harboring occult gallstones was also significantly associated with age, recurrent pancreatitis and altered liver function tests results during an index episode.²² In contrast, in one study, the yield of positive EUS findings did not differ between single or recurrent idiopathic pancreatitis.⁸ In another study, the yield of EUS was not influenced by sex, severity of pancreatitis or recurrent disease.¹³

TABLE 1. Studies evaluating EUS in idiopathic pancreatitis

Study	No. of patients	Year	Overall yield of EUS, %	Follow-up, month	Imaging before EUS	Mean age, year	Gender (female), %
Liu et al ¹²	18	2000	78	22 (median)	CT, ERCP, US, repeated US	*	*
Ardengh et al ¹	36	2010	75	*	CT, US	47.1	58.4
Vila et al ¹³	44	2010	79	29 (mean)	CT, US	61.5	29.5
Yusoff et al ⁸	370	2004	59	*	CT, US	53.4	55.4
Tandon and Topazian ⁴	31	2001	68	16 (mean)	CT, US, MRI, MRCP, EUS	48.8	61.2
Frossard et al ¹⁴	168	2000	78	25 (mean)	US	50.0	39.0
Norton and Alderson ¹⁵	44	2000	73	3–28 (range)	US	53.5 (median)	54.5
Thevenot et al ¹⁶	41	2013	29	22 (mean)	CT, US	55.3	37.5
Zhang et al ¹⁷	33	2011	42	*	CT, US, MRCP	46.5	60.6
Kim et al ¹⁸	31	2011	42	36 (mean)	US, CT, MRCP, ERCP, SOM	51.3 (median)	64.5
Ortega et al ⁷	49	2011	51	16 (mean)	CT, US	58.0	51.0
Morris-Stiff et al ¹⁹	42	2009	41	74 (median)	CT, US, ERCP, MRCP	53.0	40.5
Mariani et al ²⁰	44	2009	80	*	US, CT	48.9	54.5

* Follow up not provided.

CT, computed tomography; ERCP, endoscopic retrograde cholangiopancreatography; MRCP, magnetic resonance cholangiopancreatography; MRI, magnetic resonance imaging; SOM, sphincter of oddi manometry.

Download English Version:

<https://daneshyari.com/en/article/5931657>

Download Persian Version:

<https://daneshyari.com/article/5931657>

[Daneshyari.com](https://daneshyari.com)