



Left ventricular filling patterns and its relation to left ventricular untwist in patients with type 1 diabetes and normal ejection fraction

G. Nallur Shivu^{a,*}, K. Abozguia^b, T.T. Phan^b, P. Narendran^b, M. Stevens^b, M. Frenneaux^c

^a University Hospital of Wales, Cardiff, UK

^b University of Birmingham, Birmingham, UK

^c School of Medicine and Dentistry, University of Aberdeen, UK

ARTICLE INFO

Article history:

Received 18 September 2011

Received in revised form 20 November 2011

Accepted 17 December 2011

Available online 11 January 2012

Keywords:

Untwisting

Type 1 diabetes mellitus

Diabetic cardiomyopathy

Left ventricular filling

Left atrial function

ABSTRACT

Background: We evaluated young patients with type 1 diabetes (T1DM) who had normal left ventricular (LV) ejection fraction and used speckle tracking echocardiography to assess changes in LV untwisting. We used cardiac magnetic resonance imaging (MRI) to assess the LV filling patterns in these subjects.

Methods: We recruited 33 T1DM patients and 32 age-matched healthy controls (HC) into the study. Study participants underwent echocardiography, cardiac MRI and metabolic exercise testing.

Results: The early peak LV untwisting rate (E) was similar in T1DM and HC (-11.9 ± 4.6 0/cm/s vs -11.3 ± 4.7 0/cm/s, $P=0.29$) but the late peak LV untwisting rate (A) was significantly increased in T1DM (-6.2 ± 3 0/cm/s vs -4.9 ± 3.9 0/cm/s, $P<0.05$). The time to early peak untwisting rate was not different ($50.9 \pm 9.6\%$ vs $48.4 \pm 7.3\%$, $P=0.12$) but the time to late peak untwisting rate was significantly delayed in T1DM patients ($80.4 \pm 12.5\%$ vs $72.7 \pm 14.6\%$, $P<0.05$). The LV filling patterns demonstrated a significantly increased left atrial (LA) contribution to LV filling in T1DM. On linear regression peak late filling rate ($r=0.60$, $P<0.000$), trans-mitral A wave ($r=0.25$, $P<0.05$) and A' ($r=0.30$, $P<0.01$) were predictors of LA contribution to LV filling.

Conclusion: We demonstrate for the first time using speckle tracking that LV untwisting rate E is preserved and untwisting rate A is increased and delayed in young patients with uncomplicated T1DM. The LA contribution to LV filling is increased in these patients and is directly related to increases in other indices of LA function like peak late filling rate, trans-mitral A wave and A'.

© 2011 Elsevier Ireland Ltd. All rights reserved.

1. Introduction

Diabetes is a metabolic disorder characterized by hyperglycemia and insulin deficiency. The prevalence of type 2 diabetes (T2DM) in particular is rapidly increasing throughout the western world [1–3]. Heart failure occurs more frequently in diabetes [4] and although frequently due to coronary artery disease (CAD) [5–7] and/or hypertension, may occur in the absence of these, when it is termed diabetic cardiomyopathy. Although there is increasing evidence for the presence of diabetic cardiomyopathy as a separate entity, detection of early changes in the myocardium is challenging in patients with diabetes. Previous studies with 2D and tissue Doppler echocardiography have demonstrated abnormalities in various diastolic parameters

prior to the onset of overt systolic dysfunction [8–10]. Echocardiographic trans-mitral pulsed wave Doppler demonstrates abnormalities in mitral inflow velocity and deceleration time (DecT), isovolumetric relaxation time and filling patterns [11]. In healthy young subjects the majority of left ventricular (LV) filling is accomplished in the early filling phase and the left atrium (LA) contributes the remainder. In heart failure the early phase of filling is hampered by the stiff ventricle [12,13]. In this situation the LA contributes significantly more to LV filling. Similarly there is increased reliance on the LA for LV filling in the presence of diastolic dysfunction. We demonstrated recently that increased LA filling compensates for impaired early relaxation during exercise in patients with heart failure with preserved ejection fraction [14]. LA function may be an important factor in maintaining ejection fraction in the early stages of diabetic cardiomyopathy. Loss of atrial function with the onset of atrial fibrillation may precipitate worsening heart failure in any of the above situations.

In this study we investigated various aspects of LV diastolic function in young T1DM subjects with no underlying heart failure or CAD. Speckle tracking echocardiography has recently been used to measure LV torsion and untwisting. LV untwisting, which follows LV

Abbreviations: T1DM, type 1 diabetes; LV, left ventricular; MRI, magnetic resonance imaging; LA, left atrial; HC, healthy controls; T2DM, type 2 diabetes; CAD, coronary artery disease; DecT, deceleration time; EF, ejection fraction; EDV, end-diastolic volume; ESV, end-systolic volume; PFR, peak filling rates.

* Corresponding author at: Department of Cardiology, University Hospital of Wales, Heath Park, Cardiff, CF14 4XW, UK. Tel.: +44 2920 747747; fax: +44 2920 522172.

E-mail address: drgani23@gmail.com (G.N. Shivu).

torsion contributes significantly to LV filling. We studied the untwisting patterns in patients with T1DM using speckle tracking echocardiography. The early untwisting corresponds to the early phase of filling and late untwisting corresponds to the LA filling phase. We used cardiac MRI to directly measure LV volumes during the cardiac cycle and hence LV filling patterns in these subjects.

1.1. Aims of the study

1. To assess the LV untwisting patterns and other parameters of diastolic function in type 1 diabetes patients in the absence of CAD and heart failure.
2. To assess the LV diastolic filling patterns in these subjects as compared to healthy controls and calculate the LA contribution to LV filling.
3. To assess the relationship between LA contribution to LV filling and other parameters of LA function and LV diastolic function.

1.2. Hypotheses of the study

1. LV torsion and untwist are altered in patients with type 1 diabetes prior to onset of overt heart failure.
2. Increased reliance on LA for LV filling which is related to the diastolic dysfunction in these subjects with no overt heart failure.
3. Study of LV untwisting patterns and LV filling will provide insights into early changes in diastolic function in diabetes.

2. Methods

This was a prospective study in patients with type 1 diabetes as compared to HC. All subjects were fasted overnight and fasting blood samples were taken. Following this, subjects were allowed to have a light breakfast and the diabetes patients had their morning dose of insulin. All subjects then underwent echocardiography, cardiac MRI and metabolic testing on the same day which completed their participation in the study. All the investigations were undertaken in the University of Birmingham and the project was approved by Multicenter Regional Ethics Committee in Birmingham. Data on LV torsion in these subjects have been published previously by our group [15].

2.1. Patients

We recruited 33 subjects who met the following criteria from the Heart of England NHS Foundation Trust and University Hospital Birmingham NHS Trust, Birmingham, UK:

2.2. Inclusion criteria

- Patients with T1DM according to WHO definition with HbA1C <10%.
- No history of chest pain or breathlessness.
- No evidence of CAD or heart failure based on history, 12 lead electrocardiogram, a normal ejection fraction (EF) on echocardiography and metabolic exercise testing.
- Above 18 years of age.
- Able to provide informed consent.

2.3. Exclusion criteria

- Less than 18 years of age or unable to provide informed consent.
- Previous history of CAD, heart failure or renal failure.
- Pregnant subjects.

2.4. Healthy controls

32 age and sex matched controls with no cardiac history or diabetes mellitus were recruited. All HC had a normal 12 lead electrocardiogram, echocardiogram and metabolic exercise test. HC were recruited by general adverts in the University of Birmingham and blood banks.

2.5. Echocardiography

Echocardiography was performed with participants in the left lateral decubitus position with a Vivid 7 (GE Vingmed) echocardiographic machine and a 2.5-MHz transducer. Standard echocardiographic views were obtained from parasternal (short axis gray scale images at mitral, papillary and apical levels) and apical (4 and 2 chamber views) windows. Trans-mitral flow profiles and tissue Doppler measurements of mitral annular velocities were performed. The ejection fraction was calculated with

Simpson's rule [16]. The left atrial volume was determined using the area-length method. The gray scale images obtained in the parasternal short axis view were used to compute LV rotation using commercially available speckle tracking software on an Echopac system.

2.6. Metabolic exercise testing

Subjects underwent a symptom-limited erect treadmill exercise testing using a standard ramp protocol with simultaneous respiratory gas analysis on a Schiller CS-200 Ergo-Spiro exercise machine. Samplings of expired gases were performed continuously, and data were expressed as 30-second means. Minute ventilation, oxygen consumption, carbon dioxide production, and respiratory exchange ratio (RER) were obtained. Peak oxygen consumption (VO₂ max) was defined as the highest value of oxygen consumption measured during the exercise period. Blood pressure and ECG were monitored throughout. Subjects were encouraged to exercise to exhaustion with a minimal requirement of RER > 1.

2.7. Cardiac MRI

Cardiac MRI was performed on a 3 T Phillips Achieva MRI scanner. The subjects were positioned supine with a dedicated cardiac sense coil wrapped around the chest. The subjects were positioned in the scanner with their heart at the isocenter of the magnet. Survey images were obtained and the short axis images were planned from this. The LV was divided into slices of 8 mm with a gap of 2 mm. A total of 12 slices ensured complete coverage of the LV from apex to base. Short axis cine images were acquired at breath hold using sense cine sequences.

3. Analysis

3.1. Speckle tracking echocardiography (STE)

Myocardial deformation was measured using a commercially available speckle tracking system on an ECHOPAC (version 4.2.0) workstation. In this system, the displacement of speckles of myocardium in each spot was analyzed and tracked from frame to frame. We selected the best-quality digital two-dimensional image and the LV endocardium was traced at end-systole. The region of interest width was adjusted as required to fit the wall thickness. The software package then automatically tracked the motion through the rest of the cardiac cycle. The onset of the QRS complex was taken as the beginning of systole. Adequate tracking was verified in real time. Counter-clockwise rotation was marked as a positive value and clockwise rotation as a negative value when viewed from the apex. In order to calculate LV torsion, torsion rate and untwist rates, the rotation traces of the basal and apical LV cross-sections were exported into DPlot graph software (Version 2.2.1.4, HydeSoft Computing, LLC, Vicksburg, USA). The LV twist curve was generated by calculating the difference between apical and basal rotations at each corresponding time point. LV untwist rates were derived from the first derivative of the LV twist curve (Fig. 1). Peak LV torsion was derived from LV twist divided by LV diastolic longitudinal length. In order to adjust for the differences in heart rate between individuals the RR interval was normalized to 100%. Of the 65 subjects in the study, 61 (94%) subjects had both adequate LV basal and apical images for speckle tracking to complete analysis of all LV rotational parameters.

3.2. Reproducibility of STE

Reproducibility of STE determined within our department using a randomly selected group of 10 controls has been published previously [17]. Inter-observer measurement variability was determined by two independent observers who measured LV torsion in the 10 controls. To obtain the intra-observer variability, the first observer performed the analysis on two separate occasions 1 month apart. We performed Bland-Altman plots to assess variability of measurement. Our results showed that for LV torsion, intra-observer reproducibility was 0.24 ± 0.58 (bias ± 1.96 standard deviation of the difference (SD)) and inter-observer reproducibility was 0.15 ± 0.69 (bias ± 1.96 SD), which are acceptable.

Download English Version:

<https://daneshyari.com/en/article/5975820>

Download Persian Version:

<https://daneshyari.com/article/5975820>

[Daneshyari.com](https://daneshyari.com)