



ELSEVIER

Contents lists available at ScienceDirect

Journal of Cardiology Cases

journal homepage: www.elsevier.com/locate/jccase

Case Report

Stress-induced cardiomyopathy accompanied by heat stroke



Chiaki Watanabe (MD, PhD)*, Naoto Nishina (MD), Chuichi Kawai (MD, PhD, FJCC)

Department of Cardiology, Takeda General Hospital, Kyoto, Japan

ARTICLE INFO

Article history:

Received 8 December 2014
 Received in revised form 10 March 2015
 Accepted 25 March 2015

Keywords:

Stress-induced cardiomyopathy
 Takotubo-cardiomyopathy
 Heat stroke
 Ventricular tachycardia
 Aberrant conduction

ABSTRACT

Heat stroke is a severe illness characterized by a core temperature $>40^{\circ}\text{C}$ (104°F) and central nervous system abnormalities, such as delirium, convulsions, or coma, resulting from exposure to environmental heat or strenuous physical experience. We describe a case with high body temperature and coma, who also presented with wide QRS tachycardia and shock. He was initially treated for heat stroke and wide QRS tachycardia. Serial electrocardiographic change and echocardiographic findings suggested this was complicated with stress-induced cardiomyopathy. This case demonstrates the importance of recognizing that serious myocardial damage can result from heat stroke.

<Learning objective: Heat stroke can cause multiple organ failure and the presentation of circulatory failure in heat stroke may be the sign of myocardial dysfunction. To distinguish acute coronary syndrome and stress-induced cardiomyopathy, both of which could be evoked by heat stroke, the evaluation of coronary artery is necessary. Stress-induced cardiomyopathy may cause lethal arrhythmia or circulatory collapse in acute phase. Invasive circulatory monitoring is recommended in patients with heat stroke.>

© 2015 Japanese College of Cardiology. Published by Elsevier Ltd. All rights reserved.

Introduction

Heat stroke is a medical emergency commonly encountered in small children and socioeconomically deprived elderly people, but can also occur in healthy young persons after strenuous activities or in specific environments. The increase in victims of heat stroke, even in warm climates, may be attributed to global warming. Heat stroke can lead to multi-organ failure (MOF) and cause myocardial damage.

On the other hand, stress-induced (takotsubo) cardiomyopathy is a transient myocardial dysfunction precipitated by physical or emotional stress, mimicking the presentation of acute coronary syndrome [1].

Here, we report a case of heat stroke presenting with wide QRS tachycardia and shock, the cause of which was considered to be stress-induced cardiomyopathy.

Case report

An 87-year-old man was brought to our emergency department because of consciousness disturbance and generalized convulsion on a hot summer morning. He had a 30-year history of epilepsy and hypertension, which had been treated by a neurologist until

17 months previously, with no subsequent follow-ups and medications. He lived in a house without air conditioning. He was barely able to walk indoors, and had not been eating properly recently. On the morning of admission, his son found him conscious but immobile in the bathroom. His son called an ambulance because the patient gradually became unresponsive and had a convulsion.

On physical examination, he was unconscious (Glasgow coma scale 6 points), his body temperature was 41.2°C , heart rate 160/min, and blood pressure 110/43 mmHg. Respiration rate was 40/min, and oxygen saturation 96% with O_2 9L/min mask inhalation. Examination of the chest and abdomen was unremarkable.

Blood examination revealed leukocytosis, and elevated levels of C-reactive protein (1.37 mg/dl: normal value <0.30 mg/dl), aspartate transaminase (82 U/L: normal range <40 U/L), lactate dehydrogenase (444 U/L: normal range <245 U/L), γ -glutamyl transferase (109 U/L: normal range <75 U/L), blood sugar (356 mg/dl: normal range 70–109 mg/dl), and uric acid (12.4 mg/d: normal range <7.0 mg/d). Renal impairment (blood urea nitrogen/creatinine = 38/1.88 mg/dl: normal range $<22/1.04$ mg/dl), and hyponatremia (126 mEq/L: normal range 135–147 mEq/L) were also noted.

A 12-lead electrocardiogram revealed regular wide QRS tachycardia of a rate of 160/min with superior axis and atypical complete right bundle branch block (CRBBB) pattern (Fig. 1). A chest radiograph showed cardiomegaly without pulmonary edema. Echocardiography showed diffuse severe left ventricular hypokinesis.

* Corresponding author at: Takeda General Hospital, 28-1 Ishidamoriminami, Fushimi-ku, Kyoto 601-1495, Japan. Tel.: +81 75 572 6331.

E-mail address: c-watanabe@takedahp.or.jp (C. Watanabe).



Fig. 1. 12-lead electrocardiogram at initial presentation. It showed regular wide QRS tachycardia with superior axis and atypical complete right bundle branch block morphology.

As a tentative diagnosis of heat stroke and wide QRS tachycardia due to unknown cause was made, rapid infusion of 2 L cold normal saline and surface cooling with ice pack were started. Intravenous administration of lidocaine (50 mg) and magnesium sulfate (2.46 g) resulted in repetitive transient termination and relapse of tachycardia. Thus, the patient was intubated with sedation by intravenous propofol, which led to gradual slowing of the wide QRS complex tachycardia, and transition to atrial fibrillation (AF). A 12-lead electrocardiogram after wide QRS tachycardia termination showed AF with left anterior hemiblock and CRBBB. These findings, except AF rhythm and slightly different morphology in precordial leads, had been also noted 17 months prior to admission. Acute myocardial infarction was excluded because neither definite ST elevation nor local asynergy in echocardiogram was recognized.

His body temperature returned to 37.5 °C in 4 h by vigorous cooling. Although blood pressure was preserved (100–110 mmHg) during wide QRS tachycardia, he fell into shock (blood pressure 50–60 mmHg) after conversion to AF. Continuous infusion of norepinephrine (0.3 µg/kg/min) to maintain blood pressure above 90 mmHg and heparin to prevent thromboembolism were initiated.

On the next day, blood examination revealed marked increase of transaminase [aspartate transaminase (7521 U/L), alanine transaminase (2636 U/L), and lactate dehydrogenase (5321 U/L)], indicating rhabdomyolysis and/or liver shock. Myoglobinuria (37,600 mg/dl) and elevated creatine phosphokinase (CPK) (4154 U/L: normal range <250 U/L) indicating rhabdomyolysis due to heat stroke and positive troponin-T (1.65 ng/ml: normal range <0.012 ng/ml) indicating myocardial damage were also noted. Thrombocytopenia ($4.4 \times 10^4/\mu\text{l}$) accompanied by prolongation of prothrombin time (56.2%: normal range 66.0–127.6%), activated partial thromboplastin time (46.8 s: normal range 25.1–39.8 s), and increased fibrin degradation product (21.3 µg/ml:

normal range <5.0 µg/ml) suggested disseminated intravascular coagulation (DIC).

Electrocardiogram on the following day showed giant negative T wave in II, III, aVF, and V2–6 (Fig. 2). Re-examination of echocardiography demonstrated akinesis or severe hypokinesis at mid to apical wall with preserved basal wall motion (Fig. 3). Temporal change of electrocardiogram and echocardiographic findings suggested stress-induced (takotsubo) cardiomyopathy. He was given carvedilol (2.5 mg/day) and enalapril (2.5 mg/day) orally.

There was no recurrence of wide QRS tachycardia and hemodynamics were stable after tapering and cessation of noradrenaline infusion. On the 4th day after admission, he recovered consciousness without evident neurologic deficit and was extubated. One delivery of electrical shock with 100 J for AF on the 5th day restored sinus rhythm, with no recurrence of AF thereafter. On the 7th day, apical wall motion was still hypokinetic but clearly improved. On the 14th day, echocardiogram showed complete recovery of left ventricular wall motion with ejection fraction of 66%. Liver enzyme, CPK, coagulation disorders, and renal function were gradually normalized. Multi-detector computed tomography (MDCT) showed normal coronary tree. His condition continued to improve and he was transferred to another hospital to continue rehabilitation.

Discussion

Heat stroke is an acute thermoregulatory failure due to exposure to high temperatures. Excessive heat denatures proteins, destabilizes phospholipids and lipoproteins, and liquefies membrane lipids, leading to diffuse tissue derangement. Complications of heat stroke include acute renal failure, rhabdomyolysis, DIC, acute respiratory distress syndrome, and brain damage. MOF and death will ensue in severe heat stroke.

Download English Version:

<https://daneshyari.com/en/article/5984445>

Download Persian Version:

<https://daneshyari.com/article/5984445>

[Daneshyari.com](https://daneshyari.com)