



Short communication

Automated assessment of early hypoxic brain edema in non-enhanced CT predicts outcome in patients after cardiac arrest[☆]

Uta Hanning^{a,b,1}, Peter Bernhard Sporns^{a,1}, Pia Lebiedz^c, Thomas Niederstadt^a, Tarek Zoubi^a, Rene Schmidt^d, Stefan Knecht^e, Walter Heindel^a, André Kemmling^{f,g,*}

^a Department of Clinical Radiology, University Hospital of Muenster, Germany

^b Institute of Epidemiology and Social Medicine, University of Muenster, Germany

^c Department of Cardiovascular Medicine, University of Muenster, Germany

^d Institute of Biostatistics and Clinical Research, University of Muenster, Germany

^e Department of Neurology, Mauritius Hospital Meerbusch and Clinical Neuroscience, University Duesseldorf, Germany

^f Institute of Neuroradiology, University Hospital of Luebeck, Germany

^g Department of Diagnostic and Interventional Neuroradiology, University Medical Center Hamburg-Eppendorf, Germany

ARTICLE INFO

Article history:

Received 18 October 2015

Received in revised form 26 February 2016

Accepted 13 March 2016

Keywords:

CT

Cardiac arrest

Coma

Outcome

ABSTRACT

Introduction: Early prediction of potential neurological recovery in patients after cardiac arrest is challenging. Recent studies suggest that the densitometric gray-white matter ratio (GWR) determined from cranial computed tomography (CT) scans may be a reliable predictor of poor outcome. We evaluated an automated, rater independent method to determine GWR in CT as an early objective imaging predictor of clinical outcome.

Methods: We analyzed imaging data of 84 patients after cardiac arrest that underwent noncontrast CT within 24 h after arrest. To determine GWR in CT we applied two methods using a recently published automated probabilistic gray-white matter segmentation algorithm (GWR.aut) and conventional manual measurements within gray-white regions of interest (GWR.man). Neurological outcome was graded by the cerebral performance category (CPC). As part of standard routine CPC was assessed by the treating physician in the intensive care unit at admission and at discharge to normal ward. The performance of GWR measures (automated and manual) to predict the binary clinical endpoints of poor (CPC3–5) and good outcome (CPC1–2) was assessed by ROC analysis with increasing discrimination thresholds. Results of GWR.aut were compared to GWR.man of two raters.

Results: Of 84 patients, 55 (65%) showed a poor outcome. ROC curve analysis revealed reliable outcome prediction of GWR.aut (AUC 0.860) and GWR.man (AUC 0.707 and 0.699, respectively). Predictive power of GWR.aut was higher than GWR.man by each rater ($p=0.019$ and $p=0.021$, respectively) at an optimal cut-off of 1.084 to predict poor outcome (optimal criterion with 92.7% sensitivity, 72.4% specificity). Interrater reliability of GWR.man by intra-class correlation coefficient (ICC) was moderate (0.551).

Conclusion: Automated quantification of GWR in CT may be used as an objective observer-independent imaging marker for outcome in patients after cardiac arrest.

© 2016 Elsevier Ireland Ltd. All rights reserved.

Introduction

Recent advances in emergency medicine and resuscitation management such as mild therapeutic hypothermia have increased the

number of survivors of cardiac arrest.^{1,2} However cognitive and neurologic outcomes are still poor in patients following resuscitation, especially in initially comatose patients.³ Early assessment of the potential of neurological recovery in comatose cardiac arrest survivors remains challenging.^{4,5} As the number of survivors of cardiac arrest increase and more effective therapies become available the need to identify those patients who remain comatose with likely good neurologic outcome becomes essential.^{4,6} Recent studies suggest that a reduced GWR in cranial computed tomography (CT) scans indicating hypoxic brain edema may be an early predictor of poor outcome in cardiac arrest patients.^{4,7–9} However, in these studies GWR measurements were not objective and only

[☆] A Spanish translated version of the summary of this article appears as Appendix in the final online version at <http://dx.doi.org/10.1016/j.resuscitation.2016.03.018>.

* Corresponding author at: Institut für Neuroradiologie, Universitätsklinikum Schleswig-Holstein, Campus Luebeck, Ratzeburger Allee 160 (Zentralklinikum), 23538 Luebeck, Germany.

E-mail address: andre.kemmling@uksh.de (A. Kemmling).

¹ Both authors contributed equally.

assessed visually or by manual placement of regions of interest (ROI) in gray and white matter.

We hypothesise that an observer independent method to determine in CT may be superior to the manual ratings limited by rater variability. We therefore present an automated method to determine GWR in CT as an early objective imaging predictor of clinical outcome and compared the results to conventional manual measurements.

Methods

Study population

Between 1-2011 and 12-2014 we consecutively included 87 data sets acquired from comatose cardiac arrest patients who underwent non-contrast spiral CT of sufficient quality within 24 h after arrest. Electronic patient records were anonymized prior to analysis and our study design was entirely retrospective. In this setting, the need for patient consent was waived by our institutional review board. In our department, routine neurological evaluation of patients after cardiac arrest consists of laboratory, electrophysiological and clinical examinations including the Cerebral Performance Category score (CPC).¹⁰ The CPC was assessed by the treating physician in the intensive care unit (ICU) at admission and at the time of discharge to normal ward. Neurological evaluation included pupillary light reflex (PLR), corneal reflex (CR) (day 3) and Glasgow coma scale (GCS) (days 1 and 3). Exclusion criteria were haemorrhage or ischemic infarction in CT.

GWR determination

For automated GWR determination we employed a recently published brain segmentation algorithm to decompose the CT density of brain into tissue-specific components.¹¹ In brief, the algorithm uses normalized probabilistic maps of white matter (WM) and gray matter (GM) in standard space derived from partial tissue segmentations of 600 unsuspicious brain MRIs (3.0T, T1-3D-Turbo-Field-Echo). The voxel-specific partial tissue component

(tissue probability) in CT space was obtained by precise non-linear deformation of tissue probability maps in standard space to the individual CT space using a custom CT reference image.¹¹ The tissue-specific density within GM- and WM-space was then determined by the mean of all voxel densities weighted by probabilistic GM- and WM-content, respectively (Eq. (1)).¹¹ All CT image operations were performed using the FMRIB Software Library v5.0, Oxford, UK.

$$\frac{\sum(HU_{xyz} \times P_{xyz}[tissue])}{\sum(P_{xyz}[tissue])} \quad (1)$$

(HU_{xyz} = density of voxel xyz in Hounsfield units; P_{xyz} = partial tissue component at voxel xyz for GM or WM, respectively).

GWR.aut was then calculated as the ratio between the mean CT density within GM- and WM-space (Fig. 1). Manual ratings by ROI placement were performed independently by two readers who were blinded to outcome. ROI measurements were performed under a standardized protocol and included bilateral manual placement of circular ROIs (area = 0.1 cm²) in the putamen and the posterior limb of internal capsule.⁸

Statistical analysis

Univariable distribution of metric variables is described by median and interquartile range. For categorical data, absolute and relative frequencies are given.

Patients with good (CPC of 1–2) versus poor (CPC of 3–5) outcome were compared by Mann–Whitney *U* test for metric outcome variables and by chi-square test for categorical outcome variables. The performance of GWR measures (automated and manual) to predict poor and good outcome was assessed by ROC analysis with increasing discrimination thresholds. Pairwise ROC comparisons between GWR.aut and both GWR.man were performed by assessing differences in the areas under the empirical ROC curves.¹² Interrater reliability of GWR.man was quantified using intra-class correlation coefficient (ICC).

No adjustment for multiple testing was performed and “significance” refers to local statistical significance defined as a local,




Fig. 1. Tissue specific decomposition of a native CT to determine GWR in global cerebral hypoxia. Automated probabilistic assessment of gray matter (middle column) and white matter (right column) in computed tomography (left column): GWR's in two patients resuscitated after cardiac arrest with good outcome versus poor outcome (lower row).

Download English Version:

<https://daneshyari.com/en/article/5997256>

Download Persian Version:

<https://daneshyari.com/article/5997256>

[Daneshyari.com](https://daneshyari.com)