

Conditional Sox9 ablation improves locomotor recovery after spinal cord injury by increasing reactive sprouting

William M. McKillop, Elisa M. York, Luc Rubinger, Tony Liu, Natalie Maria Ossowski, Kathy Xu, Todd Hryciw, Arthur Brown

PII: S0014-4886(16)30146-7
DOI: doi: 10.1016/j.expneurol.2016.05.028
Reference: YEXNR 12305

To appear in: *Experimental Neurology*

Received date: 19 January 2016
Revised date: 18 May 2016
Accepted date: 21 May 2016

Please cite this article as: McKillop, William M., York, Elisa M., Rubinger, Luc, Liu, Tony, Ossowski, Natalie Maria, Xu, Kathy, Hryciw, Todd, Brown, Arthur, Conditional Sox9 ablation improves locomotor recovery after spinal cord injury by increasing reactive sprouting, *Experimental Neurology* (2016), doi: 10.1016/j.expneurol.2016.05.028

Conditional Sox9 ablation improves locomotor recovery after spinal cord injury by increasing reactive sprouting

William M. McKillop[a,b], Elisa M. York[a], Luc Rubinger[a], Tony Liu[a], Natalie Maria Ossowski, Kathy Xu[a], Todd Hryciw[a] and Arthur Brown[a,b]

Corresponding Author: Dr. Arthur Brown
Robarts Research Institute, Schulich School of Medicine,
University of Western Ontario
100 Perth Drive, London, Ontario, Canada, N6A 5K8
Email: abrown@robarts.ca
Telephone: 519-663-3776 ext. 24308

[a] Robarts Research Institute, University of Western Ontario, London, Canada

[b] Department of Anatomy and Cell Biology, University of Western Ontario, London, Canada

Keywords: SOX9, spinal cord injury, neuroplasticity, CSPG, reactive sprouting, perineuronal nets