

Accepted Manuscript

Decompressive Craniectomy in Conjunction with Evacuation of Intracranial Hemorrhagic Lesions is Associated with Worse Outcomes in Elderly Patients with Traumatic Brain Injury: A Propensity Score Analysis

Takahiro Kinoshita, MD, Kazuhisa Yoshiya, MD, PhD, Yasunori Fujimoto, MD, PhD, Ryuichiro Kajikawa, MD, Takeyuki Kiguchi, MD, PhD, Masahiko Hara, MD, PhD, Akatsuki Wakayama, MD, PhD, Toshiki Yoshimine, MD, PhD

PII: S1878-8750(16)00159-5

DOI: [10.1016/j.wneu.2016.01.071](https://doi.org/10.1016/j.wneu.2016.01.071)

Reference: WNEU 3675

To appear in: *World Neurosurgery*


Received Date: 7 November 2015

Revised Date: 17 January 2016

Accepted Date: 19 January 2016

Please cite this article as: Kinoshita T, Yoshiya K, Fujimoto Y, Kajikawa R, Kiguchi T, Hara M, Wakayama A, Yoshimine T, Decompressive Craniectomy in Conjunction with Evacuation of Intracranial Hemorrhagic Lesions is Associated with Worse Outcomes in Elderly Patients with Traumatic Brain Injury: A Propensity Score Analysis, *World Neurosurgery* (2016), doi: 10.1016/j.wneu.2016.01.071.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Original Research Article

Decompressive Craniectomy in Conjunction with Evacuation of Intracranial Hemorrhagic Lesions is Associated with Worse Outcomes in Elderly Patients with Traumatic Brain Injury: A Propensity Score Analysis

Running Head Title: Craniotomy vs DC in elderly patients with TBI

Takahiro Kinoshita, MD¹⁾²⁾, Kazuhisa Yoshiya, MD, PhD³⁾, Yasunori Fujimoto, MD, PhD¹⁾,
Ryuichiro Kajikawa, MD¹⁾, Takeyuki Kiguchi, MD, PhD²⁾, Masahiko Hara, MD, PhD⁴⁾⁵⁾,
Akatsuki Wakayama, MD, PhD¹⁾, Toshiki Yoshimine, MD, PhD⁶⁾

1: Department of Neurosurgery, Osaka Neurological Institute, Toyonaka, Japan

2: Department of Emergency and Critical Care, Osaka General Medical Center, Osaka, Japan

3: Department of Traumatology and Acute Critical Medicine, Osaka University Graduate School of Medicine, Suita, Japan

4: Department of Clinical Epidemiology and Biostatistics, Osaka University Graduate School of Medicine, Suita, Japan

5: Department of Medical Innovation, Osaka University Hospital, Suita, Japan

Download English Version:

<https://daneshyari.com/en/article/6043714>

Download Persian Version:

<https://daneshyari.com/article/6043714>

[Daneshyari.com](https://daneshyari.com)