

# Accepted Manuscript

Endoscopic treatment of temporal arachnoid cysts in 34 patients

Tim Couvreur, MD, Giorgio Hallaert, MD, Tatjana Van Der Heggen, MD, Edward Baert, MD, Frank Dewaele, MD, Jean-Pierre Kalala Okito, MD PhD, Dimitri Vanhauwaert, MD, Marc Deruytter, MD, Dirk Van Roost, MD PhD, Jacques Caemaert, MD PhD



PII: S1878-8750(15)00486-6

DOI: [10.1016/j.wneu.2015.04.053](https://doi.org/10.1016/j.wneu.2015.04.053)

Reference: WNEU 2882

To appear in: *World Neurosurgery*

Received Date: 31 December 2014

Revised Date: 22 April 2015

Accepted Date: 22 April 2015

Please cite this article as: Couvreur T, Hallaert G, Van Der Heggen T, Baert E, Dewaele F, Kalala Okito J-P, Vanhauwaert D, Deruytter M, Van Roost D, Caemaert J, Endoscopic treatment of temporal arachnoid cysts in 34 patients, *World Neurosurgery* (2015), doi: 10.1016/j.wneu.2015.04.053.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## Highlights:

- 1: We present the largest single centre study on endoscopic TAC fenestration
- 2: First description of the use of a YAG-cutting laser for fenestration
- 3: Intracranial hypertension is the best indication for surgery
- 4: Specific attention to epilepsy originating from the ipsilateral temporal lobe

Download English Version:

<https://daneshyari.com/en/article/6045394>

Download Persian Version:

<https://daneshyari.com/article/6045394>

[Daneshyari.com](https://daneshyari.com)