

---

## Maxillary sinus floor augmentation in patients with maxillary sinus pseudocyst: case report

Nukhet Celebi, DDS, PhD,<sup>a</sup> Zeynep Burcin Gonen, DDS,<sup>b</sup> Erdem Kilic, DDS, PhD,<sup>a</sup> Osman Etoz, DDS, PhD,<sup>a</sup> and Alper Alkan, DDS, PhD,<sup>c</sup> Kayseri, Turkey  
ERCIYES UNIVERSITY

The maxillary sinus floor elevation procedure has gained popularity with predictable results, and is a safe, acceptable technique for bone augmentation, providing a base for dental implant treatment. Faint radiopaque lesions at the base of the maxillary sinus are frequent diagnoses on radiographs and must be identified during dental implant planning. Pseudocysts classically appear hemispheric, homogeneously opaque, and well delineated in panoramic and periapical radiographs. The great majority of these lesions are asymptomatic and do not require surgical treatment. In this case report, we present 4 patients who had a maxillary sinus floor elevation procedure using either crestal or lateral approaches in the presence of antral pseudocysts. No complications were encountered during follow-up periods in these patients and all implants are functioning successfully. (*Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2011;112:e97-e102)

Maxillary sinus floor elevation is a safe and predictable surgical procedure to vertically augment the residual alveolar bone,<sup>1</sup> and allows the placement of dental implants with adequate length in the edentulous posterior maxilla.<sup>2</sup> Currently, a sinus floor elevation procedure can be performed with either a lateral or a crestal approach.<sup>3</sup> The lateral approach was originally described by Tatum and subsequently published by Boyne and James.<sup>4</sup> The classic procedure for this augmentation entails the preparation of a trap door to elevate the Schneiderian membrane in the lateral sinus wall. The space created beneath the lifted sinus membrane is then grafted with different fillers consisting of autogenous bone, bone substitutes, or a mixture of these materials.<sup>4-7</sup> More recently, Summers<sup>1</sup> suggested the crestal approach for sinus floor elevation with subsequent placement of implants, using a set of osteotomes of varying diameters to prepare the implant site. The crestal approach is proposed when there is at least 5 to 6 mm of residual alveolar bone height.<sup>8</sup>

Regardless of the method used, risk assessment must be considered during the preoperative surgical planning

to reduce intra- and postsurgical sinus complications.<sup>9</sup> Radiopaque lesions, such as mucoceles, retention cysts, and pseudocysts at the floor of the maxillary sinus may present obstacles during sinus elevation.<sup>9</sup> Sinus cysts present different clinicopathologic features and behavior.<sup>10</sup> Mucoceles are extravasations of mucous into the surrounding soft tissues following trauma or obstruction of salivary flow. The expansive nature of the lesion can cause bone resorption. Retention cysts are thought to arise from blockage of an antral seromucous gland, resulting in a ductal epithelium-lined cystic structure filled with mucin. Pseudocysts are inflammatory in origin and result from fluid accumulation within the sinus membrane. This lesion lacks an epithelial lining.<sup>10</sup> Antral pseudocysts have recently been reported not to be a contraindication for sinus floor augmentation procedures with lateral approach.<sup>11,12</sup>

In this case report, we present 4 patients who had a maxillary sinus floor elevation procedure using either crestal or lateral approaches in the presence of antral pseudocysts.

### MATERIAL AND METHODS

Four patients (2 men, 2 women) with an edentulous atrophic maxilla, median age 42.7 (range, 37-50), underwent maxillary sinus floor bone augmentation procedures (Table I). All of the patients were referred to Erciyes University, Oral and Maxillofacial Surgery Hospital, for implant-supported prosthesis in a partially edentulous maxilla. The panoramic radiographic examination revealed insufficient alveolar bone height for the placement of implants with adequate length on the posterior maxilla, and a dome-shaped, faintly radiopaque lesion, compatible with an antral pseudocyst,

This study was presented as an oral presentation at the 5th ACBID International Conference, May 25-29, 2011.

<sup>a</sup>Assistant Professor, Department of Oral and Maxillofacial Surgery, Faculty of Dentistry, Erciyes University, Kayseri, Turkey.

<sup>b</sup>Resident, Department of Oral and Maxillofacial Surgery, Faculty of Dentistry, Erciyes University, Kayseri, Turkey.

<sup>c</sup>Professor, Department of Oral and Maxillofacial Surgery, Faculty of Dentistry, Erciyes University, Kayseri, Turkey.

Received for publication May 3, 2011; accepted for publication Jun 2, 2011.

1079-2104/\$ - see front matter

© 2011 Mosby, Inc. All rights reserved.

doi:10.1016/j.tripleo.2011.06.001

**Table I.** Patients and procedures

Case	Gender	Age	Radiographic examination	Diameter of pseudocyst (width × height)	Residual bone height	Sinus augmentation procedure	Implants
1	Male	38	OPG + CT	25 × 18 mm	4 mm	Crestal approach	4.0 × 11.5 mm 3I
2	Female	46	OPG	13 × 10 mm	5 mm	Crestal approach	4.0 × 11.5 mm 3I
3	Female	37	OPG	18 × 16 mm	2 mm	Lateral approach	4.1 × 12.0 mm ITI
4	Male	50	OPG + CT	16 × 13 mm	2 mm	Lateral approach	4.1 × 12.0 mm ITI

OPG, orthopantomograph; CT, computerized tomography.

located in the related maxillary sinuses in all of the patients. Antral pseudocyst was confirmed with either computerized tomographic (CT) examination and/or panoramic radiographic examination (Figs. 1-4). The patients' medical histories were uneventful except for case 1, who had been smoking for 20 years.

### Crestal-approach patients

In the first 2 cases, 3I implants (3I Dental Implant System, BIOMET, Inc, Warsaw, IN) were inserted with the crestal approach using osteotomes of gradually increasing diameters in the right posterior maxillas of both patients. No graft materials were used for sinus lifting. Amoxicillin/clavulanic acid, 625 mg twice daily for 10 days, chlorhexidine gluconate-benzylamine HCl mouth rinse 3 times daily for 7 days, and flurbiprofen 100 mg twice daily for 7 days, and a nasal decongestant spray regimen were prescribed to all patients after the operations. The patients had no discomfort and no complaints postoperatively. Final prosthetic restorations were reconstructed at least 4 months after the surgery.

### Lateral-approach patients

In these 2 cases, a 2-stage procedure was performed using the lateral approach. To augment the sinus floor, 1 g of bovine bone xenograft (Bio Oss, Geistlich Sons, Ltd., Wolhusen, Switzerland) was used. The lateral sinus window was covered with platelet-rich fibrin membrane in case 3, and with 20 × 20-mm collagen membrane (OsteoBiol, Tecness Dental, Torino, Italy) in case 4. The same prescriptions as with the first 2 cases were delivered. ITI implants (ITI, Straumann, Basel, Switzerland) were inserted 7 months after the first surgery.

No problems occurred related either to the cyst or to the implants for 6 to 8 months postoperatively for all patients. All implants were osseointegrated successfully in all patients (Figs. 5-8).

## DISCUSSION

Sinus augmentation has a high percentage of success, but presents a number of complications, such as membrane perforation, obstruction of the maxillary ostium,



Fig. 1. Right antral pseudocyst was confirmed with CT examination in case 1.

hemorrhage, acute or chronic infection, graft loss, and dental implant failure.<sup>9</sup> Proper maintenance of normal physiology is necessary for decreasing the incidence of complications. The ostium, located 25 to 35 mm above the antral floor, connects the maxillary sinus to the nasal cavity.<sup>13</sup> Ostium in patients with a pseudocyst may be blocked by obliteration of the sinus cavity by overfilling with bone graft in sinus augmentation procedures.

Faint radiopaque lesions at the base of the maxillary sinus are frequently diagnosed on radiographs<sup>9</sup> and must be identified during dental implant planning. Pseudocysts classically appear hemispheric, homogeneously opaque, and well delineated in panoramic and periapical radiographs. They usually demonstrate an attachment to the floor of the maxillary sinus. The great majority of these lesions are asymptomatic and do not require surgical treatment.<sup>14</sup>

Although the literature indicates that sinus augmentation procedures are predictable and safe,<sup>9</sup> the presence of maxillary antral pseudocysts and retention cysts may present an obstacle during sinus elevation and might result in future complications and potential fail-

Download English Version:

<https://daneshyari.com/en/article/6059417>

Download Persian Version:

<https://daneshyari.com/article/6059417>

[Daneshyari.com](https://daneshyari.com)