Accepted Manuscript

The aldo-keto reductase AKR1B10 is upregulated in keloid epidermis, implicating retinoic acid pathway dysregulation in the pathogenesis of keloid disease

Natalie Jumper, Tom Hodgkinson, Guyan Arscott, Yaron Har-Shai, Ralf Paus, Ardeshir Bayat

PII: S0022-202X(16)30985-X

DOI: 10.1016/j.jid.2016.03.022

Reference: JID 272

To appear in: *The Journal of Investigative Dermatology*

Received Date: 14 December 2015

Revised Date: 9 February 2016

Accepted Date: 7 March 2016

Please cite this article as: Jumper N, Hodgkinson T, Arscott G, Har-Shai Y, Paus R, Bayat A, The aldo-keto reductase AKR1B10 is upregulated in keloid epidermis, implicating retinoic acid pathway dysregulation in the pathogenesis of keloid disease, *The Journal of Investigative Dermatology* (2016), doi: 10.1016/j.jid.2016.03.022.

The aldo-keto reductase AKR1B10 is upregulated in keloid epidermis, implicating retinoic acid pathway dysregulation in the pathogenesis of keloid disease

Natalie Jumper[1], Tom Hodgkinson[1], Guyan Arscott[5], Yaron Har-Shai[4], Ralf Paus[2, 3], Ardeshir Bayat[1, 2]

[1]*Plastic and Reconstructive Surgery Research, Manchester Institute of Biotechnology, University of Manchester, 131 Princess St, M1 7DN UK*

[2]*Centre for Dermatology Research, Institute of Inflammation and Repair, University of Manchester, Manchester, UK*

[3]*Department of Dermatology, University of Münster, D-48149, Münster, Germany*

[4]*Plastic Surgery Unit, Carmel Medical Center, Haifa, Israel*.

[5]*Department of Plastic and Reconstructive Surgery, University of West Indies, Kingston, Jamaica*

Correspondence to: Dr Ardeshir Bayat, *Plastic & Reconstructive Surgery Research*, *Institute of Inflammation & Repair*, *University of Manchester*, *Stopford Building*, *Manchester M13 9PT*.

Phone: *0161 306 0607 Email: Ardeshir.Bayat@manchester.ac.uk*