

Wounds with complicated shapes tend to develop infection during negative pressure wound therapy



Masaki Fujioka^{a,b,*}, Kenji Hayashida^c, Chikako Senju^c

^a Department of Plastic and Reconstructive Surgery, Nagasaki University, Nagasaki, Japan

^b Department of Plastic and Reconstructive Surgery, Clinical Research Center, National Hospital Organization Nagasaki Medical Center, Nagasaki, Japan

^c Department of Plastic and Reconstructive Surgery, National Hospital Organization Nagasaki Medical Center, Nagasaki, Japan

ARTICLE INFO

Article history:

Received 12 August 2013

Accepted 25 October 2013

Available online 19 November 2013

Keywords:

Complicated wounds

Wound infection

Negative pressure wound therapy

Vacuum-assisted closure therapy

Wound geometry

ABSTRACT

Introduction: While negative pressure wound therapy (NPWT) has been shown to be useful, we felt that patients with wounds of complicated shapes were likely to develop infection during performing NPWT. We conducted an investigation to determine the factors of wound shape responsible for the occurrence of infection.

Materials and methods: A total of 55 patients with wounds were treated using NPWT in our unit in 2011. Eight whose wounds formed a pocket, 7 whose wounds were deep, and 40 whose wounds did not come under the above 2 types were eligible for this retrospective study.

Results: Fifteen patients (27.3%) with NPWT showed a relapse of local infection. Six of the 8 patients (75.0%) in the wound with pocket group, 5 of the 7 (71.4%) in the deep wound group, and 4 of the 40 (10.0%) in the other wounds developed infection. The wound infection development ratio of the wound with pocket and deep wound groups was significantly higher than that of the other wound group.

Conclusion: Wounds with complicated shapes are more likely to develop infectious complications during the management of NPWT. More careful observation is required when negative pressure therapy is used for wounds with a complicated shape.

© 2013 Elsevier GmbH. All rights reserved.

1. Introduction

Negative pressure wound therapy (NPWT) of infected wounds has recently gained popularity among various surgical specialties [1–3]. This system is based on the application of negative pressure by controlled suction to the wound surface. The effectiveness of the NPWT for microcirculation and the promotion of granulation tissue proliferation owing to removing excessive exudates, increasing blood flow, and decreasing bacterial colonization has been verified. Thus, it has allowed uncomplicated wounds to heal quickly [4]. However, it sometimes leads to local wound infection, including: erythema, swelling, increased pain, exudates or pus, and fever, which can cause long-term distress for the patient, increase the hospitalization time, and, consequently, decrease the quality of life. In our experience, we have seen an increased tendency of wounds with complicated shapes to develop infection during NPWT.

Abbreviation: NPWT, negative pressure wound therapy.

* Corresponding author at: Department of Plastic and Reconstructive Surgery, National Hospital Organization Nagasaki Medical Center, 1001-1 Kubara 2, Ohmura City 856-8562, Japan. Tel.: +81 0957 52 3121; fax: +81 0957 54 0292.

E-mail address: mfujioka@nmc.hosp.go.jp (M. Fujioka).

The present study investigates this by comparing patients who underwent NPWT with deep wounds, wounds with large pockets, and shallow wounds without pockets.

2. Patients and methods

NPWT has been employed in our department since 2011 as a device to bridge the period between debridement and definite surgical closure in full-thickness wounds. A total of 575 patients with wounds (acute wounds: 345, chronic wounds: 230) were treated in the Department of Plastic and Reconstructive Surgery, National Organization Nagasaki Medical Center, in 2011. Of these patients, 55 underwent negative pressure wound therapy using the Vacume-Assisted Closure System (V.A.C.ATS[®], KCI Inc., San Antonio, TX, USA). Wound diagnoses of patients who received NPWT are shown in Fig. 1. All participants received surgical debridement, as well as nutrition and hemodynamic support. After debridement, cleansing and wet-to-dry dressing or continuous irrigation were performed for several days. After recognizing symptoms of infection disappeared, subatmospheric pressure (125 mmHg below ambient) was applied and transmitted to the wound continuously using a pump. We used black foam. In cases wounds formed a pocket with over-hanging skin, the form was not




Fig. 1. Wound diagnosis in patients who received NPWT (N = 55).

inserted to the pocket, but laid over the area of ulcer to stick the inner wall of pocket firmly each other.

The dressing foam was usually changed every 48 h; however, this varied depending on the presence of infection. As no wounds showed inflammatory signs at the initiation of NPWT, no patients were administered antibiotics during NPWT use. The diagnosis of wound infection was based on the clinical signs and symptoms of the patient, including: erythema or skin discoloration, edema, warmth, induration, increased pain, purulent wound exudate, elevated temperature, and elevated white blood cell count.

Of these patients, 8 whose wounds formed a pocket with more than 1 cm of over-hanging skin (wound with pocket group), 7 whose wounds were deep, extending to the bone or penetrating into the muscle (deep wound group), and 40 whose shallow wounds were without a pockets (other group) were eligible for this retrospective study (Fig. 2). In this study, wound with pocket group was defined as having a more than 1 cm of over-hanging skin to differentiate wounds with pocket from those with thick rolled wound margin, but without undermining pocket. All information was obtained from patients' medical records, examination, and interview at the first examination.

3. Results

Fifteen of the 55 patients (27.3%) with NPWT showed a relapse of local infection. Patients developing infection ranged in age from




Fig. 2. Wound shape in patients who received NPWT.

Table 1
The cause of wound in patients with and without wound infection.

	Pressure ulcer	Infection	Trauma	Ischemia	Chronic ulcer
Infection (15)	6	6	1	1	1
Non-infection (40)	9	14	11	3	3

Table 2
The location of wound in patients with and without wound infection.

	Buttock	Extrimities	Trunk
Infection (15)	7	6	2
Non-infection (40)	7	26	6

Table 3
The complications which influence on development of infection in patients with and without wound infection.

	Diabetes mellitus	Renal failure	Collagen disease	Cancer	Steroid usage
Infection (15)	8	1	1	3	1
Non-infection (40)	17	2	4	8	4

31 to 84 years (mean age, 57.5 years), and patients without infection ranged in age from 8 to 95 years (mean age, 51.5 years) (no significant difference, Wilcoxon rank sum test). The etiology of wounds in patients with and without wound infection is shown in Table 1. There was no significant difference between the groups for each cause of wound ($p > 0.05$, Chi-square test). The locations of wounds in patients with and without wound infection are shown in Table 2. There was no significant difference between the groups in each wound location ($p > 0.05$, Chi-square test). Complications which may influence the development of infection, such as diabetes mellitus, renal failure, collagen disease, cancer, and steroid usage, in patients with and without wound infection are shown in Table 3. There was no significant difference between the groups for each complication ($p > 0.05$, Chi-square test).

Six of the 8 patients (75.0%) in the wound with pocket group, 5 of the 7 (71.4%) in the deep wound group, and 4 of the 40 (10.0%) in the other group developed infection. The patients' sex, age, characters of wounds, location, and the interval between the start of NPWT and the development of infection are shown in Table 4. The mean number of negative pressure wound therapy treatment days was 10 (range: 1–19). The wound infection development ratio

Table 4
Cases of wound infection development during VAC therapy.

Case	Shape of wound	Sex	Age	Wound	Location	NPWT duration (days)
1	Pocket	M	75	Pressure ulcer	Sacrum	15
2	Pocket	M	31	Pressure ulcer	Ischial tuberosity	19
3	Pocket	M	57	Pressure ulcer	Ischial tuberosity	14
4	Pocket	F	78	Sacral pressure ulcer	Sacrum	11
5	Pocket	M	43	Abscess	Iliopsoas	8
6	Pocket	M	54	Pressure ulcer	Sacrum	12
7	Deep	F	40	Osteomyelitis	Trochanter	10
8	Deep	F	72	Post-surgical wound infection	Abdomen	1
9	Deep	F	74	Osteomyelitis	Foot	11
10	Deep	M	61	Intra-muscular abscess	Thigh	10
11	Deep	F	54	Gas gangrene	Leg	3
12	Others	M	8	Laceration	Knee	14
13	Others	F	88	Ischemic necrosis	Toe	11
14	Others	M	70	Chronic ulcer	Leg	8
15	Others	M	65	Pressure ulcer	Heel	12

Download English Version:

<https://daneshyari.com/en/article/6078530>

Download Persian Version:

<https://daneshyari.com/article/6078530>

[Daneshyari.com](https://daneshyari.com)