



Zone 2 sacral fractures managed with partially-threaded screws result in low risk of neurologic injury



Amir Herman^{a,b,*}, Emily Keener^a, Candice Dubose^a, Jason A. Lowe^a

^aThe Orthopaedic Trauma Unit, Division of Orthopaedics, University of Alabama at Birmingham, United States

^bTalpiot Medical Leadership Program, Sheba Medical Centre, Israel

ARTICLE INFO

Article history:

Accepted 9 April 2016

Keywords:

Trans-sacral screws

Sacroiliac screws

Partially threaded screws

ABSTRACT

Background: Zone 2 sacral fractures account for 34% of sacral fractures with reported neurological deficit in 21–28% of patients. The purpose of this study was to examine the risk factors for neurological injury in zone 2 sacral fractures. The authors hypothesized that partially thread iliosacral screws did not increase incidence of neurologic injury.

Methods: A retrospective review of consecutive patients admitted to a level 1 trauma center with zone 2 sacral fractures requiring surgery from September 2010 to September 2014 was performed. Patients were excluded if no neurological exam was available after surgery. Fractures were classified according to Denis and presence/absence of comminution through the neural foramen was noted. Fixation schema was recorded (sacral screws or open reduction and internal fixation with posterior tension plate). Any change in post-operative neurological exam was documented as well as exam at last clinic encounter. **Results:** 90 patients met inclusion criteria, with zone 2 fractures and post-operative neurological exam. No patient with an intact pre-operative neurologic exam had a neurological deficit after surgery. 86 patients (95.6%) were neurologically intact at their last follow-up examination. Four patients (4.4%) had a neurological deficit at final follow-up, all of them had neurological deficit prior to surgery. 81 patients were treated with partially threaded screws of which 1 (1.2%) had neurological deficit at final follow-up.

Fifty-seven fractures (63.3%) were simple fractures and 33 fractures (36.7%) were comminuted. All four patients with neurological deficit had comminuted fractures. The association between neurologic deficit in zone 2 sacral fracture and fracture comminution was found to be statistically significant (p -value = 0.016). No nonunion was observed in this cohort.

Conclusions: The use of partially threaded screws for zone 2 sacral fractures is associated with low risk for neurologic injury, suggesting that compression through the fracture does not cause iatrogenic nerve damage. The low rate of sacral nonunion can be attributed to compression induced by the use of partially threaded compression screws. There is a strong association between zone 2 comminution and neurologic injury.

© 2016 Elsevier Ltd. All rights reserved.

Introduction

Sacral fractures occur in 23–45% of all pelvic ring injuries, and when found to be unstable, require operative stabilization [1–3]. Sacral fractures are classified according to Denis: extra foraminal (Zone 1), involving the neural foramina (Zone 2) or involving the neural canal (Zone 3) [3]. Zone 2 sacral fractures are reported to

range between 34 and 47.5% of all sacral fractures. Denis originally described the influence of fracture zone on neurological deficit and prognosis, with zone 2 fractures characterized by injury to the L5 through lower sacral nerve roots. Other authors reported the incidence of neurologic damage to be as high as 21–28% of the patients with most patients recovering at least one functional level [4,5].

Several fixation methods for of zone 2 sacral fractures have been examined: posterior plating [6,7], triangular fixation using pedicular screws [8] and sacroiliac or trans-sacral screws [9–13]. Percutaneous fixation with trans-sacral or sacral screws has become a popular method of fixation owing to, low complication rates and good clinical outcomes [12,13]. In addition, the

* Corresponding author at: The University of Alabama at Birmingham, Department of Surgery, Division of Orthopaedic Spine and Trauma Faculty, Office Tower 901, 510, 20th Street South, Birmingham, AL 35294, United States.

E-mail address: amirherm@gmail.com (A. Herman).

biomechanical properties of the sacral screws show 80–85% return of pelvic biomechanical rigidity after instrumentation [14,15].

One potential concern with partially threaded screws is the potential for compression and narrowing of the neural foramen particularly through comminuted fractures, which may result iatrogenic neurologic injury [16,17]. Fully threaded screws represent an alternative to partially threaded screws as a means to avoid this potential complication [18,19].

The purpose of this study was to define the incidence of post-operative neurologic injury and identify risk factors for neurological deficit after zone 2 sacral fractures. The authors hypothesized that partially threaded posterior screws (trans-sacral, sacral) would not result in iatrogenic neurologic injury, and that fracture comminution through the neural foramen would predict injury-related neurologic compromise.

Methods

Following institution IRB approval, a retrospective review of all pelvic ring fractures and fracture dislocations fixed at a single level 1 trauma academic center from September 2010 to September 2014 was performed using CPT and ICD-9 codes. Patient inclusion criteria included: age > 18 years at the time of injury and closed zone 2 sacral fracture according to Denis. Exclusion criteria were: lack of neurological examination after surgery, patients that did not have neurologic examination prior to surgery and had a neurologic deficit after surgery, open fractures, and any patient transferred after fixation to an outside facility.

We extracted patient demographic data including age and gender; as well as injury profiles (mechanism of injury, abbreviated injury score (AIS) and the injury severity score (ISS)). Neurological status at arrival, first available neurologic exam after surgery and last clinical follow-up neurological examination were obtained from consult, post-operative, in-patient progress, and clinic notes, respectively. Further data collection included: comminuted vs simple fractures, the surgical intervention of open vs closed reduction, and posterior tension band plate fixation versus percutaneous screws, as well as screw type (partially threaded vs fully threaded), and screw number.

Neurologic deficit at final follow up was defined as motor strength of less than five (out of five) of the injured extremity.

Statistical analysis

Statistical analysis was performed by an experienced biostatistician (A.H.) using SPSS © 23.0 (Chicago, IL, USA). Categorical data are presented as count (percent). Continuous data are presented as mean (\pm standard deviation). The data were divided to patients with intact and deficient neurological examination at final follow-up. All the collected variables were compared between these two study groups. Comparisons between continuous variables were done by the Wilcoxon–Mann–Whitney rank sum test. Comparisons between categorical data were done with chi-square test or the Fisher exact test. The later test was used if expected count was less than five in any cell. All *p*-values reported are two-sided.

Results

Initial data review included 100 patients with zone 2 sacral fractures. After reviewing the patients' data, seven patients were excluded because of lack of neurological examination at follow-up. These included one patient that died shortly after admission, five patients that were transferred intubated and followed at another hospital and one patient that had an above knee amputation the same side as the sacral fracture. Three additional patients were

excluded because they did not have a neurological examination prior to surgery and had a neurologic deficit after surgery.

The study population included 90 patients, mean age was 39.10 (\pm 15.03). Of the 90 patients, 36 patients (40.0%) were male and 54 patients (60.0%) were female. The most common cause of injury was motor vehicle accident – 52 patients (57.8%) followed by fall from height – 12 patients (13.3%). There was no statistically significant difference in the demographic characteristics between neurologically intact and deficient patients (Table 1).

Of the 90 patients, 86 (95.6%) had no neurological deficit at any time point. Four patients (4.4%) had a neurological deficit at final follow-up. None of the four patients with final neurological deficit had an intact examination at time of admission. These four patients presented with weak but present initial motor exam defined as 2–3/5 motor strength in the tibialis anterior, extensor hallucis longus, flexor hallucis longus or gastrocnemius–soleus complex. No patient presenting with an initially intact neurologic exam developed a neurologic deficit following operative fixation (Table 2).

Table 1
Demographic and injury related data.

	Neurologically intact PostOp (N=86)	Neurological deficit PostOp (N=4)	P-value
Age	39.21 (\pm 15.22)	36.75 (\pm 10.43)	0.887
Gender			
Male	34 (39.5%)	2 (50.0%)	
Female	52 (60.5%)	2 (50.0%)	0.676
Mechanism of injury			
Assault	1 (1.2%)	00 (00%)	
Crush Injury	5 (5.8%)	00 (00%)	0.931
Fall from height	12 (14.0%)	00 (00%)	
Fall from horse	2 (2.3%)	00 (00%)	
MCC	5 (5.9%)	00 (00%)	
MVC	48 (55.8%)	4 (100.0%)	
Pedestrian vs MV	10 (11.6%)	00 (00%)	
Tornado	3 (3.5%)	00 (00%)	
AIS–Abdomen	2.38 (\pm 0.61)	3.33 (\pm 1.53)	0.226
AIS–Extremities	3.02 (\pm 0.88)	3.25 (\pm 0.96)	0.595
AIS–Injury severity score	22.34 (\pm 10.62)	30.5 (\pm 8.34)	0.087

AIS=Abbreviated Injury Score; MVC=Motor vehicle collision; MCC= motor cycle collision; MV= motor vehicle.

Table 2
Pelvic injury and surgery related data.

	Neurologically intact PostOp (N=86)	Neurological deficit PostOp (N=4)	P-value
Side of zone 2 fracture			
Left	42 (48.8%)	2 (50.0%)	
Right	33 (38.4%)	2 (50.0%)	0.886
Bilateral	11 (12.8%)	0 (00.0%)	
Fracture pattern			
Simple	57 (66.3%)	0 (0.0%)	
Comminuted	29 (33.7%)	4 (100.0%)	0.016
Fixation method			
ORIF and plate	5 (5.8%)	2 (50.0%)	
CRPP and screw	81 (94.2%)	2 (50.0%)	0.001
Type of screws			
Sacroiliac screws	35 (43.2%)	0 (00.0%)	
Trans-sacral screws	46 (56.8%)	2 (100.0%)	0.506
Number of screws			
1 screw	48 (59.3%)	2 (100.0%)	
2 screws	25 (30.9%)	0 (00.0%)	0.717
3 screws	5 (6.2%)	0 (00.0%)	
4 screws	3 (3.7%)	0 (00.0%)	
Type of screw threads			
Partially threaded	80 (98.8%)	1 (50.0%)	
Fully threaded	1 (1.2%)	1 (50.0%)	0.049

Download English Version:

<https://daneshyari.com/en/article/6082776>

Download Persian Version:

<https://daneshyari.com/article/6082776>

[Daneshyari.com](https://daneshyari.com)