



The prevalence of hyper- and hypothyroidism in patients with ulcerative colitis

Giovanni Casella ^a, Elisabetta De Marco ^a, Elisabetta Antonelli ^c,
Marco Daperno ^d, Vittorio Baldini ^a, Stefano Signorini ^b, Federica Sannella ^c,
Antonio Morelli ^c, Vincenzo Villanacci ^e, Gabrio Bassotti ^{c,*}

^a Division of Internal Medicine, Italy

^b Clinical Pathology Section, Desio General Hospital, Italy

^c Gastroenterology and Hepatology Section, Department of Clinical and Experimental Medicine, University of Perugia, Italy

^d Gastroenterology and Digestive Endoscopy Unit, Mauriziano Hospital, Torino, Italy

^e 2nd Pathology Section, Spedali Civili di Brescia, Italy

Received 30 July 2008; received in revised form 23 August 2008; accepted 1 September 2008

KEYWORDS

Hyperthyroidism;
Hypothyroidism;
Thyroid disease;
Ulcerative colitis

Abstract

Background: The association between ulcerative colitis and thyroid disorders has been previously reported. However, most reports consist of single case description, and a systematic assessment of this relationship has only sporadically been investigated.

Aims: To study a cohort of patients with ulcerative colitis to establish the prevalence of hyper- and hypothyroidism.

Material and methods: During a four-year period, we studied thyroid function in 162 ulcerative colitis patients (62 men, 100 women, age range 18–78 years).

Results: Thyroid dysfunction was present in 4 patients (2.5%) of the overall population and was represented by both hypo- (3 patients) and hyperthyroidism (1 patient). The incidence of this kind of thyroid dysfunction was significantly ($p=0.03$) lower than that found in a large (more than 5000 subjects) control group.

Conclusions: We conclude that the prevalence of hyper-/hypothyroidism is relatively low in patients with ulcerative colitis, at least in our country, and does not justify a systematic investigation of the thyroid function, except in selected cases, probably those with scarce or no response to standard therapeutic measures.

© 2008 European Crohn's and Colitis Organisation. Published by Elsevier B.V. All rights reserved.

1. Introduction

The association between ulcerative colitis (UC) and thyroid disease was firstly reported in 1968, with an emphasis on the increased therapeutic difficulties when thyroid abnormalities

* Corresponding author. Clinica di Gastroenterologia ed Epatologia, Ospedale Santa Maria della Misericordia, Piazza Menghini, 1, 06156 San Sisto (Perugia), Italy. Fax: +39 075 584 7570.

E-mail address: gabassot@tin.it (G. Bassotti).

are not recognized.¹ Since then, several reports appeared in the literature, mostly describing single case reports or very small groups of patients,^{2–6} and only two studies reported series including more than 10 subjects; one investigated 300 UC patients and the other reported on a mixed series of 31 UC and Crohn's disease patients.^{7,8}

The purpose of the present study was to assess the prevalence of hyper- and hypothyroidism in a series of UC patients.

2. Materials and methods

Retrospective data of a five-year period (December 2002–December 2007) on one hundred sixty-two consecutive UC patients (62 men, 100 women, mean age 43 (range 18–78) years) were obtained. The disease was classified according to standard endoscopic and histological criteria.^{9,10} Disease activity was defined as mild, moderate and severe according to Truelove and Witt's classification.¹¹

Thyroid hormones (fT3, fT4, TSH) were assessed in all UC patients by immunochemiluminescent assay (Roche Diagnostics GmbH, Mannheim, Germany) and, only if hormonal alteration were present, further evaluation was carried out by assessing anti-thyroglobulin and anti-thyroperoxidase antibodies in hypothyroidism (immunochemiluminescent assay, Roche) and anti-TSH receptor (2nd generation TRAb assay TRAK human DYNOfest, BRAHMS Diagnostica GmbH, Berlin, Germany)¹² in hyperthyroidism.¹³ These cases were also investigated with ultrasound and nuclear (scintigraphy) imaging.

Incidence data of thyroid dysfunction in UC patients was compared to that obtained in a group of 5721 subjects (1037 men, 4684 women, mean age 48 (range 20–75) years) undergoing a screening program for thyroid disorders with the above methods in the general population of the same geographic area in the period January 2003–December 2007.

3. Statistical analysis

Patients' and controls' data were compared by means of chi-square test with Yates' correction for continuity and Mann–Whitney *U* test, where appropriate. Values of $p < 0.05$ were chosen for rejection of the null hypothesis.

4. Ethical considerations

After carefully explaining the aims of the investigation the patients gave informed consent, and the study was carried out in accordance with local ethical guidelines, following the recommendations of the Declaration of Helsinki (Edinburgh revision, 2000).

5. Results

Table 1 summarizes the disease characteristics of the UC study population. Thyroid disorders were present in 4 patients (2.5%) of the overall UC population and were represented by both hypo- and hyperthyroidism. A 60 year old woman had hyperthyroidism, diagnosed 1 year before the onset of UC. Thyroid function showed a marked increase of FT4 (25 ng/ml n.v. 7–12), inhibition of TSH (<0.01 U/L) and

Table 1 Disease characteristics of the UC study population

	UC patients (total)	Subgroup with UC and thyroid disease
	<i>n</i>	<i>n</i>
Total number	162	4
Age yr (range)	45 (18–78)	65 (55–77)
Women n	100	2
Duration of disease yr (range)	9 (2–15)	18 (12–26)
Location		
Proctitis	8	
Left-sided colitis	132	3
Extensive colitis	22	1
Surgery		1
Disease activity		
Severe	31 (15 women)	4
Moderate	82 (44 women)	
Mild	49 (41 women)	

the presence of antibodies anti-TSH receptor. Ultrasonographic features and thyroid scintigraphy were consistent with Graves' disease. In this patient the clinical condition improved after the starting of methimazole therapy; however after 5 years of follow-up total colectomy was performed for the presence of severe dysplasia in the left colon. Thyroid function remained stable after surgery. In the other 3 patients (1 woman and 2 men, age range) autoimmune hypothyroidism with the presence of both anti-thyroglobulin and anti-microsomal antibodies was diagnosed. In two cases the diagnosis was made 1 and 10 years after the onset of ulcerative colitis. In these three patients a marked increase of TSH (15 mU/L n.v. 0.7–5) and low blood value of FT4 (5 ng/ml n.v. 8–19) was documented. Ultrasonography findings revealed a normal size of the thyroid gland with altered ultrasound pattern, suggesting an autoimmune thyroid process, whereas thyroid scintigraphy showed a decreased uptake of the radioisotope. All the three patients have been successfully treated by oral L-thyroxine therapy. None of these three patients had extraintestinal manifestations of UC.

As potential predictive factors associated with thyroid dysfunction we were able to identify a more advanced age ($p=0.009$), severity ($p=0.001$) and duration of the disease ($p<0.0001$) (Table 1), in that all four patients with thyroid dysfunction were also more than 55 years old, had severe UC, and the onset of disease was >10 years for each patient.

The incidence of thyroid dysfunction (both hypothyroidism, 360 subjects, and hyperthyroidism, 69 subjects) in the general population was 7.4% (429 out 5721 subjects, mean age 62 (range 54–71) years), significantly higher than that found in the patients' population cohort (chi-square 4.8, $p=0.027$, 95% CI 2.4–7.4).

6. Discussion

An increased prevalence of thyroid disorders (2–4 times higher than in the general population) has been previously reported in UC patients.¹ In subsequent studies on the topic (mostly case reports), both hypo- and hyperthyroidism were

Download English Version:

<https://daneshyari.com/en/article/6100885>

Download Persian Version:

<https://daneshyari.com/article/6100885>

[Daneshyari.com](https://daneshyari.com)