

**LONGITUDINAL CHANGES OF BLOOD PRESSURE AFTER WEIGHT LOSS:
FACTORS INVOLVED.**

Authors: L Flores, PhD^{1,2}, J Vidal, PhD^{1,2,3}, I Núñez, PhD⁴, S Rueda, MD¹, J Viaplana, MD³, E Esmatjes, PhD^{2,3,5}.

Affiliations:

¹ Obesity Unit, Department of Endocrinology and Nutrition, Hospital Clinic.

² CIBERDEM: Spanish Biomedical Research Centre in Diabetes and Associated Metabolic Disorders.

³ IDIBAPS: Institut d'Investigacions Biomèdiques August Pi i Sunyer.

⁴ Diagnostic Imaging Center, Hospital Clinic.

⁵ Diabetes Unit, Department of Endocrinology and Nutrition. Hospital Clinic i Provincial, Barcelona, Spain.

Short title: Obesity: effect of weight loss on blood pressure.

Source of funding: This study received a grant (PI07/0124) from the Fondo de Investigación Sanitaria (FIS) del Instituto de Salud Carlos III.

Correspondence to:

Dr Lilliam Flores, PhD, MD;

Obesity Unit; Department of Endocrinology and Nutrition;

Hospital Clinic i Universitari;

Villarroel, 170, Escalera 11, 2^o piso. CP 08036, Barcelona, Spain.

Phone: 0034658023562; Fax: 0034934516638; e-mail: lflores@clinic.ub.es

Word Count: 2962; Tables: 5

Material Supplementary: Tables 4

Download English Version:

<https://daneshyari.com/en/article/6111434>

Download Persian Version:

<https://daneshyari.com/article/6111434>

[Daneshyari.com](https://daneshyari.com)