Accepted Manuscript

Title: Rapid emergence of high-level tigecycline resistance in *Escherichia coli* strains harbouring *bla_{NDM-5}* in vivo

Author: Xi Li Xinli Mu Yunxing Yang Xiaoting Hua Qing Yang Nanfei Wang Xiaoxing Du Zhi Ruan Xiaoqiang Shen Yunsong Yu



PII: DOI: Reference:	S0924-8579(16)00039-X http://dx.doi.org/doi:10.1016/j.ijantimicag.2016.01.005 ANTAGE 4743				
To appear in:	International	Journal	of	Antimicrobial	Agents
Received date: Revised date: Accepted date:	15-10-2015 15-1-2016 20-1-2016				

Please cite this article as: Li X, Mu X, Yang Y, Hua X, Yang Q, Wang N, Du X, Ruan Z, Shen X, Yu Y, Rapid emergence of high-level tigecycline resistance in *Escherichia coli* strains harbouring *bla*_{NDM-5} in vivo, *International Journal of Antimicrobial Agents* (2016), http://dx.doi.org/10.1016/j.ijantimicag.2016.01.005

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ACCEPTED MANUSCRIPT

Highlights

- This report is the first to demonstrate high-level tigecycline (TIG) resistance in *Escherichia coli* in vivo.
- We report the first treatment-emergent MIC development of TIG from 1 mg/L to 64 mg/L in *E. coli*.
- Clinicians should be aware of the risk of an increase in the MIC of TIG under therapy.

Page 1 of 24

Download English Version:

https://daneshyari.com/en/article/6117598

Download Persian Version:

https://daneshyari.com/article/6117598

Daneshyari.com