



Short communication

Production of equine herpesvirus 1 recombinant glycoprotein D and development of an agar gel immunodiffusion test for serological diagnosis



Nadia Fuentealba^{a,e,*}, Guillermo Sguazza^a, Mariela Scrochi^{a,e}, Maria Bravi^{a,g},
Carolina Zanuzzi^{b,c,e}, Santiago Corva^d, Eduardo Gimeno^{c,e}, Marcelo Pecoraro^a,
Cecilia Galosi^{a,f}

^a Department of Virology, School of Veterinary Sciences, National University of La Plata, 60 & 118, P.O. Box 296, 1900 La Plata, Buenos Aires, Argentina

^b Department of Histology and Embryology, School of Veterinary Sciences, National University of La Plata, 60 & 118, P.O. Box 296, 1900 La Plata, Buenos Aires, Argentina

^c Department of Pathology, School of Veterinary Sciences, National University of La Plata, 60 & 118, P.O. Box 296, 1900 La Plata, Buenos Aires, Argentina

^d Department of Epidemiology, School of Veterinary Sciences, National University of La Plata, 60 & 118, P.O. Box 296, 1900 La Plata, Buenos Aires, Argentina

^e National Research Council (CCT-CONICET-La Plata), Argentina

^f Scientific Research Commission of the Province of Buenos Aires (CIC-PBA), Argentina

^g Argentine Agency for the Promotion of Science and Technology (ANPCyT), Argentina

A B S T R A C T

Article history:

Received 31 October 2013

Received in revised form 24 February 2014

Accepted 27 February 2014

Available online 11 March 2014

Keywords:

Agar gel immunodiffusion test

Equine herpesvirus 1

Recombinant glycoprotein D

Virus neutralization

Equine herpesvirus 1 and 4 (EHV-1 and 4) infect most of the world's horses, causing serious clinical illness. Viral glycoproteins have been identified as the immunodominant antigens that generate the antiviral serological responses to EHV-1 and EHV-4 in infected horses. Here, glycoprotein D of EHV-1 was expressed by a recombinant baculovirus, purified and evaluated by a simple agar gel immunodiffusion test (AGID). Compared with virus neutralization, serological analysis by AGID showed good specificity (100%) and sensitivity (99.5%). The estimated Kappa values for repeatability and reproducibility were satisfactory. Thus, this rapid, inexpensive, simple and highly specific AGID test seems to be a valuable alternative tool for serological detection of antibodies against both EHV-1 and EHV-4.

© 2014 Elsevier B.V. All rights reserved.

Equine herpesvirus 1 (EHV-1) and Equine herpesvirus 4 (EHV-4) are ubiquitous herpesviruses that infect most of the world's horses at some time during their lives, causing respiratory disease, abortion and neurological disorders (Allen and Bryans, 1986). After the initial lytic phase of infection in the equine respiratory tract, EHV-1 and EHV-4 enter a latent state in both circulating lymphocytes and lymphocytes in draining lymph nodes, as well as in sensory nerve-cell bodies within the trigeminal ganglia. Recurrent shedding of the virus from asymptomatic carriers may

spread the disease in the equine population (Baxi et al., 1996). Several studies have identified glycoproteins B, C, and D (gB, gC and gD, respectively) as the immunodominant antigens that generate antiviral serological responses to EHV-1 and EHV-4 in infected horses (Allen et al., 1992). EHV-1 and EHV-4 are closely related both genetically and antigenically. Diagnoses of both viruses are based on virus isolation, histopathology, immunohistochemical staining, or antibody detection, but they can show cross-reactivity in many serological tests such as virus neutralization and some ELISAs (http://www.oie.int/fileadmin/Home/eng/Health_standards/tahm/2.05.09.EQUINE.RHINO.pdf, access September 24, 2013). Serological tests such as virus neutralization and ELISA require special facilities or commercial equipment. In contrast, the agar gel immunodiffusion test (AGID), which was designed for other equine viruses as an effective tool for the detection of specific antibodies (Coggins and Norcross, 1970), is a fast technique that does not require special conditions and can be performed at any veterinary diagnostic laboratory.

In this work, a purified recombinant gD of EHV-1 was produced and evaluated by a simple AGID test performed to examine a serum

* Corresponding author at: Department of Virology, School of Veterinary Sciences, National University of La Plata, 60 & 118, P.O. Box 296, 1900 La Plata, Buenos Aires, Argentina. Tel.: +54 221 4824956; fax: +54 221 4824956.

E-mail addresses: nadiafuatealba@fcv.unlp.edu.ar, nadiafuatealba@hotmail.com (N. Fuentealba), sguazza@fcv.unlp.edu.ar (G. Sguazza), scrochimariela@gmail.com (M. Scrochi), murguita@hotmail.com (M. Bravi), carozanuzzi@gmail.com (C. Zanuzzi), sgcorva@fcv.unlp.edu.ar (S. Corva), ejgimeno@gmail.com (E. Gimeno), peco@fcv.unlp.edu.ar (M. Pecoraro), cmgalosi@fcv.unlp.edu.ar (C. Galosi).

samples with the same specificity and sensitivity as virus neutralization in a short time.

For recombinant gD production, rabbit kidney RK13 cells (Argentine Cell Bank Association, Argentina) infected with the EHV-1 AR8 strain were used. Infected cells were collected, treated with proteinase K (Promega, Madison, Wisconsin, USA) and total DNA was extracted with a commercial kit (Promega). DNA was quantified by measuring absorbance at an OD₂₆₀/OD₂₈₀ ratio in a SmartSpec™ 3000 spectrophotometer (BIO-RAD, California, USA). A polymerase chain reaction (PCR) was used to generate a 1373-bp fragment encompassing the EHV-1 gD open reading frame (ORF) flanked by 5'SnaBI and 3'XmaJI sites, using a primer pair (5'-AGTTACGTATTATGCGCTGTGTGCTG-3' and 5'-GTTTACCTAGGCTGGGTATATTAACATCC-3'). The PCR was carried out in a Mastercycler Gradient (Eppendorf, Hamburg, Germany), in a final volume of 25 µl, using 2.5 µl of 10X Taq buffer with KCl, 2.5 U of Taq DNA polymerase (Fermentas, Tecnolab, Buenos Aires, Argentina), 1.5 µM MgCl₂, 0.2 mM each of dATP, dCTP, dGTP, and dTTP (Promega), 0.5 µM of each primer and 2 µl of DNA sample (concentration 5.6 µg/ml). The DNA was amplified with an initial denaturation step of 94 °C for 4 min, followed by 30 cycles consisting of 94 °C denaturation for 30 s, 60 °C annealing for 20 s, and 72 °C extension for 1 min. The PCR products were run on 1.5% agarose gel in TBE buffer (45 mM Tris-borate, 1 mM ethylenediamine tetra-acetic acid-EDTA-, pH 8), stained with ethidium bromide at a final concentration of 0.5 µg/ml and purified using a gel extraction kit Wizard SV Gel & PCR Clean Up (Promega). The PCR product coding for EHV-1 gD was cloned into the vector pCR2.1-TOPO using the TOPO TA Cloning Kit (Life Technologies Corporation, Carlsbad, California, USA) and then digested with restriction endonucleases corresponding to restriction sites added to the primers. The fragment obtained corresponding to the gD gene was inserted into the multiple cloning site of the transfer plasmid pFastBacHTB between the StuI and XbaI sites (isoschizomers of SnaBI and XmaJI, respectively) and transformed into DH10Bac cells (Invitrogen, Carlsbad, California, USA). These restriction sites allowed fusion of EHV-1 gD in frame with the N-terminal 6X His tag and controlled by *Autographa californica* multiple nuclear polyhedrosis virus (AcMNPV) polyhedrin promoter. The pFastBac-gD construction was sequenced using the primers pFastBacF: 5'-TATTCCGGATTATTCATACC-3' (3994–4014) and pFastBacR: 5'-TTCAGTTCAGGGGAGGTG-3' (4298–4318) and the sequence of the resulting ORF was examined using the Vector NTI Advance™ software version 11.0 (Invitrogen). Recombinant baculovirus was obtained in *Spodoptera frugiperda* (Sf21) cells (Gibco, Grand Island, NY, USA) transfected with recombinant bacmid DNA. For glycoprotein expression High Five cells (Invitrogen) were infected with 1 PFU of recombinant baculovirus/cell. The infected cell extracts were harvested at 12, 24 and 48 h post-infection and examined by 10% SDS-PAGE and Western Blot according to standard procedures. The location of gD EHV-1 in infected insect cells was determined by immunofluorescence using horse anti-EHV-1 polyclonal antibody, provided by Dr. T. Kumanomido (Equine Research Institute, Japan Racing Association, Tochigi, Japan) and anti-horse fluorescein isothiocyanate conjugated (Sigma–Aldrich, Saint Louis, Missouri, USA). Visualization was performed on an epifluorescence microscope Olympus System – Model BHS (Olympus Corporation, New York, USA).

After determining the time of greatest expression of gD, infected cells were harvested, lysed by freezing and thawing, clarified by centrifugation and purified by immobilized metal ion affinity chromatography (IMAC) using Ni-NTA agarose (Qiagen, Maryland, USA). The elution was performed with 50 mM sodium phosphate buffer and 500 mM sodium chloride at decreasing pHs (6, 5 and 4). SDS-PAGE and Western blot were used to detect the purified protein. Purified gD was concentrated in a cellulose membrane (M.W.

14,000, Sigma–Aldrich) with polyethylene glycol 6000 (Biopack, Buenos Aires, Argentina) and its concentration was determined by the standard method of Bradford, 1976.

The AGID test was performed according to the previously described method for Equine infectious anemia (EIA) virus (Coggins and Norcross, 1970), but using two different agar solutions: one consisting of 1% Difco™ agar granulated (Becton Dickinson, Sparks, Maryland, USA) in 0.145 M borate buffer (9 g H₃BO₃, plus 2 g NaOH per liter) pH 8.6 (±0.2) and the other consisting of 1% Difco agar in Tris buffer 0.2 M and NaCl 0.85%, pH 7.2. The gD antigen was assayed at decreasing concentrations of 300, 150, 75, 50, 37.5, 18.7 and 9.4 µg/ml using one EHV-1-positive reference horse serum (1:64 and 1:8 neutralizing antibody titers for EHV-1 and -4, respectively) and one negative horse serum (Equine Research Institute, Japan Racing Association, Tochigi, Japan). The AGID was carried out in a Petri dish using a punch of six wells of 4 mm (~40 µl) in diameter and 3 mm apart and incubated at room temperature in a humid chamber. The results were read at 48 h. After determining the optimal concentration of gD, reference sera for EIA virus, Equine Influenza virus (EIV), Equine arteritis virus (EAV) and Equine herpesvirus 2 (EHV-2) were tested. Then, two-fold serial dilutions in phosphate buffer saline (PBS) of an EHV-1-positive horse serum (neutralizing antibody titers 1:128) were examined by AGID. A total panel of 396 serum samples previously tested by virus neutralization and some of which (n = 38) showed severe hemolysis were also examined by AGID. In addition, the 10 sera of the panel with 1:256 neutralizing antibody titers were examined in two-fold serial dilutions. Repeatability and reproducibility of AGID were evaluated four times with 32 sera (16 positive and 16 negative) and by four different operators. The Kappa index of concordance was determined at both intra- and inter-operator level.

The virus neutralization test was performed in microplates using two-fold serial dilutions of sera (two replicate wells) mixed with an equal volume (25 µl) of 100 TCID₅₀ of the AR8 EHV-1 strain and after 60 min of incubation in a 5% CO₂ atmosphere, RK-13 cells (100 µl of 5 × 10⁵ cells/ml) were added. The results were read 5 days later and neutralizing antibody titers were expressed as the highest serum dilution that protects 100% of the cell monolayer from virus destruction in both wells. The effectiveness of the present AGID test was compared with the virus neutralization test, and the sensitivity and specificity were evaluated with ROC Curve analysis (Dohoo et al., 2009).

Western blot results showed that EHV-1 gD was expressed in readily detectable antigenic forms of gD after 12 h of infection with a maximum concentration at 48 h. The High Five cells infected with the recombinant baculovirus showed a similar banding pattern in all samples at different times post-infection, with one main band of 65 kDa and another band at a lower concentration (Fig. 1). After purification using Ni-NTA, SDS-PAGE also revealed two specific bands of 58 and 65 kDa in cell samples at 48 h post-infection. Fluorescence specificity was clearly seen on the cell surface and in the cytoplasm but not in cell nuclei (data not shown). After purification, recombinant gD was obtained at 300 µg/ml with a high purity level, as evaluated by the Western blot.

The AGID reaction was observed when the borate buffer was used. Clear precipitation lines were obtained using antigen concentrations of 50 µg/ml and EHV-1-positive reference horse serum (Fig. 2A) and up to a 1/8 dilution of the EHV-1-positive horse serum (Fig. 2B). No positive reactions were observed with reference EIA, EIV, EAV and EHV-2 sera, since these viruses are antigenically, genetically, and pathogenetically distinct from EHV-1 and EHV-4 (Allen et al., 2004).

The AGID test detected 182 negative and 213 positive sera with different neutralizing antibody titers (1:256 n = 10; 1:128 n = 21; 1:64 n = 22; 1:32 n = 46; 1:16 n = 82 and 1:8 n = 32). One serum with doubtful neutralizing antibody titer at a 1:4 dilution was considered

Download English Version:

<https://daneshyari.com/en/article/6133746>

Download Persian Version:

<https://daneshyari.com/article/6133746>

[Daneshyari.com](https://daneshyari.com)