



Development and validation of TaqMan quantitative PCR for detection of frog virus 3-like virus in eastern box turtles (*Terrapene carolina carolina*)

Matthew C. Allender^{a,*}, David Bunick^a, Mark A. Mitchell^b

^a Department of Comparative Biosciences, College of Veterinary Medicine, University of Illinois, 2001 South Lincoln Avenue, Urbana, IL 61802, United States

^b Department of Veterinary Clinical Medicine, College of Veterinary Medicine, University of Illinois, 1008 West Hazelwood Drive, Urbana, IL 61802, United States

ABSTRACT

Article history:

Received 8 August 2012

Received in revised form 4 October 2012

Accepted 17 December 2012

Available online 26 December 2012

Keywords:

Reptile
Ranavirus
Epidemiology
Molecular

Ranavirus has caused disease epidemics and mass mortality events globally in free-ranging fish, amphibian, and reptile populations. Viral isolation and conventional PCR are the most common methods for diagnosis. In this study, a quantitative real-time PCR (qPCR) assay was developed using a TaqMan probe-based assay targeting a highly conserved region of the major capsid protein of frog virus 3-like virus (FV3-like) (Family *Iridoviridae*, genera *Ranavirus*). Standard curves were generated from a viral DNA segment cloned within a plasmid. The assay detected viral DNA 1000 times lower than conventional PCR. Thirty-one clinical samples (whole blood and oral swabs) from box turtles were tested using these assays and the prevalence of the virus determined. Quantitative PCR allows for a superior, rapid, sensitive, and quantitative method for detecting FV3-like virus in box turtles, and this assay will be useful for early detection and disease monitoring.

© 2012 Elsevier B.V. All rights reserved.

1. Introduction

Ranavirus is one of five genera from the family *Iridoviridae* and one of three genera reported to infect reptiles. They are large, icosahedral, DNA viruses that have emerged as a significant cause of mortality events in free-ranging herpetofauna throughout the world (Green et al., 2002; Johnson et al., 2008; Miller et al., 2011). *Ranavirus* has specifically been linked as a contributing factor in amphibian declines, with numerous outbreaks occurring in the US (Green et al., 2002; Jancovich et al., 2003; Johnson et al., 2008; Miller et al., 2011). Recently, ranaviral disease in amphibians was placed on the World Organization for Animal Health list of reportable diseases. Disease events in amphibians are often clustered into local epizootics, with significant impact on the local population (Green et al., 2002). These epizootics have been scattered across numerous habitats and landscapes in the US; however, disease predictability has not been successful. Furthermore, there have been increasingly more reports and cases in chelonians, and specifically box turtles, in the US (De Voe et al., 2004; Allender et al., 2006; Johnson et al., 2008).

Diagnostic assays that are validated and optimized to detect the presence of pathogens in certain species are key to characterizing the disease and understanding disease ecology. Quantitative PCR

has not been developed previously for detection of FV3-like virus in turtles. Its use would provide much greater sensitivity in detection of FV3-like virus as compared with conventional PCR, and additionally would allow the quantitation of virus levels within specific tissues. Identifying clinical samples with the highest viral load will allow for more efficient sampling methods that target those tissues and potentially direct therapy toward those sites. Additionally, the greater sensitivity of qPCR will allow for the detection of ranavirus in animals with lower titers of virus, as might be seen in early or subclinical infections. In studies investigating iridoviruses in fish, TaqMan real-time PCR was shown to be 100 times more sensitive than conventional PCR, and crucial for the identification of subclinical disease states (Getchell et al., 2007; Pallister et al., 2007). Animals with subclinical infections may serve as important carriers or reservoirs for infectious disease; therefore, it is critical to develop assays capable of detecting the pathogen in these animals. To date, no such assay has been reported for ranavirus quantification in free-ranging or captive chelonians.

The purpose of this study was to develop and evaluate diagnostic methods for characterizing an emerging pathogen, FV3-like virus (genera *Ranavirus*, Family *Iridoviridae*) in box turtles. The hypotheses tested in this study were that a qPCR TaqMan based assay would be both sensitive and specific for characterizing FV3-like virus in box turtles. This is essential when considering the application of these assays to additional free-ranging populations and/or experimental models. Furthermore, it allows for evaluating potential climatic and environmental impacts of the disease, treatment, and management options.

* Corresponding author. Tel.: +1 217 265 0320.

E-mail addresses: mattallender@vetmed.illinois.edu, mcallend@illinois.edu (M.C. Allender).

2. Materials and methods

2.1. DNA extraction

Whole blood and an oral swab were collected from a positive eastern box turtle that presented to the University of Tennessee (Knoxville, TN, USA) wildlife clinic in 2007. DNA was extracted following manufacturer's instructions (QIAamp DNA blood mini kit, Qiagen, Valencia, CA). Concentrations and purity of DNA were determined using a spectrophotometer (Nanodrop spectrophotometer, Thermo Scientific, Wilmington, DE).

2.2. Conventional PCR, sequencing, and cloning

Conventional PCR was performed using developed previously primers (sense: 5'-GACTTGGCCACTTATGAC-3' and antisense: 5'-GTCTCTGGAGAAGAAGAA-3') (MCP 4) targeting a portion of the major capsid protein (Mao et al., 1997). Products were sequenced in both directions using the University of Illinois Core DNA Sequencing Facility (Urbana, IL, USA) and compared to known sequences in GenBank using TBLASTX. For cloning, the PCR product (531 bp) from the MCP 4 primers was then cloned in *Escherichia coli* (TOPO TA Cloning® kit, Invitrogen, Carlsbad, CA). The cloning product was verified through sequencing in both directions. Plasmids were linearized with EcoR1, purified (QIAfilter plasmid Maxi kit, Qiagen, Valencia, CA), and quantified using spectrophotometry. Ten-fold serial dilutions of linearized plasmids were made from 10.0×10^2 ng/μl to 10×10^{-8} ng/μl. Viral genome (DNA) copy number was calculated using the following formula:

$$\# \text{copies}/\mu\text{L} = \frac{(\text{ng DNA of plasmid} + \text{clone}/\mu\text{L})(6.022 \times 10^{23} \text{ copies/mol})}{(\text{bp length})(1 \times 10^9 \text{ ng/g})(650 \text{ g/mol of bp})}$$

The final copy number for 10-fold serial dilutions ranged from 5.29×10^9 to 5.29×10^1 viral copies per reaction.

2.3. Real time qPCR assay

A primer/probe assay for a TaqMan-MGB (TaqMan® primers, FAM dye labeled, Applied Biosystems, Carlsbad, CA) based qPCR assay were designed using a commercial software program (Primer Express®, Applied Biosystems, Carlsbad, CA) based on published sequences of the major capsid protein of FV3 (Mao et al., 1997). TaqMan assay was performed using forward (AACGCCGACCGAAACTG), reverse (GCTGCCAAGATGTCGGGTAA), and probe (CCGGCTTTCCGGC) targeting a 54bp segment of the major capsid protein (MCP) of frog virus 3. Real-time qPCR assays were performed using a real-time PCR thermocycler (7500 ABI real-time PCR System, Applied Biosystems, Carlsbad, CA) and data was analyzed using associated software (Sequence Detection Software v2.05, Applied Biosystems, Carlsbad, CA). Each reaction contained 12.5 μl of 2× TaqMan Platinum PCR Supermix-UDG with ROX (TaqMan Platinum PCR Supermix-UDG with ROX, Invitrogen, Carlsbad, CA), 1.25 μl TaqMan primer-probe, 2.5 μl turtle-derived FV3 dilution, and water to a final concentration of 25 μl. Cycling parameters were as follows: 1 cycle at 50 °C for 2 min followed by 95 °C for 10 min, then 40 cycles at 95 °C for 15 s and 60 °C for 60 s, and a final cycle of 72 °C for 10 min.

2.4. Standard curve, specificity, and sensitivity

To determine the sensitivity, assays were performed in three technical repeats on dilutions of turtle-derived positive control plasmid of FV3 MCP DNA (5.29×10^9 – 5.29×10^1 copies/rxn) within a single run. Standard curves were generated using the cycle threshold values of the positive control plasmid dilutions. Intra-assay variation was determined for both assays by calculating the mean C_t

values, standard deviations, and coefficient of variations separately for each control plasmid DNA dilution. Efficiency curves of the dilutions were performed in uninfected cell culture lysates (spiked with plasmid dilutions), infected cell lysates, and turtle whole blood extracts from a positive sample.

2.5. Box turtle samples

Thirty-one eastern box turtles that were presented to the University of Tennessee College of Veterinary Medicine Wildlife clinic were examined during March through October 2007. Whole blood and swabs of the oral cavity were collected as described previously (Allender et al., 2011). The University of Tennessee Institutional Animal Use and Care Committee approved all animal use (protocol 1864). Whole blood and oral swab extracts of samples from animals with unknown disease status were evaluated. Results for detection were compared to published conventional PCR results for the same samples (Allender et al., 2011).

2.6. Statistical analysis

The quantity of ranavirus target DNA in infected whole blood and oral swabs was determined using a standard curve method. The copy number of the target DNA was determined from the standard curve generated with 10-fold dilutions of the positive control plasmid that contained the target sequence of the respective qPCR assay.

Copy numbers were tabulated and evaluated for normality using the Shapiro–Wilk test. Mean, median, standard deviation, 95% confidence interval, and 10–90% percentiles were determined for positive cases (copy number) for each assay. The Mann–Whitney *U* test was used to evaluate between assay differences. The prevalence of ranavirus was determined for each assay (categorical variable assigned; 1 = positive, 0 = negative). Exact 95% binomial confidence intervals were determined for all proportions. Level of agreement (kappa) was determined between both the real-time PCR assays and the conventional PCR reported previously based on prevalence. All statistical analysis was performed using statistical software (IBM SPSS Statistics 19, Chicago, IL).

3. Results

3.1. Conventional PCR, standard curve, reproducibility

Conventional PCR was performed to evaluate turtle-derived FV3-like virus dilutions using MCP 4 (Mao et al., 1997). The TaqMan primer set was designed to detect a 54 bp length gene segments of a conserved portion of the MCP gene of FV3.

Serial 10-fold dilutions of positive control plasmids were assayed using TaqMan assay and standard curves were generated based on C_t values (Fig. 1). The linear range for TaqMan was between 5.29×10^9 and 5.29×10^4 viral copies with an R^2 of 0.999 (slope = −3.277) (Fig. 1). Efficiency curves carried out on spiked-uninfected cell lysates (controlled for total DNA per rxn), infected cell lysates, and turtle whole blood extracts performed the equally well (data not shown). The amplification plots of TaqMan qPCR are presented in Fig. 2.

The intra- and inter-assay reproducibility was evaluated for the serial dilutions of the control plasmids (Table 1). The intra-assay CVs for TaqMan were between 0.04 and 0.24%. The inter-assay CVs were 0.14–0.3%, respectively. These results indicate high reproducibility between assays at all dilutions. The dynamic range for qPCR assays was from 5.29×10^9 to 5.29×10^1 .

Download English Version:

<https://daneshyari.com/en/article/6134673>

Download Persian Version:

<https://daneshyari.com/article/6134673>

[Daneshyari.com](https://daneshyari.com)