



Short communication

The interaction between human initiation factor eIF3 subunit c and heat-shock protein 90: A necessary factor for translation mediated by the hepatitis C virus internal ribosome entry site

Saneyuki Ujino^a, Hironori Nishitsuji^a, Ryuichi Sugiyama^a, Hitoshi Suzuki^a, Takayuki Hishiki^b, Kazuo Sugiyama^c, Kunitada Shimotohno^b, Hiroshi Takaku^{a,*}

^a Department of Life and Environmental Sciences, Chiba Institute of Technology, 2-17-1 Tsudanuma, Narashino-shi, Chiba 275-0016, Japan

^b Research Institute, Chiba Institute of Technology, 2-17-1 Tsudanuma, Narashino, Chiba 275-0016, Japan

^c Center for Integrated Medical Research, School of Medicine, Keio University, 35 Shinanomachi, Shinjyuku-ku Tokyo 160-8582, Japan

ARTICLE INFO

Article history:

Received 10 June 2011

Received in revised form 4 October 2011

Accepted 6 October 2011

Available online 14 October 2011

Keywords:

eIF3c

Hsp90

HCV IRES

17-AAG

Translation initiation

ABSTRACT

Heat-shock protein 90 (Hsp90) is a molecular chaperone that plays a key role in the conformational maturation of various transcription factors and protein kinases in signal transduction. The hepatitis C virus (HCV) internal ribosome entry site (IRES) RNA drives translation by directly recruiting the 40S ribosomal subunits that bind to eukaryotic initiation factor 3 (eIF3). Our data indicate that Hsp90 binds indirectly to eIF3 subunit c by interacting with it through the HCV IRES RNA, and the functional consequence of this Hsp90–eIF3c–HCV-IRES RNA interaction is the prevention of ubiquitination and the proteasome-dependent degradation of eIF3c. Hsp90 activity interference by Hsp90 inhibitors appears to be the result of the dissociation of eIF3c from Hsp90 in the presence of HCV IRES RNA and the resultant induction of the degradation of the free forms of eIF3c. Moreover, the interaction between Hsp90 and eIF3c is dependent on HCV IRES RNA binding. Furthermore, we demonstrate, by knockdown of eIF3c, that the silencing of eIF3c results in inhibitory effects on translation of HCV-derived RNA but does not affect cap-dependent translation. These results indicate that the interaction between Hsp90 and eIF3c may play an important role in HCV IRES-mediated translation.

© 2011 Elsevier B.V. All rights reserved.

The hepatitis C virus (HCV), a member of the *Flaviviridae* family, has a positive-strand RNA genome (Taylor et al., 1999; Bartenschlager and Lohmann, 2001) encoding a large precursor polyprotein that is cleaved by host and viral proteases to generate at least 10 functional viral proteins: core, envelope 1 (E1), E2, p7, nonstructural protein (NS2), NS3, NS4A, NS4B, NS5A, and NS5B (Grakoui et al., 1993; Hijikata et al., 1993). Ishii et al. identified an HCV replicon system in which the full HCV genomic RNA autonomously replicates in the Huh-7 human hepatoma cell line (Fig. 1A) (Ishii et al., 2006). This HCV replicon system allows researchers to study HCV genome replication in cell culture. HCV protein synthesis is initiated by the HCV RNA genome. This genome contains a conserved structure in its 5'-untranslated region (5'-UTR) that acts as an internal ribosome entry site (IRES) (Lukavsky, 2008). Briefly, the small ribosomal subunit (40S) and the eukaryotic initiation factor eIF3 bind specifically to the HCV IRES RNA, allowing for direct recognition of the start codon present in the 5'-UTR of the viral mRNA (Spahn et al., 2001; Collier et al., 2002; Kieft

et al., 2002; Fraser and Doudna, 2007; Julien et al., 2009). Consistent with its diverse functions, eIF3 is the largest and most complex initiation factor. The mammalian version, for example, contains 13 nonidentical subunits designated eIF3a to eIF3m. The eIF3 core subunit (eIF3a–c, g, and i) is essential for translation (Kieft et al., 2002; Hinnebusch, 2006; Masutani et al., 2007; Zhou et al., 2008), and eIF3 specifically associates with the apical half of domain III of the HCV IRES (Kieft et al., 2001, 2002; Siridechadilok et al., 2005; Fraser and Doudna, 2007).

Hsp90 is a heat-shock protein that is abundant in the cytosol of eukaryotes and prokaryotes. In contrast to other chaperones, a number of substrates are known to contain Hsp90 (Schulte et al., 1995). Studies of eukaryotes have revealed that these Hsp90 client proteins include a variety of transcription factors (Coumaillieu et al., 1995; Garcia-Cardena et al., 1998; Nagata et al., 1999; Sato et al., 2000; Richter and Buchner, 2001; Xu et al., 2001; Waza et al., 2005). Recently, many studies have reported that Hsp90 is involved with not only HCV RNA replication and viral protein but also HCV IRES-mediated translation (Waxman et al., 2001; Kim et al., 2006; Okamoto et al., 2006; Nakagawa et al., 2007; Ujino et al., 2009). In the present study, we demonstrate that eIF3 forms a complex with Hsp90 that is critical for HCV IRES-mediated translation.

* Corresponding author. Tel.: +81 47 478 0407; fax: +81 47 478 0407.

E-mail address: hiroshi.takaku@it-chiba.ac.jp (H. Takaku).




Fig. 1. Inhibition of IRES-mediated translation by an Hsp90 inhibitor. (A) The structure of the HCV replicon RNA molecules comprising the HCV 5'-UTR, including the HCV IRES, the neomycin phosphotransferase gene (Neo^r), and the coding region for the HCV proteins core to NS5B (in the HCV full-length replicon). (B) A schematic representation of the bicistronic HCV IRES or EMCV IRES reporter construct pRenilla-HCV IRES-firefly luciferase (RL-HCV IRES-FL) or pRenilla-EMCV IRES-firefly luciferase (RL-EMCV IRES-FL) driven by the CMV promoter to direct cap-dependent translation of renilla luciferase (RL) and HCV IRES or EMCV IRES-dependent translation of firefly luciferase (FL). The vector construct for HCV IRES-mediated translation of firefly luciferase, pHCV IRES-firefly luciferase (HCV IRES-FL) (Ujino et al., 2010). (C) The structure of the Hsp90 inhibitor 17-AAG (17-allylamino-17-demethoxygeldanamycin, Sigma-Aldrich Chemical Co.). (D) Inhibition of IRES-mediated translation by 17-AAG. Huh-7 cells (1×10^5 cells/well on 12-well plates) treated with 17-AAG (25 and 50 nM) and DMSO as a control for 24 h, and were then transfected with pRenilla-HCV IRES-firefly luciferase (RL-HCV IRES-FL) using Lipofectamine 2000 (Invitrogen). At 24 h post-transfection, Renilla luciferase (cap-dependent translation) and firefly luciferase (HCV IRES-dependent translation) activities were measured with a Dual-Luciferase Reporter Assay System (Promega). The data represent the mean \pm standard deviations (SDs) from the experiments performed in triplicate.

To investigate the effects of the Hsp90 inhibitor 17-AAG on HCV IRES translation, a bicistronic reporter system was used that consisted of an upstream reporter, Renilla luciferase (RL), expressed by cap-dependent translation and a downstream reporter, firefly luciferase (FL), which is under the translational control of the HCV IRES. To construct pcDNA-HCV IRES-firefly Luc, pHCV IRES-firefly Luc (HCV IRES-FL) (Ujino et al., 2010) (Fig. 1B) was digested with BamHI and Sall. The IRES-firefly Luc fragments were inserted into the BamHI-XhoI site of pcDNA3.1 (Invitrogen, Carlsbad, CA). To construct pRenilla-HCV IRES-firefly luciferase (RL-HCV IRES-FL), Renilla luciferase fragments were amplified by PCR from a pFN11A Flexi vector (Promega, Madison, WI), and the PCR products were inserted into the BamHI site of pcDNA-HCV IRES-firefly Luc. The human hepatoma cell line Huh-7 was maintained in Dulbecco's modified Eagle's medium (DMEM; Invitrogen) containing 10% fetal bovine serum (FBS). The benzoquinone ansamycin, the antibiotic geldanamycin (GA) and its less toxic analogue 17-allylamino-17-demethoxygeldanamycin (17-AAG) (Fig. 1C) (Sigma-Aldrich Chemical Co., St Louis, MO) directly bind to the ATP/ADP binding pocket of Hsp90, thus preventing ATP binding and the completion of client protein refolding (Neckers, 2003). The client proteins of Hsp90 appear to shift the role of the primary chaperone from Hsp90 to Hsp70 in cells treated with Hsp90 inhibitors (Doong et al., 2003). It is also well known that 17-AAG causes a modest increase in Hsp70 levels (Morimoto, 1998; Bagatell et al., 2000; Guo et al., 2005). In our previous report, a significant induction of Hsp70 was detected (Ujino et al., 2009).

For the reporter gene assay, Huh-7 cells were treated with different concentrations of the Hsp90 inhibitor, 17-AAG, or DMSO as

a control for 24 h. They were then transfected with the bicistronic reporter construct RL-HCV IRES-FL using Lipofectamine 2000 (Invitrogen), which directs cap-dependent translation of the RL gene and HCV IRES-dependent translation of FL genes (Invitrogen). At 24 h post-transfection, the Renilla luciferase (cap-dependent translation) and firefly luciferase (HCV IRES-dependent translation) activities were measured with a Dual-Luciferase Reporter Assay System (Promega, Madison, WI). In cells treated with 50 nM 17-AAG, firefly luciferase activity was reduced by 55% with RL-HCV IRES-FL, whereas Renilla luciferase activity was mostly maintained (Fig. 1D). The inhibition of HCV IRES-mediated translation occurred in a dose-dependent manner. Recently, Kim et al. (2006) demonstrated that Hsp90 regulates ribosomal function by maintaining the stability of 40S ribosomal proteins such as rpS3 and rpS6. The interaction between the 40S ribosomal proteins and Hsp90 has also been associated with ribosomal activities such as protein synthesis. We also found that the Hsp90 inhibitor 17-AAG influences HCV IRES-mediated luciferase activity, suggesting that 17-AAG inhibited HCV RNA replication and HCV IRES-mediated translation.

The HCV IRES is recognized specifically by the small ribosomal subunit and eIF3 before the initiation of viral translation. Although the degradation of rpS3, a component of the small ribosomal subunit, has been shown to occur in the presence of the Hsp90 inhibitor (Kim et al., 2006), the influence of Hsp90 inhibition on eIF3 is not understood. To determine whether 17-AAG affects the expression of the eIF3 subunit, we analyzed eIF3a, eIF3b, eIF3c, eIF3g and eIF3i protein expression by western blot analysis. The HCV replicon cell line NNC#2 (NN/1b/FL), which carries a full genome replicon, was cultured in DMEM with 10% FBS, nonessential amino acids,

Download English Version:

<https://daneshyari.com/en/article/6143326>

Download Persian Version:

<https://daneshyari.com/article/6143326>

[Daneshyari.com](https://daneshyari.com)