

Accepted Manuscript



Incidence of posterior reversible encephalopathy syndrome in eclamptic and pre-eclamptic patients with neurological symptoms

Michinori Mayama, MD, Kaname Uno, MD, Sho Tano, MD, Masato Yoshihara, MD, Mayu Ukai, MD, Yasuyuki Kishigami, MD, Yasuhiro Ito, MD, PhD, Hidenori Oguchi, MD, PhD

PII: S0002-9378(16)00339-2

DOI: [10.1016/j.ajog.2016.02.039](https://doi.org/10.1016/j.ajog.2016.02.039)

Reference: YMOB 10963

To appear in: *American Journal of Obstetrics and Gynecology*

Received Date: 1 December 2015

Revised Date: 28 January 2016

Accepted Date: 16 February 2016

Please cite this article as: Mayama M, Uno K, Tano S, Yoshihara M, Ukai M, Kishigami Y, Ito Y, Oguchi H, Incidence of posterior reversible encephalopathy syndrome in eclamptic and pre-eclamptic patients with neurological symptoms, *American Journal of Obstetrics and Gynecology* (2016), doi: 10.1016/j.ajog.2016.02.039.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Incidence of posterior reversible encephalopathy syndrome in eclamptic and

ACCEPTED MANUSCRIPT

pre-eclamptic patients with neurological symptoms

Michinori MAYAMA^{1,*}, MD; Kaname UNO¹, MD; Sho TANO¹, MD; Masato

YOSHIHARA¹, MD; Mayu UKAI¹, MD; Yasuyuki KISHIGAMI¹, MD; Yasuhiro ITO²,

MD, PhD; Hidenori OGUCHI¹, MD, PhD

¹ Department of Obstetrics, Perinatal Medical Center, TOYOTA Memorial Hospital,
Toyota, Aichi, Japan

² Department of Neurology, TOYOTA Memorial Hospital, Toyota, Aichi, Japan

Disclosure: The authors received no financial support for this work and report no conflict
of interest

*** Corresponding author:**

Michinori Mayama

Department of Obstetrics, Perinatal Medical Center, Toyota Memorial Hospital

20-2, Misato-cho, Toyota, Aichi, Japan

Phone: +81-90-9527-4848

Fax: +81-565-24-7170

E-mail: marimo.mayama@gmail.com

Word count: Abstract: 354 words; main text: 1840 words

Condensation:

Posterior reversible encephalopathy syndrome was detected in 92.3% eclamptic and 19.2% pre-eclamptic patients with neurological symptoms, and these patients had similar clinical and radiological backgrounds.

Download English Version:

<https://daneshyari.com/en/article/6143677>

Download Persian Version:

<https://daneshyari.com/article/6143677>

[Daneshyari.com](https://daneshyari.com)