### Accepted Manuscript

Two-group time-to-event continual reassessment method using likelihood estimation


Amber Salter, John O'Quigley, Gary R. Cutter, Inmaculada B. Aban

PII: S1551-7144(15)30091-4 DOI: doi: 10.1016/j.cct.2015.09.016

Reference: CONCLI 1282

To appear in: Contemporary Clinical Trials

Received date: 16 June 2015 Revised date: 18 September 2015 Accepted date: 21 September 2015



Please cite this article as: Salter Amber, O'Quigley John, Cutter Gary R., Aban Inmaculada B., Two-group time-to-event continual reassessment method using likelihood estimation, *Contemporary Clinical Trials* (2015), doi: 10.1016/j.cct.2015.09.016

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## **ACCEPTED MANUSCRIPT**

# Two-Group Time-To-Event Continual Reassessment Method Using Likelihood Estimation

Amber Salter, PhD1, John O'Quigley, PhD2, Gary R. Cutter, PhD1 Inmaculada B. Aban, PhD1

<sup>1</sup> Department of Biostatistics, University of Alabama at Birmingham, Birmingham, AL USA

<sup>2</sup> Université Pierre et Marrie Curie - Paris VI, 75005 Paris, France

Corresponding Author:

Amber Salter

University of Alabama at Birmingham

School of Public Health

1665 University Blvd. Room 327

Birmingham, AL 35294-0022

Phone: 205 934-6725

asalter@uab.edu

Keywords:dose finding, phase I trial, patient heterogeneity, continual reassessment method, time-to-event, maximum likelihood

#### Download English Version:

# https://daneshyari.com/en/article/6150557

Download Persian Version:

https://daneshyari.com/article/6150557

<u>Daneshyari.com</u>