



## Effects of fear of falling and activity restriction on normal and dual task walking in community dwelling older adults

Orna A. Donoghue\*, Hilary Cronin, George M. Savva, Claire O'Regan, Rose Anne Kenny

*The Irish Longitudinal Study on Ageing (TILDA), Trinity College Dublin, Ireland*

### ARTICLE INFO

#### Article history:

Received 25 April 2012

Received in revised form 12 September 2012

Accepted 30 October 2012

#### Keywords:

Gait  
Anxiety  
Falls  
Ageing  
Balance

### ABSTRACT

Fear of falling (FOF) is associated with poor physical and psychosocial health and can have debilitating consequences especially when it leads to activity restriction. This study examined whether normal and dual task gait disruptions were independently associated with FOF and activity restriction or if they were fully explained by impaired health status. Data was obtained from The Irish Longitudinal Study on Ageing (TILDA). Community dwelling adults  $\geq 65$  years, with a Mini-Mental State Examination score  $\geq 18$  and who completed a gait assessment ( $n = 1307$ ) were divided into three groups: no FOF, FOF but no activity restriction (FOF-NAR), FOF with activity restriction (FOF-AR). Physical, psychosocial and cognitive measures were obtained and gait characteristics were assessed using a GAITrite<sup>®</sup> mat during normal and dual task (cognitive) walking. After adjusting for sociodemographics, physical, mental and cognitive health, FOF was associated with reduced gait speed and stride length and increased double support phase and step width in normal and dual task conditions; these changes were most pronounced in those who restrict activities as a result of FOF. These gait changes may be associated with an increased fall risk, however some changes especially increased step width may also reflect positive, compensatory adaptations to FOF. The results also highlight the importance of treating underlying health impairments and preventing the transition from FOF to activity restriction.

© 2012 Elsevier B.V. All rights reserved.

### 1. Introduction

Fear of falling (FOF) is common in older adults, particularly females and is associated with a history of falls, gait and balance disorders, depression, poor self-rated health and reduced social activity [1–5]. It can affect an individual in two ways. The first is increased awareness leading to more cautious behaviour and positive fall prevention strategies. The second is avoidance of activities which can lead to social isolation, physical deconditioning, reduced functional capacity and frailty which can increase the risk of falls and exacerbate FOF [1,5–7].

Despite the strong association between falls and gait, limited research has examined independent associations between FOF and gait. Existing studies have found reduced gait speed [8–10], reduced step or stride length [9,11], increased step width and increased double support phase (DSP) [9] in those with FOF but similar changes are observed with increasing age [12,13], depression [14] and with a history of falls [15], all of which are also associated with FOF. RoCHAT et al. [16] found similar gait

changes between groups with no FOF, FOF alone and FOF with activity restriction but while the differences between the no FOF and FOF alone groups were fully explained by confounding variables, those with FOF and activity restriction had decreased gait performance and increased gait variability independent of health and functional status.

Gage et al. [17] suggested that anxiety associated with FOF may reduce the cognitive resources available for gait and balance control which may lead to gait disruptions. As dual task walking requires greater attention than normal walking, greater disruptions would be expected in dual task conditions. To date, only one study has examined dual task gait in older adults with and without FOF [10]. Reelick et al. [10] reported reduced velocity which explained increased gait variability among those with FOF but found no effect of FOF on trunk sway or cognitive task performance.

Gait disturbances are modifiable risk factors for falls, functional decline and other adverse health outcomes [18] and so it is important to understand how FOF affects gait. Adjusting for known confounders such as socio-demographics, depression, co-morbidity and cognitive function will also help determine if gait changes associated with FOF can be attributed to differences in physical, mental or cognitive health. Therefore, there were two aims in this study: (i) to examine the relationship between FOF, activity

\* Corresponding author at: The Irish Longitudinal Study on Ageing (TILDA), Lincoln Gate, Trinity College Dublin, Dublin 2, Ireland. Tel.: +353 1 896 4391; fax: +353 1 896 2451.

E-mail address: [odonogh@tcd.ie](mailto:odonogh@tcd.ie) (O.A. Donoghue).

restriction and gait characteristics in normal and dual task walking and (ii) to determine if these relationships persist after adjusting for potentially underlying factors.

## 2. Methods

### 2.1. Study design

The Irish Longitudinal Study on Ageing (TILDA) is a large prospective cohort study examining the social, economic and health circumstances of community dwelling older adults in Ireland. This study is based on the first wave of data collected from October 2009 to July 2011. Details of the study design and data collection are available elsewhere [19]. Briefly, the sampling frame is the Irish Geodirectory, a listing of all residential addresses in the Republic of Ireland. A clustered sample of addresses was chosen and all household residents aged 50 or over and their spouses/partners (of any age) were eligible to participate. Ethical approval was obtained from Trinity College Dublin Research Ethics Committee and written informed consent was obtained from 8504 participants.

### 2.2. Data collection

There were three parts to data collection: (i) a computer-assisted personal interview that included detailed questions on socio-demographics, wealth, health, lifestyle, social support and participation, use of health and social care and attitudes to ageing; (ii) a self-administered questionnaire; and (iii) a health assessment carried out by research nurses. Inclusion criteria for this study were age  $\geq 65$  years, no history of Parkinson's disease, Alzheimer's disease or dementia, Mini-Mental State Examination (MMSE) score of  $\geq 18$  and ability to complete the gait assessment without the use of physical aids and with a minimum of eight steps. Data was available on 1307 participants following these exclusions.

Sociodemographic variables such as age, sex and education (primary, secondary, tertiary) were obtained. The number of doctor diagnosed chronic conditions was obtained from the following list: heart attack or heart failure or angina, cataracts, stroke, diabetes, lung disease, asthma, arthritis, osteoporosis, cancer, peptic ulcer, hip fracture, hypertension and high cholesterol. Grip strength was assessed using a Baseline<sup>®</sup> hydraulic hand dynamometer. Two trials were averaged on each hand and the highest value was used in the analysis. Depression was assessed using the 20-item Centre for Epidemiological Studies Depression (CES-D) scale.

Fear of falling was assessed with the question "Are you afraid of falling?". Those that answered 'Yes' were asked "Do you ever limit your activities, for example, what you do or where you go, because you are afraid of falling?". This allowed participants to be classified into three groups – no FOF, FOF but no activity restrictions (FOF-NAR) and FOF with activity restrictions (FOF-AR). It is important to note that this activity restriction question addresses both physical and social activity limitations. The number of falls in the previous year was recorded and categorised into 0, 1 or 2+ falls.

Global cognition was assessed with the MMSE while executive function and visual scanning were assessed using the Colour Trails Test [20]. In Colour Trail 1, participants drew a line connecting circles numbered 1–25 in consecutive order. In Colour Trail 2, participants connected these numbered circles but alternated between pink and yellow circles, i.e. pink 1, yellow 2, pink 3 etc. Each test was timed and the difference (trail 1 – trail 2) was used as a measure of executive function. A choice reaction time (CRT) test used a computer based program to assess concentration and speed of processing. Participants depressed a button on the keyboard to initiate the test and maintained this position while they waited for a Yes/No stimulus to appear on screen. When the stimulus appeared, the participant released the button and pressed another corresponding Yes/No button on the keyboard. The time from appearance of the stimulus to pressing this button was the total reaction time.

Gait assessment took place using a 4.88 m computerised walkway with embedded pressure sensors (GAITRite<sup>®</sup>, CIR Systems Inc., New York, USA). Participants performed two walks at their normal pace followed by two walks under cognitive dual task conditions. The cognitive task was to recite alternate letters of the alphabet (A–C–E, etc.). The number of letters attempted and the number correct were recorded. Participants started and finished 2.5 m before and 2 m after the walkway to allow for acceleration and deceleration. Mean gait speed, stride length, step width and DSP were obtained along with standard deviations (SDs) of stride length and stride time. As the SDs were related to the mean values for each variable [21], coefficient of variation (CV) was calculated ( $SD/mean \times 100$ ) and used to indicate stride-to-stride variability in stride length and stride time. These gait variables were chosen for analysis because of their associations with adverse outcomes such as falls in older adults [22]. While step parameters may vary between right and left sides, stride parameters include data for the left step followed by the right step (or vice versa). Therefore, there is little variation between right and left stride-based data so we combined data for the right and left sides for analysis. Based on a minimum of eight steps (range 8–22 steps for normal walk; 8–26 steps for cognitive walk), this provided a minimum of six strides for analysis.

### 2.3. Statistical analysis

Statistical analysis was conducted using Stata v12 (StataCorp LP, TX, USA). All analyses were weighted with respect to age, sex and education to the Quarterly

National Household Survey (2010) to ensure that data were nationally representative. Data were further weighted by health status and socio-demographic factors to account for those who did not attend a health assessment. Baseline characteristics of each group were compared using regression analysis for continuous variables and chi-squared tests for categorical variables. Mean gait variables were normally distributed and regression analysis was used to examine differences between groups using the no FOF group as the reference. Gait variability measures were positively skewed so quantile regression was used to estimate the differences in the medians and minimise the influence of high values. An additional advantage of this approach over transformations is that it keeps the coefficients in their original units allowing a more intuitive interpretation. For each gait parameter, two models were estimated. Model 1 adjusted for age, sex, height and education while Model 2 additionally adjusted for chronic conditions, number of medications, grip strength, CES-D score, total reaction time, Colour Trail time difference and number of falls in the previous year. Complete covariate data was available for 1207 (92.3%) and 1196 (91.5%) participants for the normal and dual task walk respectively. Complete case analysis was used for multivariate analyses, i.e. all missing data were treated as missing at random. Significance level was set at  $p < 0.05$ .

## 3. Results

In this population based sample (mean age 71.3 years; range 65–93 years), 73.5% of participants ( $n = 961$ ) reported no FOF, 19.1% reported FOF but no activity restriction ( $n = 250$ ) while 7.5% reported FOF with activity restriction ( $n = 96$ ). Baseline characteristics for each group are provided in Table 1. Both FOF groups were older, more likely to be female and a faller and had poorer physical and mental health than the no FOF group ( $p < 0.05$ ). There was no difference between groups in global cognition (MMSE) but the FOF-AR group had a longer choice reaction time and Colour Trail time difference than the no FOF group ( $p < 0.05$ ).

Table 2 outlines the gait characteristics for the normal and dual task walk for each group. In the normal walk, a progressive deterioration in gait characteristics was observed moving from the no FOF to FOF-NAR and FOF-AR groups. Specifically, both FOF groups demonstrated reduced gait speed and stride length and increased DSP and stride time variability ( $p < 0.05$ ). Similar univariate differences in gait speed, stride length and DSP were observed in the dual task walk ( $p < 0.05$ ) while gait variability was non-significantly higher in both FOF groups. In the cognitive task, both groups attempted and recited correctly more letters than the no FOF group, but there was no difference between groups in the percentage of letters recited correctly.

Table 3 provides the coefficients from the multivariate regression analysis illustrating the magnitude of differences between groups using the no FOF group as the reference. After adjusting for age, sex, height and education, both FOF groups walked more slowly, took shorter steps and had increased step width and DSP compared to the no FOF group in both normal and dual task walking conditions ( $p < 0.05$ ). Additionally adjusting for chronic conditions, number of medications, grip strength, CES-D score, total reaction time, Colour Trail time difference and number of falls reduced the coefficients for all variables but the differences observed in Model 1 remained statistically significant for almost all mean gait variables. The reduction in gait speed for the FOF-AR group ( $p = 0.051$ ) and stride length for the FOF-NAR group ( $p = 0.057$ ) in dual task conditions were also very close to the significance level chosen.

In summary, FOF with and without activity restriction was independently associated with slower gait speed, shorter stride length and increased step width and DSP both in normal and dual task walking. Effects on DSP and step width were similar in both groups however, reductions in gait speed and stride length were more pronounced in the FOF-AR group. There was no difference in stride length variability or stride time variability between groups in either normal or dual task walking ( $p > 0.05$ ).

Download English Version:

<https://daneshyari.com/en/article/6207520>

Download Persian Version:

<https://daneshyari.com/article/6207520>

[Daneshyari.com](https://daneshyari.com)