

Case Report

# Anterior retroperitoneal approach for removal of L5-S1 foraminal nerve sheath tumor—case report

Kevin Phan, BS<sup>a,b,c</sup>, Ralph J. Mobbs, MBBS MS FRACS<sup>a,b,c,\*</sup>

<sup>a</sup>NeuroSpine Surgery Research Group (NSURG), Prince of Wales Private Hospital, Randwick, Sydney, Australia

<sup>b</sup>NeuroSpine Clinic, Prince of Wales Private Hospital, Randwick, Sydney, Australia

<sup>c</sup>University of New South Wales, Sydney, Australia

Received 23 May 2015; revised 24 November 2015; accepted 2 December 2015

## Abstract

**BACKGROUND CONTEXT:** Extradural lumbar schwannomas are a rare form of nerve sheath tumors (NSTs). The typical management approach for extradural foraminal NSTs is total gross resection, which involves a midline incision and muscle exposure, followed by laminectomy and facetectomy to access the tumor for resection. Following tumor removal, spinal fusion is often indicated to reduce postoperative deformity, pain, and neurologic deficits.

**PURPOSE:** We report the case of a 34-year-old woman who presented with a 2-year history of progressive dysesthesia and left foot drop. Magnetic resonance imaging revealed a lesion in the lateral L5/S1 foramen. A novel anterior-retroperitoneal approach was used to access the tumor, via muscle splitting, retraction of peritoneum medially and psoas muscle or iliac vessels laterally.

**STUDY DESIGN/SETTING:** This study is a case report of a novel approach for extradural lumbar schwannomas.

**METHODS:** The methods involve a description of the approach and reporting of clinical findings.

**RESULTS:** The schwannoma was successfully resected without requiring additional fusion surgery. The patient recovered uneventfully and was discharged on day 2 post operation.

**CONCLUSION:** We propose that the anterior-retroperitoneal approach is a viable technique for resection of lumbar foraminal NSTs without the need for fusion surgery. © 2015 Elsevier Inc. All rights reserved.

## Keywords:

Anterior; Fusion; Laminectomy; Lumbar; Nerve sheath tumor; Retroperitoneal; Schwannoma

## Introduction

Neoplasms of the spinal canal comprise a range of tumors which may involve the spinal cord, theca, and nerves. Nerve sheath tumors (NSTs) are the most frequent form, making up 25%–35% of primary spinal neoplasms in the Western population. These tumors can be classed according to their

location, including intradural, combined intradural-extradural, and purely extradural [1]. In the intramedullary location, astrocytomas and ependymomas are more common tumors, compared with extramedullary tumors which are more likely NSTs such as schwannomas, neurofibromas, and meningiomas.

Spinal extradural foraminal NSTs, which are more likely found in the cervical and thoracic region compared with the lumbar spine, are a rare form of neoplasms [1]. The typical management approach for extradural foraminal NSTs is total gross resection, which involves a midline incision and muscle exposure, followed by laminectomy and facetectomy to access the tumor for resection. Following tumor removal, spinal fusion is often indicated to reduce postoperative deformity, pain, and neurologic deficits [2–4]. Various reports have demonstrated the safety and efficacy of mini-open tubular removal of these lesions, a posterior approach [3,5,6]. We report a case of removal of an L5/S1 foraminal schwannoma through an anterior, retroperitoneal exposure.

FDA device/drug status: Not applicable.

Author disclosures: **KP:** Nothing to disclose. **RJM:** Royalties: Stryker Spine (F); Stocks: Medtronic USA (D), J&J USA (D); Speaking/Teaching Arrangements: Stryker Spine and Synthes (B); Trips/Travel: Orthotec Australia (B, Grant); Research Support: Cerapedics (C, Grant), outside the submitted work.

The disclosure key can be found on the Table of Contents and at [www.TheSpineJournalOnline.com](http://www.TheSpineJournalOnline.com).

\* Corresponding author. Neuro Spine Clinic, Suite 7a, Level 7 Prince of Wales Private Hospital, Barker Street, Randwick, New South Wales 2031, Australia. Tel.: +61 2 9650 4766.

E-mail address: [ralphmobbs@hotmail.com](mailto:ralphmobbs@hotmail.com) (R.J. Mobbs)

## Case report

A 34-year-old healthy woman presented with a 2-year history of progressive dysesthesia and foot drop on the left side. Magnetic resonance imaging (MRI) scan revealed a lesion located in the lateral aspect of the left L5/S1 foramen, with deflection of the exiting L5 nerve root laterally. Nerve conduction studies confirmed an L5 nerve root lesion, with normal conduction velocities of the common peroneal nerve.

The patient was initially booked for a posterior approach total facetectomy, resection of the lesion, and then transforaminal lumbar interbody fusion with pedicle screw fixation at an outside institution. The primary author (RJM) adopted a non-fusion approach of an anterior retroperitoneal approach to access the lesion for resection.

Under general anesthesia, a linear incision medial to the anterior superior iliac spine (Fig. 1) and lateral to the rectus abdominus was performed. Using a muscle splitting approach through the external and internal oblique muscles, the retroperitoneal plane was exposed. An Alexis retractor (Applied Medical Resources Corporation, Orange County, CA, USA) was used to open the corridor to the psoas muscle. A SynFrame retractor (SYNTHES Spine, Paoli, PA, USA) was inserted to retract the peritoneum medially and the psoas or common iliac vessels laterally to expose the L5/S1 foramen from an anterior direction (Fig. 2).

The lesion was easily located at the L5/S1 foramen with the L5 nerve deflected laterally (Fig. 3). Using microneurosurgical technique, the false capsule of the NST was incised and opened with complete resection of the lesion. Histopathology revealed a schwannoma with no atypical features.

The patient recovered uneventfully and was discharged on day 2 post operation. The patient re-commenced her teaching




Fig. 2. Anatomy of the anterior exposure. The L5 nerve is positioned lateral to the nerve sheath tumor (white arrows).

duties at 2 weeks following surgery. Imaging at 9 months' follow-up demonstrated complete resection of the NST (Fig. 4).

## Comment

Extradural foraminal schwannomas at the lumbar spine region are an exceptionally rare type of spinal NST. Prior studies have reported prevalence for lumbar extradural schwannomas of 0.7%–4.2% in all extradural schwannomas [1,3]. The mainstay treatment approach involves total resection of the lesion [1,7], via a midline incision, bilateral muscle retraction from posterior spinous structures, laminectomy, facetectomy on the side of the tumor, followed by motion segment stabilization via fusion [1,8]. However, prior reports have used a posterior approach for tumor resection. To the best of our knowledge, this is the first case using an anterior retroperitoneal exposure for resection of a lumbar extradural foraminal schwannoma.

The posterior approach has traditionally been used for surgical access of the spine, particularly in fusion, discectomy, and pedicle screw instrumentation cases. The posterior approach allows access to posterior structures such as the lamina, ligamentum flavum, and facet joints. It also facilitates simultaneous fusion and posterior instrumentation if required in the same exposure, thereby reducing surgical trauma and dissection of soft tissues and paraspinal structures [9–12]. However, multiple studies have also emphasized several disadvantages associated with the posterior approach for spinal canal or foraminal access and tumor resection. First, the posterior or transforaminal approach may be associated with longer operation duration, because of the requirement of dissection of paraspinal structures, laminectomy, and facetectomy [13]. Second, the posterior approach may require prolonged retraction of nerve roots and opening of neural foramen for schwannoma access. Excess retraction force may lead to exacerbated dural injury and nerve root pain [9–12,14]. Third, another major concern is that the posterior approach inherently




Fig. 1. Axial T2WI magnetic resonance imaging (MRI): Foraminial schwannoma (asterisk). L5 nerve pushed laterally by lesion (white arrow). (A) Potential posterior approach with facetectomy to resect lesion. (B) Potential anterior approach to resect lesion. Inset demonstrates location and size of incision for the described approach.

Download English Version:

<https://daneshyari.com/en/article/6211778>

Download Persian Version:

<https://daneshyari.com/article/6211778>

[Daneshyari.com](https://daneshyari.com)