

Clinical Study

Hypertrophy of the anterior arch of the atlas associated with congenital nonunion of the posterior arch: a retrospective case-control study

Michael Jin, MD, Mariet Asadoorian, MD, Lucas P. Hiller, MD, Tudor H. Hughes, MD*

Department of Radiology, University of California San Diego School of Medicine, 200 W. Arbor Drive, San Diego, CA 92103, USA

Received 26 May 2012; revised 21 June 2013; accepted 25 July 2013

Abstract

BACKGROUND CONTEXT: Nonunion of the posterior arch of the atlas is an uncommon but normal developmental variant. It is usually asymptomatic in the patient but may be associated with greater incidence of fracture because of increased stress on the anterior arch.

PURPOSE: We sought to determine whether anterior arch hypertrophy is present in cases of congenital nonunion of the posterior arch of the atlas.

STUDY DESIGN/SETTING: A retrospective analysis of 1 year (February 2005–January 2006) of computed tomography cervical spine studies requested by the University of California San Diego Medical Center Trauma Department was undertaken.

PATIENT SAMPLE: All patients matching the search criteria (see *Study design*) were included.

OUTCOME MEASURES: Area density product, defined as the midline cross-sectional area of the anterior arch on sagittal reformat multiplied by the average areal radiodensity in Hounsfield units (HU) as measured by two raters, was calculated for cases and controls.

METHODS: Cases of posterior arch nonunion were identified and matched to controls. The significance of differences in area density product between cases and controls were established by the Student *t* test. Interrater correlation was calculated.

RESULTS: Posterior arch nonunion was identified in 26 individuals (3.1% of 839 studies reviewed). Compared with age- and sex-matched controls, a 21% increase in area density product of the midline anterior arch was observed in posterior arch nonunion cases (773 HU-cm² in cases vs. 637 HU-cm² in controls; *p* < .001). This increase was attributable to a 21% increase in cross-sectional area (1.05 cm² in cases vs. 0.87 cm² in controls; *p* < .002). In contrast, there was no significant difference with regard to increased average radiodensity.

CONCLUSIONS: It has long been subjectively recognized but not objectively quantified, until the present study, that the anterior arch of the atlas is hypertrophied in cases of posterior arch nonunion. Anterior arch hypertrophy may represent an adaptive response to chronically elevated mechanical stress and loss of hoop strength in cases of posterior nonunion. © 2014 Elsevier Inc. All rights reserved.

Keywords:

Cervical atlas; Congenital; Spine; Spinal fractures; Radiology; Retrospective Study

Introduction

The atlas (C1) is the topmost vertebra. Its ring-like structure is composed of an anterior arch, two lateral masses,

and a posterior arch. It supports the skull superiorly, and its odontoid articulation with the axis (C2) affords the cervical spine its liberal range of rotational motion.

During development, the atlas classically ossifies in three centers. The two lateral centers appear in the seventh week and form the lateral masses, which also extend posteriorly to form the incipient posterior arch. At birth, a narrow segment of cartilage separates the two posterior hemi-arches; fusion occurs at age 3 to 5 years. The primordial anterior arch is fibrocartilaginous at birth, the developmental remnant of the hypochordal bow. The third center of

FDA device/drug status: Not applicable.Author disclosures: **MJ:** Nothing to disclose. **MA:** Nothing to disclose.**LPH:** Nothing to disclose. **THH:** Nothing to disclose.

* Corresponding author. Department of Radiology, University of California San Diego School of Medicine, 200 W. Arbor Drive, San Diego, CA 92103-8756, USA. Tel.: (619) 543-6607; fax: (619) 543-3781.

E-mail address: thughes@ucsd.edu (T.H. Hughes)

ossification appears anteriorly at the end of the first year of life, eventually forming the mature osseous anterior arch, which fuses with the lateral masses at age 5 to 9 years.

Nonunion of the posterior arch of the atlas is an uncommon but normal developmental variant. Failure of midline fusion of the two posterior hemiarches accounts for the majority of such defects, with prevalence estimated at 1% to 4.3% of the population based on cadaveric dissections [1] and retrospective survey of neck and cervical spine computed tomography (CT) scans [2]. Non-midline nonunion of the posterior arch is less common, and anterior arch nonunion is exceedingly rare [3]. In the emergency department, it is important to distinguish variant anatomy from acute trauma [4–9], that is, Jefferson fractures [10–12] and odontoid fractures [13,14], based on criteria such as cortication, soft-tissue swelling, lateral offset [12], and/or cervical spine instability during clinical examination [15].

Most atlantal defects are present in asymptomatic patients and represent normal variants [3]. Less commonly, posterior arch abnormalities can have severe neurologic deficits when associated with conditions such as Arnold-Chiari malformation, gonadal dysgenesis, Klippel-Feil, Turner, and Down syndromes [16]. Atlantal defects are associated with greater degree of atlantoaxial instability and susceptibility to injury after minor head or neck trauma [17–19]. It has also been suggested that atlantal defects are associated with compensatory changes such as hypertrophy of the anterior arch of C1, hypertrophy of spinous process of C2, and hypertrophic downward projection of the posterior border of the foramen magnum [17].

We hypothesized that in cases of congenital nonunion of the posterior arch, additional mechanical stress would be exerted on the anterior arch; hence, the anterior arch would adaptively remodel in these individuals and achieve greater cross-sectional area, greater radiodensity, or both, compared with age- and sex-matched control patients.

Material and methods

Institutional review board approval was obtained for retrospective review of 1 year of CT cervical spine studies ordered by the University of California San Diego Medical Center Trauma Department from February 2005 to January 2006 inclusive, using Agfa picture archive communications systems (ie, PACS) search criteria. Initial exclusion criteria were: Jefferson fracture, digitized outside images, and lack of age identifier.

Age, sex, and the presence of an open or closed posterior arch were recorded for each study. If a posterior arch cleft was present, area density product (ADP) was calculated on standard 1-mm thickness midline sagittal reformat image. ADP was defined as the midline cross-sectional area of the anterior arch on sagittal reformat image, measured by free-form mark up in Agfa PACS and multiplied by the average areal radiodensity in Hounsfield units (HU) measured on this

same image that had been magnified 15 times on a window width/level of 2000/500. These same measurements and calculations were then performed on normal age- and sex-matched control patients who had no cervical spine abnormalities. If anterior arch ADP could not technically be obtained, the case-control pair was excluded. Means of anterior arch midline cross-sectional area, average areal radiodensity, and ADP values for cases and control patients were compared. Statistical significance was established by a paired Student *t* test with a two-tailed distribution. All measurements were performed by two radiologists, who were both board-certified and fellowship-trained in musculoskeletal radiology. Interrater correlation was calculated.

Results

By search criteria, 847 CT scans were found. Eight scans met initial exclusion criteria. In total, 839 scans comprising 623 male and 216 female patients were reviewed. The ages ranged from 11 to 99 years; average age was 41 years.

Nonunion of the posterior arch (Fig. 1) was noted in 26 individuals (3.1%) with an age range of 17 to 87 years, and an average age of 43 years. Twenty of these patients presented with a simple cleft; the remaining six individuals comprised 1 split atlas, 1 near-complete occipitoatlas fusion, and 4 unilateral or bilateral occipitoatlas fusions. The first two individuals were excluded from analysis because a midline anterior arch cross-sectional area could not be measured. One case of anterior arch stress fracture was observed; posterior nonunion was noted in this same individual (Fig. 2).

Anterior arch ADP means, cross-sectional area, and average areal radiodensity, averaged across both raters, were




Fig. 1. Axial view of CT cervical spine at the level of C1: nonunion of the posterior arch. CT, computed tomography.

Download English Version:

<https://daneshyari.com/en/article/6212348>

Download Persian Version:

<https://daneshyari.com/article/6212348>

[Daneshyari.com](https://daneshyari.com)