



Gastroesophageal reflux as a predictor of complications after adenotonsillectomy in young children[☆]



Michael E. McCormick^{a,*}, Anthony Sheyn^b, Michael Hauptert^c, Adam J. Folbe^{b,c}

^a Department of Otolaryngology and Communication Sciences, Division of Pediatric Otolaryngology, Medical College of Wisconsin, Milwaukee, WI, United States

^b Department of Otolaryngology – Head & Neck Surgery, Wayne State University School of Medicine, Detroit, MI, United States

^c Department of Otolaryngology, Children's Hospital of Michigan, Detroit, MI, United States

ARTICLE INFO

Article history:

Received 14 June 2013

Received in revised form 9 July 2013

Accepted 16 July 2013

Available online 9 August 2013

Keywords:

Tonsillectomy

Adenoidectomy

Complications

Airway obstruction

Young children

Gastroesophageal reflux

ABSTRACT

Objective: To examine characteristics of young children with gastroesophageal reflux (GER) who experienced complications within the first 24 h after adenotonsillectomy.

Study design: Subset analysis of a larger retrospective cohort.

Methods: A retrospective chart review was performed at a tertiary care children's hospital. Consecutive records of children 3 years old and younger undergoing adenotonsillectomy (AT) over a 5-year period were reviewed. Children with a clinical history of GER were selected for the study.

Results: 993 children were included in the initial analysis, and GER was found to be a significant independent variable predictive of early complications. 81 children with a history of GER were included in this study and 8 (9.9%) were found to have experienced complications within the first 24 h. Six of the complications were airway-related; two required re-intubation within the first 24 h. All 8 children with complications had symptoms of sleep-disordered breathing and two had documented severe obstructive sleep apnea (AHI 18.6 and 27.2). Seven children had other risk factors for complications after AT. Eighteen (22%) children had a prolonged length of stay (range 2–7 days); additional risk factors were present in these patients as well.

Conclusions: Knowledge of risk factors for complications following adenotonsillectomy is critical for identifying at-risk patients that may warrant closer post-operative observation. GER has been previously identified as a risk factor for complications in young children. Upon closer analysis, young children with GER who have other known risk factors may be at a further increased risk for airway complications and prolonged hospitalization. Parents of these children can be counseled on the post-operative risks and the possibility of a longer hospitalization.

© 2013 Published by Elsevier Ireland Ltd.

1. Introduction

Tonsillectomy with or without adenoidectomy (AT) is one of the most common outpatient procedures performed in the United States [1,2]. The overall complication rate (early and late) for AT is estimated to be between 2 and 10% [3]; early post-operative complications include pain, hemorrhage, nausea, vomiting, dehydration, and post-obstructive pulmonary edema. Known risk factors for complications after AT include obstructive sleep apnea syndrome, craniofacial abnormalities, Down's syndrome, cerebral

palsy, obesity, bleeding disorders, and age younger than 3 years old [1,4–10].

In a recent retrospective review of 993 patients age 3 or younger who had undergone AT, we identified a number of independent variables predictive of early complications using multivariate analysis [11]. Consistent with previously published studies were the identification of prematurity and cardiovascular anomalies as predictors of early complications and also the identification of younger age (less than 3) as a predictor of airway complications. A unique and novel finding of this study was the identification of gastroesophageal reflux as an independent predictor of early complications.

Gastroesophageal reflux (GER) has gained increased interest in the pediatric otolaryngology community over the last two decades [12–15]. With the increasing role of GER in otolaryngologic diseases, as well as our recent association of GER with early post-operative complications after AT, we further analyzed the subset of

[☆] Poster presentation at American Society of Pediatric Otolaryngology, San Diego, CA, April 2012.

* Corresponding author at: Division of Pediatric Otolaryngology, Children's Hospital of Wisconsin, 9000 W. Wisconsin Avenue, Suite 540, Milwaukee, WI 53226, United States. Tel.: +1 414 266 6393; fax: +1 414 266 2693.

E-mail address: mmccormick@mcw.edu (M.E. McCormick).

children 3 years old and younger with GER who underwent adenotonsillectomy.

2. Methods

We initially performed a retrospective chart review with Institutional Review Board approval of all children 3 years old and younger undergoing adenotonsillectomy at the Children's Hospital of Michigan between July 1, 2005 and June 30, 2010. Charts were reviewed for demographic data, pre-operative symptoms and conditions, and postoperative hospitalization details, including time to adequate oral (PO) intake, complications, and length of stay (LOS). Comorbid conditions, including gastroesophageal reflux, were included when identified during historical chart review of our institution's records alone. Objective diagnostic testing for GER was not always present in our institution's records, and therefore was not included in the analysis.

Primary outcomes for the initial evaluation included: total complications on post-operative days (POD) 0–1, airway complications on POD 0–1, and length of stay (LOS). Our initial analysis using multivariate logistic regression models has previously been published [11]. Gastroesophageal reflux was identified as an independent predictor of both overall complications and prolonged length of stay (≥ 2 days). A subset analysis was performed and closely examined all children in the cohort with a history of GER in an effort to further characterize these children and their post-operative course.

3. Results

Initially, 993 children were included in the primary analysis and their characteristics have been previously published [11]. Of this original cohort, eighty-one were found to have had a history of gastroesophageal reflux. Of those 81 children, 8 (9.9%) experienced complications within the first 24 h post-operatively. Their characteristics can be seen in Tables 1 and 2. Almost all of the

children in both groups had symptoms of sleep-disordered breathing (100% of children with complications, 93.2% of children without complications). Children with GER who experienced complications were younger (median age 26.0 months vs. 31.5 months) and had more associated co-morbid conditions (Fig. 1). Additionally, more severe obstructive sleep apnea (OSA) was seen in children with complications (median AHI 22.9 vs. 12.1).

The average length of stay (LOS) of the 81 children with GER was 1.09 days; children without GER went home after 0.84 days. Eighteen (22.2%) children with GER required hospitalizations longer than the expected 24 h. Longer LOS (≥ 2 days) was seen more often in younger children (median age 26.1 months vs. 32.3 months) and children with cardiovascular anomalies, abnormal airways, or craniofacial syndromes (Fig. 2). In addition, children with more severe OSA on pre-operative polysomnography experienced longer hospitalizations (median AHI 21.2 vs. 8.4).

Post-operative complications in children with gastroesophageal reflux occurred in the presence of other known risk factors in all 8 children. Two patient examples are provided:

Patient #4 was a 23-month old male with a history of prematurity, Beckwith–Wiedemann syndrome, and gastroesophageal reflux. He had previously undergone an adenoidectomy and glossectomy but he continued to have heroic snoring, gasping, and witnessed apneas. Polysomnography demonstrated an apnea–hypopnea index of 27.2. He underwent tonsillectomy with concurrent direct laryngoscopy, bronchoscopy, and esophagoscopy. He developed respiratory distress and inspiratory stridor and was transferred to the pediatric intensive care unit (PICU) for monitoring. He did well overnight and was transferred to the surgical floor where he was discharged on post-operative day #2.

Patient #6 was an 18-month old male who presented with symptoms of snoring and witnessed apneas. He also had chronic otitis media with effusion, reflux, and a history of aortic stenosis. He underwent adenotonsillectomy and bilateral

Table 1
Demographic and pre-operative characteristics of patients who experienced complications.

Patient #	Age (months)	Gender	SDB	Asthma	Prematurity	Cardiovascular anomalies	Seizures	Airway findings	PSG (AHI)
1	14	F	Y	Y	N	N	N	N	N
2	43	M	Y	N	N	N	N	N	N
3	35	F	Y	N	N	N	Y	Y ^c	Y (18.6)
4	23	M	Y	N	Y	Y ^a	N	Y ^d	Y (27.2)
5	27	M	Y	N	Y	N	N	N	N
6	18	M	Y	N	N	Y ^b	N	N	N
7	36	M	Y	Y	Y	N	N	N	N
8	25	F	Y	N	N	N	N	N	N
TOTAL	27.6	5M/3F	100%	25%	37.5%	25%	12.5%	25%	25%

^a Atrial septal defect.

^b Aortic stenosis.

^c Laryngeal cleft.

^d Beckwith–Wiedemann syndrome.

Table 2
Operative and post-operative details of patients who experienced complications.

Patient #	Primary procedure	Other procedures	Admit location	LOS	Details
1	AT	Y ^a	PICU	1	Desaturations requiring High-Flow Nasal Cannula
2	AT	Y ^a	FLOOR	2	Desaturations requiring non-rebreather mask, transferred to PICU
3	AT	Y ^a	PICU	2	Post-obstructive pulmonary edema
4	T	Y ^a	PICU	2	Desaturations and stridor in recovery unit, admitted to PICU for monitoring
5	AT	N	PICU	2	Desaturations and CO ₂ retention requiring re-intubation
6	AT	Y ^b	PICU	2	Desaturations and stridor in recovery unit requiring re-intubation
7	AT	N	HOME	0	Returned to hospital for dehydration on POD#1
8	AT	Y ^a	FLOOR	2	Primary post-tonsillectomy bleed in recovery unit requiring re-operation

^a Direct laryngoscopy, bronchoscopy, esophagoscopy.

^b Bilateral myringotomy and tympanostomy tube insertion.

Download English Version:

<https://daneshyari.com/en/article/6213546>

Download Persian Version:

<https://daneshyari.com/article/6213546>

[Daneshyari.com](https://daneshyari.com)