

Relationship between Sleep Position and Risk of Extreme Cardiorespiratory Events

George Lister, MD¹, Denis V. Rybin, MS², Theodore Colton, ScD³, Timothy C. Heeren, PhD⁴, Carl E. Hunt, MD⁵, Eve R. Colson, MD⁶, Marian Willinger, PhD⁷, and Michael J. Corwin, MD⁸, on behalf of the Collaborative Home Infant Monitoring Evaluation (CHIME) Study Group*

Objective To determine whether infants at sleep in the prone side positions are at higher risk for an extreme cardiorespiratory event compared with infants at sleep in the supine position.

Study design We used a case-control study to compare sleep position, determined with an accelerometer, in 116 infants during an extreme cardiorespiratory event with that in 231 matched control subjects (2 per case) who did not experience any extreme events during monitoring.

Results From calculation of adjusted ORs and 95% CIs, infants placed in the prone or side position were no more likely to experience an extreme cardiorespiratory event compared with infants at sleep in the supine position. We used conditional logistic regression to account for the matched design of the study and to adjust for potential confounders or effect-modifiers.

Conclusion These findings, coupled with our earlier observation that the peak incidence of severe cardiorespiratory events occurred before the peak incidence of sudden infant death syndrome, strongly suggest that the supine sleeping position decreases the risk of sudden infant death syndrome by mechanisms other than by decreasing extreme cardiorespiratory events detected by monitoring. (*J Pediatr* 2012;161:22-5).

Since 1994, and on the basis of a recommendation of the American Academy of Pediatrics,¹ there has been a national public education campaign to reduce the risk of sudden infant death syndrome (SIDS). Coincident with the reported reduction in prone sleeping in this country, there has been a substantial decrease in SIDS to approximately 50% of the rate per 1000 live births that was reported before the recommendation.²⁻⁴ Despite the success of this public health intervention, this observation immediately prompts the question of how prone sleep affects the risk of SIDS.

We previously conducted a National Institutes of Health-sponsored multicenter study of the usefulness of home monitoring in infants thought to be at increased risk for SIDS (the Collaborative Home Infant Monitoring Evaluation [CHIME]). Toward this end, we recorded cardiorespiratory data in 1070 infants for 700 000 hours to detect episodes of extremely prolonged apnea or bradycardia, because it had been presumed for many years that apnea or bradycardia was the prelude to sudden death.⁵ The “at-risk” infant groups enrolled in the CHIME study included infants born prematurely (<1750 g and ≤34 weeks at birth), siblings of SIDS victims, and infants with a history of an apparent life-threatening event, in addition to a group of healthy term infants. We reported that, when compared with healthy term infants, extreme events (EEs) of apnea or bradycardia were more likely to occur only in premature infants and only before 43 weeks postmenstrual age (PMA), well before the peak incidence of SIDS, especially in infants born full term.⁶

Although our earlier findings suggested that extreme cardiorespiratory events were not immediate precursors of SIDS, it is important to test the validity of that inference. Thus, if there is no relationship between EEs and non-supine sleep position, it would strongly suggest that supine position decreases the risk of SIDS by means other than decreasing extreme apnea or bradycardia. In contrast, if there is a strong relationship, this should prompt additional study to understand this precise mechanism. The central question we addressed was whether infants in the prone position or infants in side position are at higher risk for an extreme cardiorespiratory event compared with infants in the supine position. We had a unique opportunity to answer this question because the monitors had been equipped with a sensor to track infant position, rather than relying on the report of an observer.

From the ¹Department of Pediatrics, UTSW Medical School, Dallas, TX; ²Data Coordinating Center, ³Department of Epidemiology, and ⁴Department of Biostatistics, Boston University School of Public Health, Boston, MA; ⁵Department of Pediatrics, University of Toledo Health Sciences Center, Toledo, OH; ⁶Department of Pediatrics, Yale University School of Medicine, New Haven, CT; ⁷National Institute of Child Health and Human Development, National Institutes of Health, Bethesda, MD; and ⁸Departments of Pediatrics and Epidemiology, Boston University Schools of Medicine and Public Health, Boston, MA

*List of members of the CHIME Study Group is available at www.jpeds.com (Appendix).

Supported by the Eunice Kennedy Shriver National Institute of Child Health & Human Development of the National Institutes of Health HD (grants HD 29067, 29071, 28971, 29073, 29060, 20056, 34625, and 59207). The content is solely the responsibility of the authors and does not necessarily represent the official views of the Eunice Kennedy Shriver National Institute of Child Health & Human Development or the National Institutes of Health. The authors declare no conflicts of interest.

0022-3476/\$ - see front matter. Copyright © 2012 Mosby Inc. All rights reserved. 10.1016/j.jpeds.2012.01.035

CHIME	Collaborative Home Infant Monitoring Evaluation
EE	Extreme event
PMA	Postmenstrual age
SIDS	Sudden infant death syndrome

Methods

We designed a case-control study with infants previously enrolled in the CHIME study. We selected “cases” as infants who experienced at least one EE. For each case, we selected two control subjects from infants who did not experience any EE. For each of the cases, we assessed the infant’s position at a period immediately before the first extreme cardiorespiratory event; for each of the two matched control subjects, we assessed the infant’s position at a period comparable with the time of the case’s first extreme cardiorespiratory event. As reported previously,⁵ all home recordings were collected between May 1994 and February 1998, the institutional review board at each site approved the study (Appendix), and the parents of all subjects gave written informed consent.

The occurrence of EEs and the methods used to identify these EEs in the CHIME study were reported previously⁵ and were extensively validated to reduce the possibility of technical artifact.⁷ An EE was defined⁵ as: apnea ≥ 30 seconds or a heart rate < 60 bpm for ≥ 10 seconds (when < 44 weeks PMA) or < 50 bpm for ≥ 10 seconds (when ≥ 44 weeks PMA). On the basis of these selection criteria, the cases included all 116 infants in whom a total of 653 EEs occurred during the CHIME study. For this report, those infants who had an EE were selected and matched to the control infants as follows. An infant was selected on the basis of having the first EE within 180 days from the start of monitoring. This frame was used because it was the target duration of home monitoring in the CHIME study. When there was more than one EE, only the first event was used, because the first EE, when noticed by the caregiver, might have prompted a change in care of the infant and confounded assessment of the hypothesis. Furthermore, for a given infant, sleeping position rarely changed during a particular night or between nights. Thus, the position observed in subsequent events would not be independent of the position observed in the first event, so that inclusion of many events from the same infant would not substantially add to the power of our analyses.

For control infants, epochs were obtained from the 3-minute “non-event” recording that we obtained hourly in all infants as part of our study protocol.⁸

We matched control infants with case infants in these ways: (1) gestational age at birth (all control infants had a gestational age at birth that was within 1 week of their matched case infant); (2) PMA at event (all control infants had a 3-minute non-event epoch recorded at a PMA that was within 1 week of the PMA at which the event occurred in the matched case infant); (3) time of day of event (all control infants had a 3-minute non-event epoch recorded at a time of day that was within 1 hour of the time at which the event occurred in the matched case infant); and (4) date and site of enrollment (as the last criteria for matching, when multiple potential control infants met all the aforementioned criteria, then infants were selected with the study identification numbers that were closest in proximity). Because identification numbers were assigned sequentially, by site, this process

served to match, to the extent possible, site and date of enrollment. There were two control epochs from two different control infants chosen to compare with each EE for the cases. Our rationale for not using an infant as its own control was that these young infants rarely changed position during the course of their sleep.

Infant position (supine, prone, side, or indeterminate) was determined by using an accelerometer as the sensor.⁸ The prone and supine positions were measured directly, and the assignment of side position was inferred by comparison with when the infant was also observed sometime during the monitoring in either the supine or prone position. Specifically, an accelerometer placed on the infant’s back showed a force of plus or minus 1 g (ie, the force of gravity on a 1 g mass) when the infant was in the prone or supine position, respectively, but it showed 0 g when the child was side (neutral position) or when the accelerometer was not connected (indeterminate). Thus, registration of a change in force was confirmation that the accelerometer was indeed attached to the infant’s back. When we could not confirm attachment, 0 g was considered “indeterminate.” The period during which the position was determined was either the 75 seconds preceding any event for the cases or during the “non-event” recordings of control subjects.

Statistical Analysis

We examined the association between infant sleep position (supine, prone, side, or indeterminate) and being an infant with at least one EE (ie, being a case) by calculating ORs and their 95% CIs. Side and indeterminate position were combined because it was not always possible to determine whether an infant was in the side position or the position was indeterminate. We used conditional logistic regression to account for the matched design of the study and to adjust for these potential confounders or effect-modifiers: sex, age, race, birth weight, PMA at birth, being sibling of SIDS victim, being born preterm, history of apparent life-threatening event, mother’s age, and mother’s education. The associations were expressed as adjusted ORs with corresponding 95% CIs. In addition, we ran a model that included all first-order interactions with the aforementioned variables to explore whether there was any effect-modification with these variables.

Maternal and infant characteristics of the cases and control subjects were compared through the independent samples *t* test for continuous measures and the χ^2 test for categorical measures. For all tests, the type I error level was set at 0.05. All analyses were performed with SAS software version 9.2 (SAS Institute, Cary, North Carolina).

Results

The Table shows demographic and clinical data comparing cases and the control subjects and demonstrates no substantial differences in the groups. There were no differences in the cases and control subjects used in this

Download English Version:

<https://daneshyari.com/en/article/6224859>

Download Persian Version:

<https://daneshyari.com/article/6224859>

[Daneshyari.com](https://daneshyari.com)