



Research paper

Outcomes associated with anxiety and depression among men who have sex with men in Estonia

R. David Parker^{a,*}, Liilia Lõhmus^b, Anti Valk^b, Cara Mangine^a, Kristi Rüütel^b^a West Virginia University, School of Public Health, Morgantown, WV, USA^b Infectious Diseases and Drug Monitoring Department, National Institute for Health Development, Hiiumägi 42, Tallinn 11619, Estonia

ARTICLE INFO

Article history:

Received 31 March 2015

Received in revised form

3 April 2015

Accepted 7 May 2015

Available online 15 May 2015

Keywords:

Men who have sex with men

Depression

Anxiety

Substance use

Estonia

ABSTRACT

Background: Evidence supports that gay, lesbian, and transgender populations are at increased risk for mental health problems. The current study was conducted to estimate the prevalence of anxiety and depressive symptoms and related factors among men who have sex with men (MSM) in Estonia. This is the only known study of its type in the Eastern European region.

Methods: In 2013, an Internet-based survey was conducted among men who have sex with men, collecting data on socio-demographics, sexuality, drug and alcohol use, mental health, suicidality, and internalized homonegativity.

Results: 265 MSM completed the questionnaire (median age 31 years, 90% Estonian, 71% from the capital; 73% employed full-time, 42% with at least college education; 44% in a steady relationship; 72% gay, 23% bisexual; 23% reported illegal drug use in the last 12 months). One third met criteria for problem drinking and depressive symptoms with one quarter for anxiety symptoms. Suicidality was 44% reported lifetime suicidal thoughts and 11% with at least one suicide attempt.

Limitations: The study instruments, EST-Q and CAGE, only measure symptoms and are not diagnostic tests. A non-random sample which could possibly include persons with easier internet access. Self-reported data and cross-sectional study design are prone to issues with recall bias and temporality.

Conclusions: With demonstrated high rates of anxiety, depression, drinking, and drug use among MSM which is consistent with similar studies in Western countries, further research could help determine effective MSM focused interventions to address the broad spectrum of issues among MSM.

© 2015 Elsevier B.V. All rights reserved.

1. Introduction

Growing evidence supports the notion that gay, lesbian, and transgender populations are at increased risk for a wide range of mental health problems, including links with bipolar disorder, depression, suicidality, and drug and alcohol misuse (Cochran et al., 2013; Cochran and Mays, 2000; Farmer et al., 2013; Guerim et al., 2015). The preponderance of research was conducted in the United States and other Western cultures, while research is extremely limited on MSM from Eastern European nations, specifically independent nations in Central and Eastern European region. These nations have unique cultural and social characteristics dominating society for many years, including the importance of integration and devaluing of individualism during Soviet Union times (Kolves et al., 2006). Understanding the burden of disease among groups at increased risk is very important for any

national response to public health and for men who have sex with men (MSM) in this region, there are unique challenges, such as lack of social support, integration, tolerance, and acceptance (Donoghoe et al., 2005).

Isolation and lack of identifiable peers as well as negative behaviors and images from the larger social groups can increase negative self-feelings regarding same sex attraction (Cochran and Mays, 2000; King et al., 2008; Mosack et al., 2013; Raymond et al., 2011; Ross et al., 2013). The result is internalized homophobia or 'homonegativity'. This internalized homonegativity (IH) is a result of the treatment or perceived treatment of gay persons by society. IH has been shown to lead to increased depression and in some instances, high risk behavior in non-Western countries. The link to depression, discomfort, and anxiety as well as high risk behavior could have a cumulative effect if persons with depression are more likely to engage in high risk behaviors and experience high levels internalized depression and IH. Similar findings have included a disproportionate representation of MSM among persons with depressive and bipolar disorder, specifically persons who identify as bisexual (Akiskal, 2005; Guerim et al., 2015).

* Correspondence to: West Virginia University School of Public Health, 1 Medical Center Drive, Morgantown, West Virginia 26506. Tel.: +1 304 293 7486; fax: +1 304 293 2700.

E-mail address: rdparker@hsc.wvu.edu (R.D. Parker).

Recreational drug use, lower educational attainment, income, and more negative health outcomes have also been linked. The burden of mental health issues and substance use have been estimated in many European countries, using systematic reviews from national studies

Findings indicate that more than one quarter (27%) of adults aged 18–65 exhibited a symptomatology consistent with a mental health disorder in the previous one year. Findings using data from the Global Burden of Disease Study identify Eastern Europe as the global region with the highest proportion of depression, substance misuse, and suicidality at close to three times that of the global burden (Ferrari et al., 2014; Whiteford et al., 2013). In this project, we sought to understand factors associated with anxiety and depression pertinent to MSM in Estonia, a Northern European nation with strong Eastern European culture.

2. Methods

We conducted an Internet based study among MSM in 2013. Eligibility criteria included: self-identified as male; living in Estonia; age 18 years and older; sexually attracted to men and/or have had sex with a man. The study was approved by the Tallinn Medical Research Ethics Committee. Before answering the questions, all participants were required to consent to participate.

An assessment with 144 primarily closed-ended questions was created using Lime Survey (Schmitz, 2014). The questionnaire's design was based on the investigator's previous experiences (Tripathi et al., 2009; Weatherburn et al., 2013) and included: demographic data, such as income, education, number of people in home, sexual orientation, openness about being gay, partners, anxiety and depression history, and antidepressant/tranquilizer use.

Symptoms of depression and anxiety were assessed using Emotional State Questionnaire (EST-Q), a self-rating scale containing six subscales and reflecting symptoms of depressive and anxiety disorders according to ICD-10 and DSM-IV (Aluoja et al., 2004, 1999; Oopik et al., 2006). Consisting of 28 items, the depression subscale contains eight items and the anxiety subscale is six items. Each item is rated on a five-point scale, from 0 to 4 (0=not at all; 4=all the time). The participants reported the extent to which specific problems troubled them during the past 4 weeks.

Alcohol use was assessed using the CAGE questionnaire (Ewing, 1984). The CAGE questionnaire is widely used to screen patients from the general population for alcohol abuse or dependence in a clinical setting. Assessment uses four yes/no domain based questions. Subjects responding affirmatively to two or more questions were classified as CAGE positive, with a high likelihood of the presence of alcoholism (lifetime prevalence). Suicidality was assessed over the last 12 months and lifetime. Specifically, we focused on reporting of suicidal ideation and suicide attempts. Each was assessed through two questions, one for 12 months and one for lifetime. The questions used were the same as in a general population study, therefore, we should be able to make cross comparisons (Tekkel and Veideman, 2013).

3. Recruitment and promotion

The study was promoted through Estonia-based gay online social media, gay community organizations, national network of anonymous HIV testing sites and youth counseling centers. Advertisements directed users to a study splash page which described study aims. Additionally, potential respondents were informed their data would be anonymized, including neither IP-address tracking nor utilization of cookies on their local machine. Following submission, all respondents were automatically directed to a landing page which provided further

information about HIV and STI testing and prevention possibilities. All participants who completed the questionnaire were offered voluntary, anonymous and free of charge HIV and STI testing. All study materials (advertisements, opening page, questionnaire and landing page) were presented in Estonian and Russian allowing a participant to choose his preferred language.

4. Analyses

Statistical analyses were performed with STATA 11.0 (StataCorp LP, College Station, TX). Descriptive statistics were used to characterize participants. Associations between participant characteristics and depression/anxiety scores were evaluated by using the Wilcoxon rank-sum test or Fisher exact test, followed by univariate and multivariable logistic regressions. Multivariable logistic regression was used for multi-variable modeling. Bayesian Information Criteria (BIC) was used, along with 95% confidence intervals and odds ratio effect sizes to determine variables to be included in the model. The adjusted r square, while not a definitive indicator of variance assumed by the data, was used as a measure of fit and one of the factors in model selection. Models were initiated using an investigator derived model based on factors perceived to be most influential according to the current literature and identified as statistically significant through bivariate analyses. Each initial model, one for symptoms of anxiety and one for symptoms of depression, included: the ability to cope with the current financial situation, number of people with whom the participant lived, proportion of friends who knew of same sex attraction, current steady relationship status, length of time in current steady relationship, number of steady male sex partners with whom participant had anal sex in the last 12 months, unprotected anal intercourse with non-steady partner with an unknown HIV status, participant was paid for sex, participant's reported perceived HIV status, if participant reported ever being diagnosed with syphilis or hepatitis B, and the participant's consideration of his overall health status.

5. Results

Overall, 430 people began the questionnaire with a 70.2 % ($n=301$) completion; 29.8 % ($n=129$) aborted the survey prior to completion. Of persons completing the survey, 12.3% ($n=36$) did not meet the inclusion criteria (18 were not Estonian residents, 10 were not MSM, and 8 did not meet the minimum age). This yielded a total sample size of 265 for analyses. Table 1 presents socio-demographic and health related data of the participants based on depressive symptoms cut-off level.

Almost one third (32.1%, $n=85$) of the participants demonstrated EST-Q depression scores above the cut-off level, indicating possible depression. The mean depression score of the total sample was 9.4 (median 8; SD 6.7; range 0–29). Close to one quarter (23.4%, $n=62$) had EST-Q anxiety scores above the cut-off level, indicating symptoms associated with anxiety. The mean anxiety score was 7.9 (median 7; SD 5.1; range 0–24). Using a pairwise correlation test, given the dependence of each score, with a Bonferroni adjustment for the alpha, depression and anxiety scores were highly correlated ($r=0.52$). Of the total sample, 17.7% ($n=47$) screened positive for both anxiety and depression. Almost 9% ($n=22$) of respondents with EST-Q depression scores above the cut-off reported being told by a doctor in the previous 12 months they had depression. 6.3% ($n=16$) of respondents with EST-Q anxiety scores above the cut-off level were told by a clinician in the previous year that he had an anxiety disorder.

Nearly one third (32.8%, $n=87$) of our sample had a positive CAGE score, that is a score above the cut-off level, mean score of

Download English Version:

<https://daneshyari.com/en/article/6231469>

Download Persian Version:

<https://daneshyari.com/article/6231469>

[Daneshyari.com](https://daneshyari.com)