



# Logistic regression model for identification of right ventricular dysfunction in patients with acute pulmonary embolism by means of computed tomography



Grzegorz Staskiewicz<sup>a,b,\*</sup>, Elżbieta Czekajska-Chehab<sup>a</sup>, Sebastian Uhlig<sup>a</sup>, Jerzy Przegalinski<sup>c</sup>, Ryszard Maciejewski<sup>b</sup>, Andrzej Drop<sup>a</sup>

<sup>a</sup> 1st Department of Radiology, Medical University of Lublin, Lublin, Poland

<sup>b</sup> Department of Human Anatomy, Medical University of Lublin, Lublin, Poland

<sup>c</sup> Department of Cardiology, Medical University of Lublin, Lublin, Poland

## ARTICLE INFO

### Article history:

Received 29 August 2011

Received in revised form 3 January 2013

Accepted 4 February 2013

### Keywords:

Pulmonary embolism

Computed tomography

Right ventricular dysfunction

## ABSTRACT

**Purpose:** Diagnosis of right ventricular dysfunction in patients with acute pulmonary embolism (PE) is known to be associated with increased risk of mortality. The aim of the study was to calculate a logistic regression model for reliable identification of right ventricular dysfunction (RVD) in patients diagnosed with computed tomography pulmonary angiography.

**Material and methods:** Ninety-seven consecutive patients with acute pulmonary embolism were divided into groups with and without RVD basing upon echocardiographic measurement of pulmonary artery systolic pressure (PASP). PE severity was graded with the pulmonary obstruction score. CT measurements of heart chambers and mediastinal vessels were performed; position of interventricular septum and presence of contrast reflux into the inferior vena cava were also recorded. The logistic regression model was prepared by means of stepwise logistic regression.

**Results:** Among the used parameters, the final model consisted of pulmonary obstruction score, short axis diameter of right ventricle and diameter of inferior vena cava. The calculated model is characterized by 79% sensitivity and 81% specificity, and its performance was significantly better than single CT-based measurements.

**Conclusion:** Logistic regression model identifies RVD significantly better, than single CT-based measurements.

© 2013 Elsevier Ireland Ltd. All rights reserved.

## 1. Introduction

Pulmonary embolism (PE) is the third most common of the “great cardiovascular killers” after myocardial infarction and stroke [1]. As opposed to the two others, the clinical symptoms and signs in patients with PE may mimic multiple other conditions, cardiovascular or pulmonary. Imaging methods have become one of the cornerstones in the diagnostic process for this group of patients.

Wide accessibility of multidetector computed tomography (MDCT), full chest submillimeter coverage with one breathhold,

concurrent assessment of pulmonary parenchyma and mediastinum, and, last but not least, chance of providing alternative diagnosis, have lead to MDCT being the gold standard in the imaging of suspected PE.

Visualization of the heart chambers and the mediastinal vessels allowed signs of right ventricular dysfunction (RVD) to be assessed during the same computed tomography pulmonary angiography (CTPA) performed to confirm presence of PE.

Even in patients with hemodynamically stable PE, RVD is associated with an increased risk of mortality. However, in normotensive patient, identification of RVD requires additional testing, including echocardiography or biomarkers [2].

Multiple CT signs of RVD were evaluated and reported in recent years, with different cutoff values, sensitivities and specificities reported. Therefore, the aim of the study was to generate a highly accurate predictive model composed of multiple measurements for the identification of RVD in the course of acute PE by computed tomography pulmonary angiography (CTPA).

\* Corresponding author at: 1<sup>st</sup> Department of Radiology, Medical University of Lublin, ul. Jaczewskiego 8, 20-090 Lublin, Poland. Tel.: +48 609 522 559.

E-mail addresses: [grzegorz.staskiewicz@gmail.com](mailto:grzegorz.staskiewicz@gmail.com) (G. Staskiewicz), [czekajska@gazeta.pl](mailto:czekajska@gazeta.pl) (E. Czekajska-Chehab), [uhlig@eranet.pl](mailto:uhlig@eranet.pl) (S. Uhlig), [jerzy.przegalinski@umlub.pl](mailto:jerzy.przegalinski@umlub.pl) (J. Przegalinski), [maciejewski.r@gmail.com](mailto:maciejewski.r@gmail.com) (R. Maciejewski), [andrzej.drop@umlub.pl](mailto:andrzej.drop@umlub.pl) (A. Drop).

## 2. Material and methods

The institutional ethical review board approved the protocol of the study with a waiver of informed consent (Decision No.: KE-0254/41/2011). For the purpose of this retrospective study, we identified 97 patients, in whom echocardiographic examination with pulmonary artery systolic pressure (PASP) assessment was performed within the 12 h of the CT scan. The CTPA studies were performed with 64-row VCT scanner (GE Medical Systems), with a standard protocol used at our institution, which includes cranio-caudal scanning of the chest,  $64 \times 0.625$  mm collimation, slice thickness 0.625 mm, continuous reconstruction. SmartPrep protocol was used for adjustment of the injection time of average 70 mL (1 mL/kg) of iodinated contrast medium (Ultravist 370, Bayer Healthcare, Germany), which was administered at a flow rate of 4–5 mL/sec by automatic injector, followed by 40 mL NaCl bolus.

Raw tomographic data was imported into the OsiriX software. The exams were reviewed by two readers in consensus. The readers were blinded to PASP values.

Obstruction of pulmonary arteries was scored according to Qanadli et al. [3]. Maximum short axis of the right (RV) and left ventricle (LV) were measured at a four chamber view, according to Quiroz et al. [4]. The diameter of the pulmonary artery (PA) was measured proximal to its bifurcation; diameter of superior vena cava (SVC) was measured at the level of arch of azygos vein; and diameter of inferior vena cava (IVC) in its thoracic part. Dimensions of each mediastinal vessel were measured in a plane perpendicular to a long axis of the vessel, with respect to the largest observed diameter. RV/LV ratio was calculated. Diameter of coronary sinus (CS) was measured in the axial plane, proximal to its opening, as described elsewhere [5]. Position of the interventricular septum was evaluated and classified as normal or convex toward left ventricle. Contrast medium reflux into the IVC was also recorded.

RVD was defined as echocardiographically measured PASP value  $>30$  mmHg, as described elsewhere [6]; based on this value, patients were divided into RVD(–) and RVD(+) groups (PASP  $\leq 30$  mmHg and  $>30$  mmHg, respectively).

Statistical analysis was performed with SPSS 16.0, SPSS Inc., Chicago, IL. Quantitative parameters are presented as minimal, maximal and median values. Qualitative parameters are presented as numbers and percentages.

$\chi^2$  test was used to compare qualitative data, and the Mann–Whitney test for quantitative data. Values of  $p \leq 0.05$  were considered to be significant. The area under ROC curve (AUC) and 95% confidence limits were used to assess predictive power for RVD of all parameters that differed significantly between the study groups. Cutoff values with best sensitivity to specificity ratio were calculated.

In order to find the best combination of CT-based indicators of RVD, a multivariate model was built by means of logistic regression analysis. A backward conditional stepwise method was used to select the parameters which were included in the final model. In this process, the weakest predictors of RVD are deleted starting with the least significant ones. The parameters which remain in the final model are independently associated with the risk of RVD. Values predicted by the model belong to (0, 1) interval, and represent probability of RVD. The model is as follows [7]:

$$p = \frac{1}{1 + e^{-z}},$$

where  $e$  is a base of natural logarithm,  $=2.71828\dots$ , and  $z$  is a linear combination of  $x_i$  variables and their estimators  $b_i$  included in the model:

$$z = b_0 + b_1x_1 + b_2x_2 + \dots + b_nx_n.$$

**Table 1**

Characteristics of the study population in patients with and without right ventricular dysfunction: RVD(+) and RVD(–), respectively. Numbers represent median, minimal and maximal values, and (\*) absolute value and percentage for qualitative values.

	RVD(–) PASP $\leq 30$ mmHg; $n = 46$	RVD(+) PASP $> 30$ mmHg; $n = 51$	<i>p</i>
Age [years]	71.5 (27–91)	67 (22–92)	NS
BMI	26.79 (23.23–38.51)	26.57 (19.15–34.29)	NS
Systolic BP [mmHg]	119 (85–168)	124 (70–163)	NS
Heart rate [bpm]	94 (60–125)	96 (48–144)	NS
PASP [mmHg]	25 (18–30)	45 (32–95)	$<0.001$
Length of hospitalization [days]	9 (1–31)	12 (1–44)	0.03
ICU admission (*)	21 (46%)	34 (67%)	0.04
Mortality (*)	4 (8.7%)	7 (13.7%)	NS

The model-based probability of  $>0.5$  classified the patient as RVD(+). The Hosmer–Lemeshow goodness-of-fit test was used to evaluate the overall fit of the final model. To compare the accuracy of the proposed model with single CT-based parameters and calculate specificity, sensitivity and accuracy of the used parameters, ROC curves were constructed, and areas under the curve (AUC) were compared with Aabel 3 package (Gigawiz Ltd., Tulsa, OK).

## 3. Results

The study group consisted of 97 patients, median age 69 years (range: 22–92). 46 patients were included in group RVD(–) and 51 patients in RVD(+) group. Patients characteristics are presented in Table 1. No significant differences of frequency of DVT, immobilization, surgery, or history of malignancy were observed between the groups. RVD(+) group was characterized by a significantly longer hospitalizations, as well as more frequent ICU admissions.

Results of comparative analysis of the CT-based parameters and the probabilities calculated by the logistic regression model are presented in Table 2.

Details of logistic regression analysis are presented in Table 3, and the obtained model for identification of RVD is as follows:

$$z = -8.361 + 0.132 \times \text{Obstruction score} + 0.078 \times \text{RV} + 0.094 \times \text{IVC}.$$

The highest sensitivity and specificity at assumed risk of RVD of 50% for the obtained predictive model were 0.79 and 0.81, respectively. Analysis of the areas under the ROC curves (AUC) showed, that the largest AUC, sensitivity and specificity were observed for the model-calculated probability, as the area was significantly larger ( $p=0.02$ ) than for any of the single CT-based parameters (AUC = 0.860, 95% CI: 0.802–0.917).

Measurements used for the final model are presented in Fig. 1, and the ROC curve of RVD probability calculated by the model is shown in Fig. 2. Hosmer–Lemeshow test showed value of 3.065 ( $p=0.930$ ), which shows a good fit of the model.

## 4. Discussion

Short-term prognosis of PE depends on the hemodynamic status of the patient as well as comorbidities [8]. Therefore, identification of RVD in PE patients is crucial for the prognosis and treatment [2]. However, the number of suggested CT-based signs of RVD is large, and different cutoffs, sensitivities and specificities are reported.

Pulmonary obstruction index according to Qanadli et al. has been linked to RVD since its introduction [3], and its feasibility has been confirmed in more recent papers by Collomb et al. [9] and van der Meer et al. [10]. Furthermore, the Qanadli index has been linked to mortality in PE patients, including a significantly higher 2-weeks mortality rate in those with an obstruction score  $>60\%$

Download English Version:

<https://daneshyari.com/en/article/6244595>

Download Persian Version:

<https://daneshyari.com/article/6244595>

[Daneshyari.com](https://daneshyari.com)