

## Characteristics of Testosterone Deficiency Syndrome in Men With Chronic Kidney Disease and Male Renal Transplant Recipients: A Cross-Sectional Study

M.G. Park, H.S. Koo, and B. Lee

---

### ABSTRACT

**Objectives.** Testosterone deficiency syndrome (TDS) is common among male patients with chronic kidney disease (CKD). We compared the characteristics of TDS in men with CKD versus renal transplantation (RT) with those of age-matched normal controls.

**Materials and Methods.** The 129 patients were: RT recipients ( $n = 25$ ) group I, CKD patients ( $n = 37$ ) group II, and controls ( $n = 67$ ). We performed estimates of testosterone, hemoglobin (Hgb), hematocrit (Hct), glucose, creatinine, and lipid profile. Self-assessment questionnaires—International Index of Erectile Function (IIEF), Aging Males' Symptoms (AMS), Center for Epidemiologic Studies Depression Scale—were used to evaluate erectile function, testosterone deficiency, and depression, respectively. We also investigated morning erection as well as the presence and duration of erectile dysfunction (ED).

**Results.** Group I (RT) showed significantly higher serum testosterone levels than group II (CKD), who displayed significantly worse erectile function, more severe testosterone deficiency symptoms, and a greater trend toward depression. Similarly, the prevalences of ED and TDS were significantly greater in group II than group I. Group I and controls differed significantly only in the results of serologic tests, such as serum creatinine, Hgb, and glucose and lipid profiles, but not in serum testosterone levels, scores of self-assessment questionnaires, or prevalence of ED or TDS. Serum testosterone levels correlated significantly with scores on the IIEF and AMS questionnaires in both group II and controls, but not group I.

**Conclusions.** RT recipients showed higher serum testosterone levels and a lower prevalence of TDS with milder symptom severity than CKD patients. RT recipients beyond the early acute posttransplant period, displayed serum testosterone levels and TDS prevalence similar to those of healthy controls. Unlike CKD patients and normal controls, serum testosterone did not significantly influence TDS symptoms in RT recipients.

---

**E**RECTILE DYSFUNCTION (ED) is commonly observed in patients with chronic kidney disease (CKD) and end-stage renal disease (ESRD).<sup>1</sup> The prevalence of ED among CKD men is estimated to be about 50%.<sup>2</sup> Multiple hormonal abnormalities have been discovered to be among the underlying causes of ED in CKD.<sup>3–5</sup> Testosterone deficiency (TD), is the most common hormonal alteration, is due to reduced prolactin clearance<sup>6</sup> and uremic inhibition of the luteinizing hormone effect on Leydig cells.<sup>7</sup> According to recommendations, the clinical and biochemical alterations associated with this condition

in CKD men can be diagnosed as TD syndrome (TDS). It markedly lowers the quality of life and decreases the functions of multiple organ systems.<sup>8</sup> Hence, one main

---

From the Departments of Urology, Nephrology, and Surgery, Seoul Paik Hospital, Inje University, Republic of Korea.

Address reprint requests to, Byungmo Lee, MD, PhD, Assistant Professor, College of Medicine, Inje University Department of Surgery, Seoul Paik Hospital, 9, Mareunnae-Ro, Jung-Gu, Seoul, 100-032, Korea. E-mail: [b0molee@gmail.com](mailto:b0molee@gmail.com)

concern is the impact of renal transplantation on TDS in CKD patients. Several studies have imported that most endocrine abnormalities are reversed after successful renal transplant (RT) with improved erectile function. However, the pattern of changes in testosterone levels varies according to the time after transplantation.<sup>9,10</sup> Furthermore, the majority of studies have not considered the changes in the prevalence and severity of ED or TDS after RT or the correlation between testosterone levels and the severity of those symptoms. Additionally, there is little information about the extent of recovery of clinical and biochemical values after RT relative to the normal range. Therefore, we investigated the characteristics of TDS among male CKD subjects and RT recipients compared with age-matched, normal controls.

## MATERIAL AND METHODS

From December 2011 to June 2012, we consecutively included 129 male RT recipients ( $n = 25$ ; group I) CKD, stage II to V, ( $n = 37$  group II), and age-matched normal controls from our health promotion center ( $n = 67$ ; control group). After obtaining consent for the study, each patient was interviewed by the same physician with regard to their past medical history, including comorbidities, onset and cause of CKD, duration of maintenance hemodialysis, time of RT, erectile function, including its presence and duration of ED as well as morning erection. Group I was questioned about ED before RT, improvement after RT and present ED. Three self-assessment questionnaires—the International Index of Erectile Function (IIEF), the Aging Males' Symptoms (AMS), and the Center for Epidemiologic Studies Depression Scale (CES-D)—were administered to evaluate the severity of ED, TD symptoms, and depression, respectively. After completing the personal interview and the self-assessment questionnaires, we performed measurements of plasma testosterone, hemoglobin (Hgb), hematocrit (Hct), creatinine, and glucose and lipid profile between 9 and 11 AM. Testosterone deficiency was defined as a plasma testosterone level  $<350$  ng/dL. Statistical analysis employed the independent samples students *t*-test, chi-square test, simple correlation test, and multiple regression analysis. A *P* value  $<.05$  was considered significant.

## RESULTS

There was no difference in age among the three groups; group I, 53.5; group II, 55.0; and controls, 52.16. The present comorbidities and causes of CKD did not differ significantly between groups I and II. The mean time from the first diagnosis of CKD was significantly by different; group I, 190 months (range, 12–480) versus group II, 108 months (range, 5–280;  $P = .023$ ). In addition, the duration of maintenance hemodialysis in group I (mean, 22.0 months; range, 3–78) was significantly shorter than that of group II (mean, 53.4 months; range, 3–156;  $P = .008$ ). The mean time since RT in group I, was 124 months (range, 11–288). Among the 25 RT recipients, 21 (84%) experienced ED before RT and 19 of the 21 (90.5%) reported improved function after RT.

Among the 37 group II patients, 18 had ESRD; those with and without ESRD differed significantly in the severity of ED and TDS, according to their total scores of IIEF and

AMS ( $P = .004$  and  $P = .000$ , respectively). Similarly, the CES-D score of men with ESRD was higher than that of patients with stage II to IV CKD ( $P = .103$ ). With respect to plasma testosterone levels, subjects with ESRD displayed lower levels of testosterone than those without ESRD: mean testosterone, 264.2 versus 371.6 ng/dL, ( $P = .070$ ). Among group II patients, the prevalence of ED and TDS differed significantly between patients with ESRD versus those with less severe CKD ( $P = .021$  and  $P = .004$ , respectively). Likewise, morning erection was reported less frequently (5.6%) by ESRD patients than by those with less severe CKD (21.1%;  $P = .168$ ; Table 1).

Comparison between group I and group II revealed the former cohort to display significantly higher serum testosterone than group II: mean, 515.7 versus 323.5 mg/dL ( $P = .001$ ). With respect to the scores of the IIEF, AMS, and CES-D, group II showed significantly worse erectile function, more severe TD symptoms, and greater tendencies toward depression than group I. Similarly, the prevalences of ED and TDS were significantly higher in group II than group I ( $P = .008$  and  $P = .019$ ); (Table 2).

Comparison of group I with the 18 group II patients with stages II to IV CKD revealed a similar pattern of significant differences. RT recipients displayed higher testosterone levels, more tolerable symptom scores on (IIEF, AMS, and CES-D), and lower prevalences of ED and TDS than patients with stages II to IV CKD.

RT recipients (group 1) were compared with normal controls. Although significant differences were noted in the results of serum creatinine, Hgb, Hct, glucose, and lipid profiles, there was no difference in serum testosterone levels (493.9 vs 515.7 ng/dL;  $P = .658$ ). The total scores on self-assessment questionnaires (IIEF, AMS, and CES-D) also did not differ significantly between these two groups ( $P = .354$ , .684, and .256 respectively). The prevalences of ED and TDS among group I were not significantly different from those of the controls ( $P = .382$  and .390, respectively; Table 2).

The correlation between serum testosterone levels and symptom severity was investigated using simple correlation tests. Both group II and controls displayed significant correlations of serum testosterone levels with IIEF and AMS scores ( $P < .05$ ). However there were no correlations in group I between serum testosterone levels and self-assessment questionnaire scores ( $P > .05$ ). Multiple regression analysis to assess the impact of plasma testosterone levels on symptom severity among the multiple variables of plasma Hgb, Hct, creatinine, and glucose and lipid profile revealed a similar pattern of significance between plasma testosterone levels and self-assessment questionnaires scores in both group II and controls ( $P < .05$ ). However, plasma testosterone levels did not show a significant impact on symptom severity in group I ( $P > .05$ ).

## DISCUSSION

The incidence of testosterone deficiency varies from 6% to 9.5% in the community and increases to 15% to 30% among

Download English Version:

<https://daneshyari.com/en/article/6249921>

Download Persian Version:

<https://daneshyari.com/article/6249921>

[Daneshyari.com](https://daneshyari.com)