



Short Communication

Resurgence of sucrose and cocaine seeking in free-feeding rats



Timothy A. Shahan*, Andrew R. Craig, Mary M. Sweeney.

Utah State University, United States

HIGHLIGHTS

- Loss of an alternative reward induced relapse of sucrose and cocaine seeking in free-feeding rats.
- It is unlikely that an acute hunger state is responsible for resurgence.
- These procedures may be useful for examining the neurobiology of resurgence.

ARTICLE INFO

Article history:

Received 19 August 2014
 Received in revised form 21 October 2014
 Accepted 30 October 2014
 Available online 15 November 2014

Keywords:

Resurgence
 Relapse
 Drug self-administration
 Alternative reinforcement
 Extinction
 Operant behavior

ABSTRACT

Resurgence is relapse of an extinguished operant response following the removal of alternative reinforcement. In animal models of resurgence to date, rats have been food deprived and food is used as the source of alternative reinforcement. Thus, when the alternative reinforcer is removed, the only remaining source of food during experimental sessions is no longer available. Acute food deprivation is known to produce reinstatement of drug seeking, thus such deprivation has been suggested a potential mechanism of resurgence. The present experiments examined whether resurgence of sucrose and cocaine seeking could be obtained with rats that were not food deprived. Free-feeding rats were trained to press a lever for either sucrose (Experiment 1) or cocaine infusions (Experiment 2). Next, lever pressing was extinguished and an alternative response (nose poking) was reinforced with sucrose. When nose poking was also placed on extinction, resurgence of both sucrose and cocaine seeking was observed. Thus, resurgence of both sucrose and cocaine seeking can be obtained in rats that are not food restricted and it appears unlikely that an acute hunger state is responsible for resurgence. In addition, the present procedures for studying resurgence in the absence of interpretive complexities introduced by the use of food-deprivation may prove useful for further investigations of the neurobiological mechanisms of resurgence.

© 2014 Elsevier B.V. All rights reserved.

An increase in an extinguished operant behavior generated by loss of an alternative source of reinforcement is known as resurgence [1–4]. Alternative reinforcement is a common component of behavioral treatments for substance abuse [5], and suspension of such alternative reinforcement when treatment ends is often associated with relapse [6–8]. Furthermore, loss of non-drug sources of reinforcement (e.g., social, vocational) is often cited as a source of relapse in human substance abusers [9–12]. Resurgence of drug-maintained behavior of rats has been demonstrated with alcohol [2] and cocaine [13] self-administration when an alternative source of food reinforcement is removed. As a result, the resurgence paradigm has been suggested as an animal model of relapse to

drug seeking induced by loss of alternative non-drug reinforcement [2,13–15].

To date, all non-human experiments on resurgence of drug [2,13,16] or food [1,17–20] seeking have used food-deprived subjects. In the resurgence test when the alternative reinforcement is removed, food is no longer available during the experimental session. Both acute and chronic food deprivation stress are known to generate relapse of drug-seeking [21–23], and it has been suggested that an acute hunger state could be the source of the resurgence effect [24]. Thus, the goal of the present experiments was to examine whether resurgence of sucrose (Experiment 1) and cocaine (Experiment 2) seeking could be obtained with rats that were not food deprived and to provide a procedure for examining resurgence that is free of the interpretive complexities introduced by food-deprivation.

In Experiment 1, 10 male Long-Evans rats (Charles River, Portage, MI, USA) were used. Rats were 71–80 days old upon arrival and were fed Harlan Teklad Rodent Diet 8604 ad libitum. Rats were

* Corresponding author at: 2810 Old Main Hill, Logan, UT 84322, United States.
 Tel.: +1 435 - 797 - 1460; fax: +1 435 - 797 - 1448.
 E-mail address: tim.shahan@usu.edu (T.A. Shahan).

allowed 2 weeks to adjust to handling and colony conditions before beginning the study. Sessions took place in four Med Associates operant chambers (St. Albans, VT, USA), described elsewhere [25]. Each chamber was equipped with two levers on either side of a food receptacle equipped with photobeam sensors to detect head entries. A small opening in the receptacle allowed a 0.1-ml dipper cup to be accessed for 4 s when the dipper arm was elevated upon completion of the scheduled response requirement. The 45-ml solution tray contained a 32% sucrose and water mixture that was newly mixed each day to ensure consistency in taste. The chamber wall opposite the levers and food aperture housed five small holes equipped with photobeams to detect nose pokes.

Prior to the start of the experiment rats were trained to consume the sucrose reinforcer for the dipper. Sucrose was delivered on a variable-time (VT) 60-s schedule in 45-min sessions. Throughout the experiment, when the dipper arm was elevated (always 4-s access), the food receptacle was illuminated and any previously lit stimulus was turned off. Only the food receptacle light was used during magazine training. Training lasted until all rats completed a session with head entries on greater than 80% of sucrose deliveries.

Next during training rats earned sucrose presentations contingent upon pressing a lever, the location of which (left or right) was counterbalanced and signaled by LEDs above the lever. During training, the ratio requirement doubled after every five sucrose deliveries from a fixed-ratio (FR) 1 to an FR 8. Next, the rats experienced a variable-interval (VI) 15-s then 30-s schedule. Training sessions began using the last schedule of reinforcement in place in the prior day's session. Session duration was 30 min, minus time for dipper delivery in training and all subsequent phases. Phase 1 began in the session following exposure to the VI 30-s schedule. Training lasted between 1 and 7 sessions.

In **Phase 1 (Target Reinforced)** rats received 50 sessions of training in which sucrose was available for presses to the target lever on a VI 45-s schedule. Presses to the inactive lever and alternative nose poke were counted but had no consequences. In **Phase 2 (Extinction + Alternative Reinforcement)**, presses to the target lever and inactive lever were recorded but had no effect. Nose pokes into the illuminated left-most nose poke aperture produced sucrose deliveries on a VI 10-s schedule. Phase 2 lasted 10 sessions. In **Phase 3 (Resurgence Test)**, sucrose was no longer available for the target or alternative response during this phase, although the solution was present in the dipper tray to keep olfactory cues consistent across phases. Phase 3 lasted four sessions.

In the last session of Phase 1, mean total target lever responses, inactive lever responses, alternative nose pokes, and sucrose deliveries were 204.7 (SD=152.44), 1.98 (SD=2.28), 1.70 (SD=2.06), and 61.90 (SD=20.15), respectively. At the end of Phase 2 (Extinction + Alternative Reinforcement) total target responses had decreased to 6.60 (SD=5.85) presses and the number of alternative responses increased to 62.00 (SD=20.11) pokes, resulting in an average of 61.90 sucrose deliveries (SD=20.15). Total inactive responses remained low at 0.9 (SD=1.37) presses.

Fig. 1 shows cumulative sucrose-lever response across successive 5-min bins in the last session of Phase 2 and the four sessions of Phase 3 (Resurgence Test). For all rats, sucrose-lever responding was greater during the first session of Phase 3 than in the last session of Phase 2. Response output varied across Phase 3 sessions, but generally remained elevated relative to last session of Phase 2. The bottom right panel of Fig. 1 shows mean cumulative sucrose-lever responses and total inactive-lever responses for the last session of Phase 2 and the first session of Phase 3. The conclusion that resurgence was obtained in the first session of Phase 3 is further supported by a 2 (last Phase 2 versus first Phase 3) \times 5 (Bin) repeated-measures ANOVA documenting significant main effects of Phase [$F(1, 5)=6.72, p<.05$] and Bin [$F(8, 40)=24.94, p<.001$] and a Phase \times Bin interaction [$F(8, 40)=3.96, p<.01$]. Resurgence

was specific to sucrose-lever responding as there was no significant increase in inactive lever pressing between the last session of Phase 2 and the first sessions of Phase 3 [$t(9)=1.53, ns$].

In Experiment 2 six experimentally naïve male Long Evans rats (Charles River, Portage, MI) served. Rats were housed individually in a temperature- and humidity-controlled colony room with a 12:12 h light/dark cycle (lights on at 7:00 AM) and had unrestricted access to food and water. Each rat was approximately 90 days of age at the time of surgery (see below). Animal care and all procedures detailed below were conducted in accordance with guidelines set forth by Utah State University's Institutional Animal Care and Use Committee. Four modular operant self-administration chambers like those used in Experiment 1 and also equipped for IV drug self-administration were used. Tygon tubing encased in a metal-spring tether was inserted through a 4 cm diameter hold cut in the center of the roof of each chamber. At all times, the Tygon tubing and tether were attached to rats' back-mounted cannulae. The other end of the tubing attached to a stainless-steel swivel (Instech, Plymouth Meeting, PA) suspended ~15 cm above the roof of the chamber. Another 60 cm of Tygon tubing attached the swivel to a 60-ml syringe mounted on an infusion pump (Med Associates) positioned outside the sound-attenuating cabinet.

Prior to the experiment, rats underwent jugular-catheterization surgery which was preceded by injections of antibiotic (gentamicin, 2.0 mg/kg, IM), analgesic/anti-inflammatory (flunixin meglumine, 1.1 mg/kg, IP), and anesthetic (ketamine, 75 mg/kg, and xylazine hydrochloride, 7.5 mg/kg, IM) drugs. An indwelling, back-mounted cannula (Plastics One, Roanoke, VA) attached to a silastic catheter (SAI-Infusions, Lake Villa, IL) was inserted through an incision in the rat's lower back and fed subcutaneously to a 0.5 cm incision centered 2 cm below the rat's shoulder blades. The catheter was fed subcutaneously from the upper-back incision to an incision made in the right ventral surface of the neck, then inserted into the right jugular vein. Following surgery, rats were given injections of flunixin meglumine (1.1 mg/kg, IP), gentamicin (2.0 mg/kg, IM), and electrolyte solution (Ringer's, 5 cc, SC) every 12 h for 2–5 days. Rats began the procedure detailed below a minimum of 5 days post-surgery. Cocaine hydrochloride was dissolved in a sterile 0.9% saline solution to a concentration 2.56 mg/ml. Drug doses were adjusted by changing the activation duration of the fixed-speed (0.0527 ml/s) syringe pump.

Prior to beginning the experiment rats experienced several sessions of training to ensure they reliably drank a 32% sucrose solution from the 0.1 ml dipper when it operated. Dipper training proceeded as described for Experiment 1. Next, rats were trained to self-administer IV cocaine infusions. Sessions during self-administration training and all phases of the experiment detailed below were 45 min long and began with illumination of the house light and target-lever LEDs. During the first session of training, a single press (i.e., fixed-ratio [FR] 1) to the target lever produced a 1 mg/kg infusion of cocaine. Here and below, each cocaine infusion was accompanied by blackout of the chamber and activation of the Sonalert for the duration of the infusion. After a minimum of three sessions of self-administration under FR 1, the number of responses required per infusion was increased to 2, and then to 3 after a minimum of three additional sessions. The dose of cocaine then was decreased across sessions to 0.75 mg/kg, 0.5 mg/kg, and, finally, 0.32 mg/kg per infusion. This phase ended when the rats reached the 0.32 mg/kg dose.

In **Phase 1 (Target Reinforced)** of the experiment, rats self-administered 0.32 mg/kg infusions of cocaine according to an FR 3 schedule for presses to the target lever. Phase 1 continued for a minimum of 14 sessions and until responding over the last three sessions occurred at stable rates (i.e., absence of reliable trend). In **Phase 2 (Extinction + Alternative Reinforcement)**, presses to the target lever and inactive lever were recorded but had no effect.

Download English Version:

<https://daneshyari.com/en/article/6257366>

Download Persian Version:

<https://daneshyari.com/article/6257366>

[Daneshyari.com](https://daneshyari.com)