

On the difficulty to communicate with fMRI-based protocols used to identify covert awareness

Alexandre Comte, Damien Gabriel, Lionel Pazart, Eloi Magnin, Elodie Cretin, Emmanuel Haffen, Thierry Moulin, Régis Aubry

PII: S0306-4522(15)00501-1
DOI: http://dx.doi.org/10.1016/j.neuroscience.2015.05.059
Reference: NSC 16309

To appear in: *Neuroscience*

Accepted Date: 24 May 2015

Please cite this article as: A. Comte, D. Gabriel, L. Pazart, E. Magnin, E. Cretin, E. Haffen, T. Moulin, R. Aubry, On the difficulty to communicate with fMRI-based protocols used to identify covert awareness, *Neuroscience* (2015), doi: http://dx.doi.org/10.1016/j.neuroscience.2015.05.059

On the difficulty to communicate with fMRI-based protocols used to identify covert awareness

Alexandre Comte[1,2,3,*], Damien Gabriel[2,3,*], Lionel Pazart[2], Eloi Magnin[1,2,3,4], Elodie Cretin[2], Emmanuel Haffen[2,3], Thierry Moulin[1,2,3,4], Régis Aubry[2,5]

[1]Department of Functional Neuroimaging Research (IFR 133 INSERM), CHU Besançon, France
[2]Center for Clinical Investigation in Technological Innovation (CIT808), CHU Besançon, France
[3]Besançon Laboratory of Integrative and Clinical Neuroscience (EA-481), Franche-Comté University, 25030, Besançon, Cedex, France
[4]Department of Neurology, CHU Besançon, France
[5]Department of Palliative care, CHU Besançon, France
[*]These authors contributed equally to this work

Corresponding author:

Alexandre Comte
Imaging Pole
Department of Functional Neuroimaging Research
J.Minjoz University Hospital
25030 Besançon Cedex
France
Tel: 0033 (0)381668948
Fax: 0033 (0)381669315
alexandre.comte@univ-fcomte.fr

Keywords: unresponsive wakefulness syndrome; awareness; communication; fMRI; correlation; support vector machine