



Research paper

Entrainment of chaotic activities in brain and heart during MBSR mindfulness training



Junling Gao^{a,b,c}, Jicong Fan^a, Bonnie Wai Yan Wu^a, Zhiguo Zhang^b, Chunqi Chang^d, Yeung-Sam Hung^b, Peter Chin Wan Fung^{a,e}, Hin hung Sik^{a,*}

^a Centre of Buddhist Studies, The University of Hong Kong, Hong Kong

^b Department of Electrical and Electronic Engineering, The University of Hong Kong, Hong Kong

^c Alzheimer's Disease Research Network, The University of Hong Kong, Hong Kong

^d School of Biomedical Engineering, Shenzhen University, PR China

^e Centre on Behavioral Health, The University of Hong Kong, Hong Kong

HIGHLIGHTS

- Wavelet entropy is used to measure the chaotic activities.
- Coordination between electronic activities of brain and heart.
- MBSR meditation can reduce the chaotic activity of brain waves.

ARTICLE INFO

Article history:

Received 9 August 2015

Received in revised form

12 November 2015

Accepted 4 January 2016

Available online 16 January 2016

Keywords:

MBSR

EEG

Wavelet entropy

Source analysis

Brain–heart coordination

ABSTRACT

The activities of the brain and the heart are dynamic, chaotic, and possibly intrinsically coordinated. This study aims to investigate the effect of Mindfulness-Based Stress Reduction (MBSR) program on the chaoticity of electronic activities of the brain and the heart, and to explore their potential correlation. Electroencephalogram (EEG) and electrocardiogram (ECG) were recorded at the beginning of an 8-week standard MBSR training course and after the course. EEG spectrum analysis was carried out, wavelet entropies (WE) of EEG (together with reconstructed cortical sources) and heart rate were calculated, and their correlation was investigated. We found enhancement of EEG power of alpha and beta waves and lowering of delta waves power during MBSR training state as compared to normal resting state. Wavelet entropy analysis indicated that MBSR mindfulness meditation could reduce the chaotic activities of both EEG and heart rate as a change of state. However, longitudinal change of trait may need more long-term training. For the first time, our data demonstrated that the chaotic activities of the brain and the heart became more coordinated during MBSR training, suggesting that mindfulness training may increase the entrainment between mind and body. The 3D brain regions involved in the change in mental states were identified.

© 2016 The Authors. Published by Elsevier Ireland Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Living organisms are nonlinear, complex systems, with certain degree of chaoticity in their physiological variables. For example, the electroencephalography (EEG) waves are known for their dynamic activity with brain networks operating between order and chaos [1]. Likewise, heart rate variability (HRV) is non-zero for a living subject. However, chaotic degree must be maintained at a

certain yet not well-known limiting level. The Lyapunov exponent (LE) of EEG can be considered as a measure of chaotic behavior. Previously, we strived to apply the LE of EEG as a parameter to predict epilepsy [2]. Similarly, we also analyzed the chaotic features of electrocardiogram (ECG) waves to look for a parameter characterizing ECG signals that would likely lead to pre-matured ventricular contractions [3]. Since the advent of high-resolution EEG machine (with ~100 or more channels), a number of methods have been developed to measure objectively the chaotic nature of signals. In particular, the concept of entropy has been applied to study the chaotic nature of various signals. Entropy describes the distribution probability of molecules in a fluid system in physics. During

* Corresponding author.

E-mail address: hinhung@hku.hk (H.h. Sik).

the evolution of a system of many signals, the degree of disorder-ness, which is measured by entropy, increases in general [4]. Since different EEG waveforms represent different states of the electrical activity of the brain, the entropy of EEG signals signifies the chaotic degree of such activity as a whole, which cannot be measured by the traditional spectrum analysis. Particularly, it measures the chaotic level over different scales and hence is more applicable and convenient for analyzing EEG data [4]. Entropy measures have been employed to study AD, epilepsy and anaesthesia in clinical settings [5].

Neurocardiology in medicine suggests a close interplay between the heart and the brain, and cardiac afferent information can affect emotion processing in the brain [6]. There is evidence of meditation-related change in the heart rate variability, when compared to pre-meditation states [7]. Another study found that during mindfulness meditation, there is also quantitative change of EEG and respiration signals [8]. However, previous studies mainly investigate the effect of meditation on the brain or the heart separately. Only a few studies report on better coordination of body and mind after meditation training and mindfulness practices [9,10]. For example, it has been shown that integrative body-mind training (IBMT) increased the correlation between frontal midline theta wave and heart rate variability (HRV), suggesting that a better regulation of autonomic nervous system by midfrontal brain region [10].

The main aims of this investigation are: (i) to measure the power of the 5 standard bands of EEG waveforms, in order to see the changes brought by MBSR. (ii) to measure the wavelet entropy of EEG before and after an eight-week MBSR program, in order to find the difference in chaotic degree as compared to the rest (control) state. (iii) to find out the 3D source positions related to the entropy changes, based on a source analysis of the EEG activity. (iv) to measure wavelet entropy of the heart rate concurrently in order to find the chaotic change, similar to (ii). (v) to find if there is correlation of the entropy changes in the two variables above, in order to see if there is heart–brain entrainment.

2. Materials and methods

2.1. Participants

The research was approved by the local Institutional Review Board (IRB). All participants provided their written informed consent prior to participate in this study. Eleven healthy participants (6 males, 5 females; mean age 35.7 year; 7 Asians and 4 Caucasians) from a local MBSR course voluntarily participated in this study. Each participant was paid 200HKD for transportation fee. All participants were above the undergraduate level. Six of them had some experience of mental exercises such as Yoga, Qigong, or other meditation methods. Nonetheless, they all did not have experience in MBSR and during the experiment, they were requested to only practice mindfulness breathing learned from the MBSR course. The course were taught following the standard MBSR program consisting of one pre-program orientation session, eight weekly classes and one all-day class when the teacher gave direct instructions. Participants also needed to make strong commitment to practice 45 min of MBSR training each day as home assignments for 8 weeks individually, including body scanning and mindful breathing.

Beck Depression Inventory (BDI) was used to exclude participants with depression [11]. The effectiveness of MBSR was measured by Five Facets Mindfulness Questionnaire (FFMQ) [12]. These five facets included non-judging, describing, non-reacting, acting with awareness, and observing. The sub-scores of each facet and a total score were calculated for each participant. The FFMQ was used to measure the subjective experience of mindfulness, and

in the following we also used EEG to measure relatively objective brain waves associated with mindfulness practice.

2.2. Experiment procedure

Ten minutes of eye-closed normal rest EEG was collected for each participant in a quiet room and participants were requested not to ruminate too much or fall asleep. They were also asked to practice mindful breathing for 10 min, following the teaching of the MBSR course. Participants were requested not to ruminate too much or fall asleep, and a post-experiment dialog showed that they could largely follow the procedures. EEG data were recorded at the beginning of the MBSR training (within two weeks), and less than one month after the MBSR course, resulting in two data sets and two conditions. The early-stage MBSR condition was not set before MBSR training but within two-weeks training time. It was to ensure that the trainees knew the right way to practice the MBSR mindful breathing at the time of EEG recording. The sequence of mindfulness state and normal rest state was counter-balanced, with half of the participants did the mindfulness condition first and half did rest condition first, randomly.

2.3. Data acquisition and analysis

The data were acquired by a 128-channel NeuroSCAN system in a quiet room. The sampling rate was 1000 Hz. The original reference point was at left mastoid, and re-referenced to both mastoids when processing the data. The ECG electrodes were placed at the left and right infraclavicular fossae.

The data was processed by EEGlab (see Supplementary materials for details). The spectra of EEG were computed using fast Fourier transform (FFT) and the log powers (dB) of delta (1–4 Hz), theta (4–8 Hz), alpha (8–12 Hz), beta (12–30 Hz), and gamma (30–80 Hz) waves were obtained. In order to evaluate the chaotic degree, wavelet entropy, an entropy of the energies under different scales given by discrete wavelet transform, of the EEG (as well as their estimated cortical sources) and heart rate were then calculated (see Supplementary materials for details on how to calculate wavelet entropy). To explore the relationship between brain and heart during the MBSR practice and normal rest states, we calculated the linear correlation coefficient (Pearson coefficient) between the EEG's and the heart rate's wavelet entropies.

3. Results

All the participants were not depressed according to their BDI scores (<13). The FFMQ data was shown in Supplementary Table 1. No significant change was found between early-stage MBSR and post-stage MBSR stages except for the facet of nonreacting ($p < 0.05$).

There were enhanced Alpha (8–12 Hz) and Beta (12–30 Hz), and reduced delta waves (1–4 Hz) during MBSR practice than during normal rest. The increase of alpha waves was globally significant, especially in the frontal and occipital lobes. The increase of beta wave was mainly in the frontal lobe. Decreased delta wave was at central-parietal areas. However, we did not find significant change between the early-stage and post-stage MBSR training. See Fig. 1 and Supplementary Table 2.

Similar to previous studies [13,14], several regions of interest (ROI) were selected to represent the occipital lobe, the middle frontal lobe and the middle parietal lobe (see Supplementary material for channels defined for these ROIs). Analysis on these ROIs showed that the increase of alpha wave and decrease of delta wave were prominent all the three ROIs, while the increase of beta was mainly in the frontal lobe. See Supplementary Table 2.

Download English Version:

<https://daneshyari.com/en/article/6279842>

Download Persian Version:

<https://daneshyari.com/article/6279842>

[Daneshyari.com](https://daneshyari.com)