



Heterogeneous relationships between abundance of soil surface invertebrates and radiation from Chernobyl

Vladimir Bezrukov^a, Anders Pape Møller^{b,*}, Gennadi Milinevsky^c, Stanislav Rushkovsky^a, Maria Sobol^a, Timothy A. Mousseau^d

^a Department of General and Molecular Genetics, Taras Shevchenko National University of Kyiv, Volodymyrska St. 64, 01601 Kyiv, Ukraine

^b Laboratoire d'Ecologie, Systématique et Evolution, CNRS UMR 8079, Université Paris-Sud, Bâtiment 362, F-91405 Orsay Cedex, France

^c Space Physics Laboratory, Taras Shevchenko National University of Kyiv, Volodymyrska St. 64, 01601 Kyiv, Ukraine

^d Department of Biological Sciences, University of South Carolina, Columbia, SC 29208, USA

ARTICLE INFO

Article history:

Received 5 October 2014

Received in revised form

10 November 2014

Accepted 13 November 2014

Keywords:

Abundance

Chernobyl

Interspecific differences

Radiation

Susceptibility to radiation

ABSTRACT

Although pollution due to nuclear accidents constitutes some of the largest environmental disasters, there is surprisingly little information available on the relationship between abundance of animals and background radiation. We sampled invertebrates with pit-fall traps at Chernobyl in areas differing in background radiation level by more than four orders of magnitude. We obtained samples from 58 traps for 19 taxa of which five showed positive associations (Acari, Araneae, *Formica* sp., Homoptera), while four showed negative associations (Coleoptera, Collembola, *Vespa*, Insecta). These relationships were independent of other environmental factors such as habitat, humidity and ecotype. Estimates of the relationship between abundance and background radiation from the pitfall trap study were similar to those from an independent study based on invertebrates found under slices of wood placed on the ground. The differences in relationship between abundance and radiation among taxa may arise from direct effects of radiation. The findings reported here have implications for choice of animal taxa for efficient monitoring of the biological impact of radiation on animals.

© 2014 Elsevier Ltd. All rights reserved.

1. Introduction

Radioactive material occurs under natural conditions due to the radionuclides contained in rocks and from cosmic rays and such radioactive hotspots show significant negative effects on plants and animals (review in Møller and Mousseau, 2013a,b). Radioactive contamination due to release from nuclear disasters has affected hundreds of thousands of square kilometers across the globe with particularly strong effects having been documented at Chernobyl and Fukushima. Studies have shown that the abundance of organisms is affected by radioactive contaminants. This applies to soil invertebrates (Krivolutski and Pokarzhevsky, 1992; Krivolutski et al., 1999; Maksimova, 2005), bacteria (Zymenko et al., 1995; Romanovskaya et al., 1998; Ragon et al., 2011), spiders (Møller and Mousseau, 2009, 2011; Møller et al., 2012b,c), insects (Møller and Mousseau, 2009, 2011; Møller et al., 2012b,c), birds (Møller and Mousseau, 2007a,b; Møller et al., 2012b,c) and mammals (Møller and Mousseau, 2013a). Such effects of radiation on abundance are

consistent among years and geographical locations (Møller et al., 2011, 2012b), but differ among taxa.

There are significant differences among bird species in susceptibility to radiation in Chernobyl with species on average being negatively affected in terms of abundance, while a few species show no or even a positive effect (Møller and Mousseau, 2007b). Similar heterogeneity has been documented for the relationship between abundance and level of background radiation in Fukushima (Møller et al., 2012b,c). While such heterogeneity in effects of radiation on abundance may be expected simply for sampling reasons, statistically significant heterogeneity beyond sampling suggests that these are inherent species-specific characteristics. Indeed, interspecific differences for birds in frequency of cataracts (Mousseau et al., 2013), reductions in brain size (Møller et al., 2013), frequency of tumors and abnormalities (Møller et al., 2013), aspermy and sperm behavior (Møller et al., 2008, 2014) and levels of antioxidants (Bonisoli-Alquati et al., 2011; Galván et al., 2014) suggest that underlying physiological processes account for such heterogeneity.

Although soil invertebrates, arachnids and insects constitute significant parts of biodiversity, and although small invertebrates are readily sampled in standard pitfall traps, there are so far no pitfall trap studies conducted in Chernobyl or Fukushima. While

* Corresponding author. Tel.: +33 1 69 15 56 88; fax: +33 1 69 15 56 96.
E-mail address: anders.moller@u-psud.fr (A.P. Møller).

decomposition of leaf litter is reduced in contaminated areas around Chernobyl (Mousseau et al., 2014), these effects are not readily interpreted because we do not know to which extent different soil-inhabiting taxa are affected.

The main objective of this study was to assess to what extent the abundance of soil surface invertebrates was affected by radioactive contamination from Chernobyl. First, we sampled invertebrates with pitfall traps to test for differences in susceptibility to radiation among taxa. Second, we identified the taxa that provide the most sensitive indicators of radiation. Third, we cross-validated our estimates of effects of radiation on abundance by comparing our pitfall trap results with those obtained from invertebrates found under slices of pine wood placed across a radiation gradient.

2. Methods

2.1. Sampling sites

We chose six sites that were ecologically similar ranging in level of contamination from 0.013 to 36 mrem/h (Fig. 1). The coordinates of the sites and their characteristics are listed in the electronic supplementary material (Table S1). We report characteristics of these six study sites in Table S1 in terms of radioactivity, soil type, amount of dead organic matter, amount of surface covered with open water as in ponds, and dominant vegetation.

Soil at the traps was classified as sand or ash (ash derived from forest fires) within a radius of 10 m from each of the trap sites. The amount of litter was classified on a three-point scale from 1 – little, over 2 – common to 3 – abundant within a radius of 10 m from the traps. We recorded if there was open water (ponds or puddles) within a distance of 50 m from the traps. Finally, we recorded whether the dominant vegetation was grass, deciduous trees or coniferous trees within a distance of 10 m from the traps.

We considered three different ecotypes based on soil, abundance of litter and other features of habitats (Ingegnoli, 2002; Bastian et al., 2003).

2.2. Cross validation

We cut 5 cm slices from the trunks of four Scots pine (two from an uncontaminated area and two from a highly contaminated area) that were approximately 60 years old. At each of 20 sites varying in radioactive contamination by more than four orders of magnitude we put our one slice of each of the four trees in early June 2013. Each slice was identifiable from a unique numbered metal tag. During early June 2014 we re-visited all 20 sites and recorded on the lower surface facing toward the soil the number of fungi and the animal taxa used in the present study and counted these animals before turning over the slice again for future study. We recorded background radiation at the site of each slice when this was originally put out and one year later as described above. For the five taxa with animals common in the two datasets we estimated the abundance of animals in relation to background radiation using Poisson regression as described below.

2.3. Measuring background radiation levels

We measured radiation levels in the field at ground level at each location where a pitfall trap was deposited using a hand-held dosimeter (Model: Inspector, SE International, Inc., Summertown, TN, USA). We measured levels 2–3 times at each of these locations and averaged the measurements for each location. Such data were correlated with data from governmental measurements at ground level published by Shestopalov (1996), estimated as the mid-point of the ranges in the published maps. These analyses showed a high degree of consistency between the two methods (Møller and

Mousseau, 2007a). Radiation levels vary by several orders of magnitude at a scale of 1 km due to heterogeneity in deposition of radioactive material after the Chernobyl accident (Shestopalov, 1996).

2.4. Pitfall traps and handling

We used 500 ml plastic containers with an inner diameter of 90 mm and 25% ethylene glycol as a preservative. They were placed in the ground so the top of the container was at the level of the soil surface and covered with CDs (compact discs) for protection from rain and solar light. The distance from the soil surface to the CD was about 2 cm. The distance among traps was 2 m, and a total of 20 traps per site we used in this survey. The traps were placed along two intersecting transects to form a “+” and oriented toward north, south, east and west. The traps were placed on 25 August 2007 and contents were collected on 8 September 2007.

2.5. Identification

Container contents were sorted under a stereo binocular microscope and transferred to 1.5 ml eppendorf tubes with 96% ethanol. The organisms were identified to order using Chinery (1993, 2007), Gibbons (1995), Gilarov (1964) and a mite identification web site (<http://www.lucidcentral.org/keys/v3/mites/Invasive.Mite-Identification/>).

Some samples were excluded from the analysis because they were destroyed by wild animals (the containers were dry or removed). One trap (site RF03) was excluded because it contained an enormous number of ants (667) and collembolas (26,922) that exceeded the general sample by more than three standard deviations. This aberrant trap was situated very close to an ants' nest hence explaining the abundance of ants.

2.6. Statistics

Because observations of animals in pitfall traps basically are count data, they have to be treated as having a Poisson distribution. Therefore, we used generalized linear mixed models with the abundance of animals as the response variable and background radiation and the first two principal component scores reflecting environmental variables (Table S2). None of these components were statistically significant predictors and hence they were excluded from the final statistical models.

We reduced the five environmental variables using principal component analysis. This analysis was based on the correlation matrix for the five environmental variables using varimax rotation. The principal component analysis revealed two factors with eigenvalues above one accounting for 60% and 25% of the variance, respectively. The first factor was strongly positively correlated with soil and ash, dead organic matter, presence of water and ecotype. The second factor showed a positive loading for presence of trees and ecotype. We subsequently extracted principal component scores for the 58 pitfall trap sites and used these scores as additional predictor variables in the analyses. All analyses were made with JMP (SAS, 2012). We report means \pm 1 SE.

3. Results

3.1. Summary statistics

We found animals of three orders of the class Arachnida, eleven orders (ten orders and one genus – Collembola) of Insecta and one order (Myriapoda) of the class Chilopoda (Table 1). In total, this study was based on 12,049 animals (Table 1) ranging in abundance

Download English Version:

<https://daneshyari.com/en/article/6294513>

Download Persian Version:

<https://daneshyari.com/article/6294513>

[Daneshyari.com](https://daneshyari.com)