



## Radiofrequency radiation injures trees around mobile phone base stations



Cornelia Waldmann-Selsam<sup>a</sup>, Alfonso Balmori-de la Puente<sup>b</sup>, Helmut Breunig<sup>c</sup>, Alfonso Balmori<sup>d,\*</sup>

<sup>a</sup> Karl-May-Str. 48, 96049 Bamberg, Germany

<sup>b</sup> C/Navarra, 1 5<sup>o</sup>B, 47007 Valladolid, Spain

<sup>c</sup> Baumhofstr. 39, 37520 Osterode, Germany

<sup>d</sup> Consejería de Medio Ambiente, Junta de Castilla y León, C/Rigoberto Cortejoso, 14 47071 Valladolid, Spain

### HIGHLIGHTS

- High frequency nonionizing radiation is becoming increasingly common.
- This study found a high level of damage to trees in the vicinity of phone masts.
- Deployment has been continued without consideration of environmental impact.

### GRAPHICAL ABSTRACT

## Bernartzky (1986), revisited:



### ARTICLE INFO

#### Article history:

Received 6 June 2016

Received in revised form 19 July 2016

Accepted 6 August 2016

Available online xxxx

Editor: D. Barcelo

#### Keywords:

Electromagnetic radiation

Effects on trees

Phone masts

Radiofrequencies

### ABSTRACT

In the last two decades, the deployment of phone masts around the world has taken place and, for many years, there has been a discussion in the scientific community about the possible environmental impact from mobile phone base stations. Trees have several advantages over animals as experimental subjects and the aim of this study was to verify whether there is a connection between unusual (generally unilateral) tree damage and radiofrequency exposure. To achieve this, a detailed long-term (2006–2015) field monitoring study was performed in the cities of Bamberg and Hallstadt (Germany). During monitoring, observations and photographic recordings of unusual or unexplainable tree damage were taken, alongside the measurement of electromagnetic radiation. In 2015 measurements of RF-EMF (Radiofrequency Electromagnetic Fields) were carried out. A polygon spanning both cities was chosen as the study site, where 144 measurements of the radiofrequency of electromagnetic fields were taken at a height of 1.5 m in streets and parks at different locations. By interpolation of the 144 measurement points, we were able to compile an electromagnetic map of the power flux density in Bamberg and Hallstadt. We selected 60 damaged trees, in addition to 30 randomly selected trees and 30 trees in low radiation areas ( $n = 120$ ) in this polygon. The measurements of all trees revealed significant differences between the damaged side facing a phone mast and the opposite side, as well as differences between the exposed side of damaged trees and all other groups of trees in both sides. Thus, we found that side differences in measured values of power flux density corresponded to side differences in damage. The 30 selected trees in low radiation areas (no visual

\* Corresponding author.

E-mail addresses: [balmara@jcyL.es](mailto:balmara@jcyL.es), [abalmori@ono.com](mailto:abalmori@ono.com) (A. Balmori).

contact to any phone mast and power flux density under  $50 \mu\text{W}/\text{m}^2$ ) showed no damage. Statistical analysis demonstrated that electromagnetic radiation from mobile phone masts is harmful for trees. These results are consistent with the fact that damage afflicted on trees by mobile phone towers usually start on one side, extending to the whole tree over time.

© 2016 Elsevier B.V. All rights reserved.

## 1. Introduction

For many years, there has been a discussion in the scientific community about whether artificial radiofrequency radiation has harmful effects on living organisms and, more specifically, on the environmental impact from mobile phone base stations (Panagopoulos et al., 2016). Trees have several advantages over animals as experimental subjects: they are continuously exposed to radiation in a constant orientation in the electromagnetic field due to their inability to move (Vian et al., 2016). Additionally, it is possible to easily document changes over time, such as disturbed growth, dying branches, and premature colour change of leaves. Moreover, the damage to trees is objective and cannot be attributed to psychological or psychosomatic factors.

Plants are specialized in the interception of electromagnetic radiation (light) but radiofrequency radiation impact on plants, which is becoming common in the environment because of the exponential use of mobile phone technology, has received little attention and his physiological effect has long been considered negligible.

Since the mid-twentieth century, several researchers have investigated the effects of electromagnetic radiation on plants, both in the laboratory (Kiepenheuer et al., 1949; Brauer, 1950; Harte, 1950, 1972; Jerman et al., 1998; Lerchl et al., 2000; Sandu et al., 2005; Roux et al., 2006, 2008; Sharma et al., 2009; Tkalec et al., 2005, 2009; Beaubois et al., 2007; Kundu and IEEE, 2013; Pesnya and Romanovsky, 2013; Cammaerts and Johansson, 2015; Grémiaux et al., 2016; Vian et al., 2016), and in nature (field observations) (Bernatzky, 1986; Volkrodt, 1987, 1991; Selga and Selga, 1996; Balodis et al., 1996; Haggerty, 2010). Both kinds of study have frequently found pernicious effects.

Around the world, phone masts have been deployed in the last two decades everywhere. Preliminary published studies have indicated deleterious effects of radiofrequency radiation on trees (Balmori, 2004; Van't Wout, 2006; Schorpp, 2011; Waldmann-Selsam, 2007; Waldmann-Selsam and Eger, 2013), cautioning that research on this topic is extremely urgent (Balmori, 2015). However, these early warnings have had no success and deployment has been continued without consideration of environmental impact.

In a review of the effects of environmental microwaves on plants (Jayasanka and Asaeda, 2013), it was indicated that effects depend on the plant family and the growth stage, as well as the exposure duration, frequency, and power density. This review concluded that most studies that address the effects of microwaves on animals and plants have documented effects and responses at exposures below limits specified in the electromagnetic radiation exposure guidelines and it is therefore necessary to rethink these guidelines (Jayasanka and Asaeda, 2013).

Since 2005, on the occasion of medical examinations of sick residents living near mobile phone base stations, changes in nearby trees (crown, leaves, trunk, branches, growth...) were observed at the same time as clinical symptoms in humans occurred. Since 2006 tree damages in the radiation field of mobile phone base stations were documented (<http://kompetenzinitiative.net/KIT/KIT/baeume-in-bamberg/>). In the radio shadow of buildings or that one of other trees, the trees stayed healthy.

Additionally, unilateral crown damage, beginning on the side facing an antenna, pointed to a possible link between RF-EMF (Radiofrequency Electromagnetic Fields) and tree damage. We carried out measurements on both sides of unilaterally damaged trees. Most of the trees had been exposed to RF-EMF for at least five years. Each time we

found considerable differences between the measured values on the damaged and on the healthy side.

The aim of the present study was to verify whether there is a connection between unusual (generally unilateral) tree damage and radiofrequency exposure.

## 2. Materials and methods

The official information of 65 mobile phone sites in the neighbouring cities Bamberg and Hallstadt was extracted from the EMF database (EMF-Datenbank) of the German Federal Network Agency (Bundesnetzagentur, in March 2011 and October 2015). Each site certificate ("Standortbescheinigung") provides information on the mounting height of antennas, the number and main beam direction of the sector antennas, the number of omnidirectional antennas (ND), the number of other transmitters, as well as the horizontal and vertical safety distances. The current specifications of the transmission facilities are available at: <http://emf3.bundesnetzagentur.de/karte/Default.aspx>

On most of the 65 mobile phone sites several sector antennas emitting RF-EMF with differences in frequency, modulation and other physical characteristics are installed (GSM 900, GSM 1800, UMTS, LTE (4th generation), TETRA). In 2011 there was a total of 483 sector antennas, in 2015 a total of 779 sector antennas.

Numerical code, address and UTM 32N coordinates for the 65 Mobile phone (base stations) sites in Bamberg and Hallstadt are shown in Table 1.

Between 2006 and 2015 there was observation and documentation of tree damages. There were some preliminary measurements on both sides of unilaterally damaged trees and approximately 700 trees in Bamberg and Hallstadt were visited. The condition of numerous trees has been documented in photographs. The photographs record the state of trees showing damage patterns not attributable to diseases, pests, drought or other environmental factors in order to monitor damage and growth over several years (in 2006, Olympus FE-100 was used; since 2007, Panasonic DMC-FZ50 was used).

In 2015 we selected a polygonal study site, with an approximate area of  $30 \text{ km}^2$ , which includes partial municipalities of Bamberg and Hallstadt ( $70 \text{ km}^2$ ). The study area with the location of the phone masts in the layer of natural areas and municipalities is shown in Fig. 1. In this area, different measurements (see below) were done both for having a radiation map and for knowing which are the incident power densities beside different trees. In spite of the fact that measurements are changing continuously, they do not show significant differences between times (own data, see below).

In this polygon, we performed 144 measurements of the radiofrequency electromagnetic fields at a height of 1.5 m at different points in the city. These measurements were taken in streets and parks and allowed the preparation of an electromagnetic map of Bamberg and Hallstadt with their interpolation. The measurements were carried out with an EMF-broadband analyzer HF 59B (27–3300 MHz) and the horizontal-isotrope broadband antenna UBB27\_G3, (Gigahertz Solutions). Measurements of the sum peak values of power flux density were in  $\mu\text{W}/\text{m}^2$ , which can be converted in V/m.

In general, a sector antenna covers an angle of  $120^\circ$  and the radiation of the sector antennas is distributed in main and secondary beams, bundled vertically and horizontally. The high-frequency emissions are reflected/diffracted and/or absorbed by buildings and trees. Therefore,

Download English Version:

<https://daneshyari.com/en/article/6320635>

Download Persian Version:

<https://daneshyari.com/article/6320635>

[Daneshyari.com](https://daneshyari.com)