



Short communication

Metal exposure influences the melanin and carotenoid-based colorations in great tits



M. Giraudeau ^{a,*}, F. Mateos-Gonzalez ^{b,1}, J. Cotín ^c, E. Pagani-Nuñez ^{b,2}, A. Torné-Noguera ^{b,3}, J.C. Senar ^b

^a Biological Sciences A08, University of Sydney, Sydney, NSW, Australia


^b Evolutionary and Behavioural Ecology Associate Research Unit, CSIC, Natural History Museum of Barcelona, P. Picasso s/n, 08003 Barcelona, Spain

^c Department Biologia Animal, Facultat de Biologia, Universitat de Barcelona, Diagonal 645, 08028 Barcelona, Spain

HIGHLIGHTS

- We measured the concentrations of metals in feathers collected on great tits.
- Melanin pigmentation was positively associated with the concentration of copper.
- Melanin pigmentation was negatively related with the concentration of chromium.
- Carotenoid-based coloration was negatively related with the concentration of mercury.

GRAPHICAL ABSTRACT



ARTICLE INFO

Article history:

Received 9 March 2015

Received in revised form 4 June 2015

Accepted 4 June 2015

Available online xxxx

Editor: D. Barcelo

Keywords:

Metal pollution

Plumage coloration

Melanin

Carotenoid

ABSTRACT

Metals are naturally found in the environment but are also emitted through anthropogenic activities, raising some concerns about the potential deleterious effects of these elements on wildlife. The potential effects of metals on bird coloration have been the focus of several recent studies since animal colored-signals often reflect the physiology of their bearers and are thus used by animals to assess the quality of another individual as a mate or competitor. These studies have shown that the melanin pigmentation seems to be positively associated and the carotenoid-based coloration negatively associated with metal exposure in wild birds. Although these studies have been very useful to show the associations between metal exposure and coloration, only few of them have actually quantified the levels of metal exposure at the individual level; always focusing on one or two of them. Here, we measured the concentrations of eight metals in great tits' feathers and then assessed how these levels of metals were associated with the carotenoid and melanin-based colorations. We found that the melanin pigmentation was positively associated with the copper concentration and negatively correlated with the chromium concentration in feathers. In addition, we have shown that the carotenoid-based coloration was negatively

* Corresponding author.

E-mail address: jcsenar@bcn.cat (J.C. Senar).

¹ Current address: Department of Integrative Biology, University of Texas, Austin, TX 78712, USA.

² Current address: Behavioral and Community Ecology, Conservation Biology Group, College of Forestry, Guangxi University, No. 100 Daxue Road, Nanning, Guangxi 530005, People's Republic of China.

³ Current address: CREAF, 08193 Cerdanyola del Vallès, Spain.

associated with the feather's mercury concentration. This study is the first one to identify some metals that might affect positively and negatively the deposition of melanin and carotenoid into the plumage of wild birds.

© 2015 Elsevier B.V. All rights reserved.

1. Introduction

Metals are naturally found in the environment (Nriagu, 1989) but are also emitted through anthropogenic activities (Azimi et al., 2005; Roux and Marra, 2007). As a consequence, the levels of metals have increased in urban areas raising some concerns about the potential deleterious effects of these elements on wildlife (Dauwe et al., 2004; Hsu et al., 2006). Metals are commonly absorbed by wild animals through their diet and some of the metals (zinc, iron, copper, cobalt and manganese) are involved in the control of various metabolic and signaling pathways (Bogden and Klevay, 2000; Valko et al., 2005). However, at high concentrations, these metals are known to have toxic effects, partly by the induction of high levels of oxidative stress (Ercal et al., 2001). Other metals like cadmium and lead do not have any known biological function but have various toxic effects (Ercal et al., 2001).

Animal colored-signals often reflect the recent nutrition, health, endocrinology, oxidative stress, or other resource-based attributes of their bearers (McGraw, 2006) and are thus involved in sexual contexts because they can be used by animals to assess the quality of another individual as a mate (Hill, 2006) or competitor (Senar, 2006). The potential effects of metals on animal coloration have been the focus of several studies in the last 15 years, with a special interest on the relationships between exposure to these metals and the deposition of the two most common pigments of the vertebrate tegument: the carotenoid and melanin pigments (Chatelain et al., 2014; Dauwe and Eens, 2008; Geens et al., 2009; McGraw, 2003, 2007, 2008, Niecke et al., 2003; Roulin et al., 2006). So far, melanin pigmentation seems to be positively associated with metal exposure in wild birds since (1) more pigmented individual feral pigeons (*Columba livia*) are found in urban compared to rural environments (Jacquin et al., 2013; Obukhova, 2007), (2) darker feral pigeons had higher concentrations of zinc in their feathers compared to paler ones after one year in standardized conditions (Chatelain et al., 2014), (3) copper deficiency influences plumage color in poultry species (Leeson, 2009; Leeson and Walsh, 2004) and (4) melanin pigmentation in great tits (*Parus major*) increases along a gradient of metal pollution caused by a lead smelter in Belgium (Dauwe and Eens, 2008). Two main hypotheses have been proposed to explain this association. First, the tyrosinase enzyme activity, a crucial component of the melanogenesis process, might be stimulated in metal-rich environment due to its copper-containing molecular structure (McGraw, 2003; Protá, 1993). Second, highly pigmented individuals might be more able to store metals in feathers due to the capacity of melanin pigments to bind metal ions (Chatelain et al., 2014; McGraw, 2003).

To the contrary, the carotenoid-based coloration was negatively affected along the same gradient of metal pollution Belgium (Dauwe and Eens, 2008; Eeva et al., 1998; Geens et al., 2009) and in urban areas (Isaksson et al., 2007). In this case, the detrimental effect of metal exposure on coloration might be explained by a decrease in the dietary availability of carotenoid pigments in polluted environments (Isaksson and Andersson, 2007) and/or the allocation of carotenoid pigment to the antioxidant and immune systems, at the detriment of coloration, since pollutants such as metals are known to induce oxidative stress in animals (Blount et al., 2003; Faivre et al., 2003; McGraw and Ardia, 2003).

Although the studies published on this topic have been very useful to show the associations between metal exposure and both melanin and carotenoid-based colorations, only few of them have actually quantified

the levels of metal contamination at the individual level; always focusing on one or two metals (Chatelain et al., 2014, Niecke et al. 2003, Roulin et al. 2006). Here, we measured the concentrations of eight metals in great tits' feathers (metal concentrations in feathers are known to reflect blood metal levels during feather growth, Burger and Gochfeld, 1992; Janssens et al., 2001) and then assessed how these levels of metals were associated with the carotenoid and melanin based colorations, allowing us to determine which specific metals might have a positive or a negative effect on both colorations.

2. Material and methods

2.1. Field procedure

From 9th February to 6th May 2009, we use baited funnel traps to capture 32 great tits (13 females and 19 males) in city parks in Barcelona (Spain, see Björklund et al. (2010) for a detailed description of the parks). At capture, sex and age were determined according to Svensson (1992) and the second left and right outermost tail rectrices (R5 and L5) were collected for later analyses of pollutant levels (see below). Feathers were stored in dry and dark conditions in individually marked polyethylene bags.

We also photographed each bird to measure the size of the melanin-based black breast-tie using Image Tool 3.0© (University of Texas, USA, Figuerola and Senar, 2000). We drew a line around the black-tie and then measured the black area within 3 cm, starting 2 cm after the junction of the black-tie and the white cheek (Järvi and Bakken, 1984; Norris, 1990, 1993; Pöysä, 1988; Quesada and Senar, 2007; Wilson, 1992). The use of digital photography standardized to a metrical reference allowed a high repeatability in the measurement of the black-tie area (Figuerola and Senar, 2000).

The breast carotenoid-based coloration (brightness, hue and chroma) was measured with a portable colorimeter Minolta CR200 (Minolta Corporation 1994). Three color measurements were taken at 90°, making contact with the surface of the feathers patch, and the mean was then calculated for each of the three components of the carotenoid-based coloration (brightness, hue and chroma). Repeatability of these measures was high ($r_1 = 0.85-0.92$, $p < 0.001$) (Figuerola et al., 1999). Values of brightness, chroma and hue were collapsed into a single variable using a Principal Components Analysis. The first eigenvalue from PCA ($\lambda = 1.89$) accounted for 63% of the variance in plumage coloration. The three values (brightness, hue and chroma) of plumage coloration loaded positively to the first component (PC1), with all the values > 0.70 (Brightness = 0.79, Chroma = 0.72, Hue = 0.86). Hence, PC1 is hereafter called yellow coloration. We have shown elsewhere that there is no correlation between tie size and the carotenoid-based coloration in great tits (Senar et al., 2003).

2.2. Trace element analysis

Feathers were first rinsed with a 0.25 M NaOH solution for 1 min and then with deionized water for another minute. Feathers (ca. 20 mg) were oven dried at 60° for 24 h and then digested in H₂NO₃ (0.5 ml) and H₂O₂ (0.5 ml) in Savilles Teflon digestion vessels for 12 h at 100 °C. Levels of mercury, copper, lead, chrome, arsenic, cadmium, antimony and zinc were measured using a ICP-MS Perkin Elmer ELAN 6000. Accuracy of analysis was checked by measuring certified reference tissue: human hair (BCR 397, Community Bureau of Reference, Commission of the European Communities). All trace element concentrations were expressed on a dry weight basis (ng·g⁻¹ dw). Mean recoveries

Download English Version:

<https://daneshyari.com/en/article/6325775>

Download Persian Version:

<https://daneshyari.com/article/6325775>

[Daneshyari.com](https://daneshyari.com)