



## Short Communication

## Obtainment of a highly concentrated pancreatic serine proteases extract from bovine pancreas by precipitation with polyacrylate



Julia Lombardi, Nadia Woitovich Valetti, Guillermo Picó, Valeria Boeris\*

Laboratorio de Físicoquímica Aplicada a Bioseparación, Facultad de Ciencias Bioquímicas y Farmacéuticas, CONICET, Universidad Nacional de Rosario, Suipacha 570 (S2002RLK), Rosario, Argentina

## ARTICLE INFO

## Article history:

Received 4 March 2013

Received in revised form 22 May 2013

Accepted 24 May 2013

Available online 4 June 2013

## Keywords:

Bovine pancreas proteases

Serine proteases

Trypsin

Chymotrypsin

Polyelectrolyte

## ABSTRACT

Serine proteases have wide application in leather, food, meat and soap powder industries, among others. There are a lot of methodologies to obtain them in large quantities but most of them are expensive or contaminating. We then propose an economical and environmentally friendly method in which proteases are separated from a crude fresh bovine pancreas homogenate using precipitation with polyacrylate, a commercially available negatively charged and weak polyelectrolyte. The zymogens of the serine proteases were activated prior to precipitation by the addition of trypsin. The proteases were precipitated by adding polyacrylate at pH 4.50 to the pancreas homogenate. Under these conditions, serine proteases are positively charged and form non-soluble complexes with the negatively charged polyacrylate. The purity of proteases was increased 5-fold with a recovery of 33% under the best conditions tested. The volume of the final product was decreased to 5% of the feedstock, in order to concentrate the sample up to 20 times. The proposed method removed up to 96% of the contaminant proteins.

© 2013 Elsevier B.V. All rights reserved.

## 1. Introduction

Meat industries are very important in our country. Therefore, great amounts of meat waste are produced. One of these products is bovine pancreas: thousands of kilos of fresh pancreas are discarded daily, contributing to environmental pollution. However, there is an increasing commercial interest in the pancreas since it is very rich in enzymes such as proteases of wide application in numerous biotechnological processes [1–3]. Proteases are among the most important hydrolytic enzymes and have been studied extensively [4].

Due to their activity and stability at alkaline pH, serine proteases are of considerable interest and have applications in different types of industries [5]: pharmaceuticals, diagnostic reagents, leather, food, meat and soap powder industries, among others.

The three serine proteases present in bovine pancreas are trypsin (Tryp), chymotrypsin (ChTRP) and elastase. In order to use these enzymes it is necessary to isolate them and to preserve them properly to keep their functionality. There are a lot of methodologies to obtain these serine proteases in large quantities but most of them are expensive or generate byproducts that are not suitable to be discarded [6–10]. We have proposed different methods to isolate ChTRP [11–14], Tryp [15,16] and their zymogens [17,18]. However, it is also interesting to obtain a concentrated extract with

protease activity (containing a mixture of the above mentioned) to be applied to many industries including the production of food products, leather, meat, soap powder, waste management and silver recovery where the different specificities are not needed.

Conventional multistep downstream processing of enzymes is often time consuming, labor intensive, requiring huge hold-up volumes to operate in batch or semibatch mode and usually accounts for up to 80% of the total process costs [6,18–20]. Some of the traditional methods suitable to be applied in scaling up are ion exchange adsorption [21] or chromatography [22]. However, chromatography is the single largest cost center in downstream processing and also the yield of a chromatography is usually low. Thus, alternatives to chromatography are an attractive option even if only a reduction in the extent of use of packed beds can be realized.

The introduction of expanded bed adsorption has allowed early process integration by protein sequestration from a crude feedstock, but reduces the overall system dynamic binding capacity for the target product due to drawback in maintaining an appropriate (close to plug-flow) hydrodynamic condition of the fluidized bed. The chromatographic resins exhibit several major limitations due to slow intra-particle diffusion within the porous beads which tend to limit the dynamic binding capacity of the resins to capture desired target product, the column design reduces throughput as a result of the increased pressure drop at higher flow rates and high material and operational cost.

As a consequence, the separation and purification schemes applied for the purification of enzymes require alternative

\* Corresponding author. Fax: +54 (0341) 480 4598.

E-mail address: [valeriaboeris@conicet.gov.ar](mailto:valeriaboeris@conicet.gov.ar) (V. Boeris).

operations to replace traditional packed or expanded-bed column resins. It is necessary to perform a biotechnological process consisting of as few steps as possible, to reduce the total mass and volume of the sample to obtain an extract rich in enzymes that can be freeze-dried [23,24].

Serine proteases have basic isoelectric pH values: 10.4 (Tryp), 8.7 (ChTRP) and 9.1 (elastase). This property opens up the possibility of using a purification technique which makes use of the positive charge of these enzymes in acid media. Charged proteins interact with polyelectrolytes (PE) to form soluble or non-soluble complexes, according to the experimental conditions of the medium. By changing these conditions, such as pH or ionic strength, the protein can be released, thus keeping its secondary and tertiary structure as well as its biological activity. Synthetic polyelectrolytes such as polyacrylate (PAA) have been used to precipitate proteases as an isolation method. PAA is not toxic; it is used in the pharmaceutical industry as a basis for preparing different edible products.

In this study, we proposed the precipitation of proteases with PAA from fresh pancreas as a generic, simple, low-cost, fast and scalable basic operation, which may be useful as first step in a purification method to concentrate the main serine proteases from its natural source, the bovine pancreas.

## 2. Materials and methods

### 2.1. Chemicals

Crystallized hemoglobin, polyacrylate sodium salt (35% w/v), benzyl-L-tyrosine ethyl ester (BTEE), crystallized Tryp and ChTRP were purchased from Sigma–Aldrich and used without previous purification. All others reagents were also analytical grade.

### 2.2. Bovine pancreas homogenate preparation

The pancreas from a recently killed bovine was removed, washed with isotonic saline solution, cut in small pieces, mixed with  $\text{CaCl}_2$  solution in final concentration of 50 mM and homogenized for 5 min. The resulting homogenate was divided in aliquots and frozen at  $-30^\circ\text{C}$ .

### 2.3. Serine proteases activation from fresh homogenate pancreas

Since serine proteases are produced as zymogens in the pancreas (chymotrypsinogen, trypsinogen and proelastase), an activation step is required. The activation of zymogens was initiated by adding a small aliquot of Tryp (0.0001%, w/w) to the pancreas homogenate (medium: 45 mM  $\text{CaCl}_2$ , 90 mM Tris–HCl buffer, pH 8.20). The time required to complete the activation process was determined by measuring the protease activity at different intervals until a maximal value was reached.

### 2.4. Solubility diagram of polyacrylate–proteases mixtures

Turbidity (absorbance at 420 nm) of solutions of 0.5 g/L PAA and different final concentrations of mixtures of Tryp and ChTRP (1:1 weight ratio) was measured and plotted against pH. The pH variations were obtained by the addition of NaOH or HCl aliquots and leaving the system to equilibrate before measuring the turbidity. These titration curves were performed in order to estimate the pH at which the system composed by proteases and PAA becomes insoluble and to analyze how the different concentrations of proteases affect the soluble–insoluble pH range.

### 2.5. Turbidimetric titration curves of serine proteases with PAA

Solutions of different concentrations of Tryp and ChTRP (1:1 weight ratio) in 50 mM acetate–phosphate buffer were titrated at  $25^\circ\text{C}$  with a 50 g/L PAA solution. The pH of all the solutions was properly adjusted to avoid changes during titration. The increase in the absorbance at 420 nm of the mixture is related to the PAA–protease insoluble complexes formation and was plotted vs. the total concentration of PAA in the tube. The amount of proteases both in the supernatant and in the redissolved precipitate was determined by measuring their proteolytic activity.

### 2.6. Determination of the ionic strength effect on serine protease complex solubility

Mixtures of proteases (0.5 mg/mL) and PAA (0.5 g/L) at three different pH values: 3.50, 4.00 and 4.50 were prepared. Each mixture was added with NaCl and the turbidity of the system was measured.

### 2.7. Determination of the effect of pH on the proteases precipitation effectiveness

PAA (0.5 g/L) was added to bovine pancreas homogenate at 3 different pH values: 3.50, 4.00 and 4.50. After a 1 h incubation, the decanted precipitate was separated from the supernatant and redissolved by addition of Tris– $\text{CaCl}_2$  buffer pH 8.20. The proteolytic activity was determined in the redissolved fraction. To calculate the percentage of recovery of proteases, the proteolytic activity in the bovine pancreas homogenate was also determined.

### 2.8. Determination of total protein concentration

Total protein concentration was determined using the bicinchoninic acid assay [25]. A fresh standard working reagent (SWR) was prepared mixing 100 vol of reagent A (Bicinchoninic acid solution purchased from Sigma–Aldrich) with 2 vol of reagent B (4%  $\text{CuSO}_4$  solution prepared from  $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$  purchased from Sigma–Aldrich). A volume of 50  $\mu\text{L}$  of protein solution (maximum concentration of 1 mg/mL) was added to 1 mL of SWR. The tubes were incubated at  $37^\circ\text{C}$  for 30 min. After leaving them to cool at room temperature, the absorbance was measured at 562 nm using a 1 cm of path length cell. The calibration curve was performed using dilutions of a standard BSA solution of 1 mg/mL.

### 2.9. Determination of protease activity

Protease enzymatic activity was estimated using hemoglobin as substrate. 250  $\mu\text{L}$  of hemoglobin solution (50 g/L of hemoglobin in HCl pH 2.00) and 250  $\mu\text{L}$  of 400 mM Tris– $\text{CaCl}_2$  buffer pH 8.20 were mixed in test tubes and added with an appropriate volume of the sample. After 30 min of incubation at  $37^\circ\text{C}$ , 500  $\mu\text{L}$  of 100 g/L TCA was added. The suspension was centrifuged at 2000g for 15 min. The supernatant was separated and its absorbance at 280 nm was measured to determine the free tyrosine residues.

ChTRP assay is based on the hydrolysis of BTEE [6]. The reaction rate was determined by hydrolysis of 0.6 mM of BTEE in 100 mM Tris– $\text{CaCl}_2$  buffer pH 8.20. The production of benzyl–tyrosine was measured by monitoring the increase in Abs at 256 nm every 2 s for 3 min. One ChTRP unit is defined as 1  $\mu\text{mol}$  of substrate hydrolyzed per minute of reaction and was calculated with the following equation:  $U \text{ (}\mu\text{mol/min)} = (\Delta A_{256 \text{ nm/min}}) \times 1000 \times 1/964$ , where 964 is the benzyl–tyrosine molar extinction coefficient.

Download English Version:

<https://daneshyari.com/en/article/641646>

Download Persian Version:

<https://daneshyari.com/article/641646>

[Daneshyari.com](https://daneshyari.com)