



On rainbow matchings in bipartite graphs

Ron Aharoni¹

*Department of Mathematics
Technion
Haifa, Israel*

Eli Berger²

*Department of Mathematics
Haifa University
Haifa, Israel*

Dani Kotlar³

*Department of Computer Science
Tel-Hai College
Upper Galilee, Israel*

Ran Ziv⁴

*Department of Computer Science
Tel-Hai College
Upper Galilee, Israel*

Abstract

We present recent results regarding rainbow matchings in bipartite graphs. Using topological methods we address a known conjecture of Stein and show that if $K_{n,n}$ is partitioned into n sets of size n , then a partial rainbow matching of size $2n/3$ exists. We generalize a result of Cameron and Wanless and show that for any n

matchings of size n in a bipartite graph with $2n$ vertices there exists a full matching intersecting each matching at most twice. We show that any n matchings of size approximately $3n/2$ have a rainbow matching of size n . Finally, we show the uniqueness of the extreme case for a theorem of Drisko and provide a generalization of Drisko's theorem.

Keywords: partial rainbow matching, full rainbow matching, bipartite graph, Ryser-Brualdi Conjecture, Stein's conjecture

1 The case of n sets of size n

Given sets F_1, F_2, \dots, F_n of edges in a graph, a (*partial*) *rainbow matching* is a partial choice function on the F_i s whose range is a matching. If the rainbow matching represents all of the F_i s, then it is a *full rainbow matching*.

A known conjecture of Ryser and Brualdi [10,16,17] states that any n matchings F_1, F_2, \dots, F_n of size n that form a partition of $K_{n,n}$ have a partial rainbow matching of size $n - 1$. The best result so far towards proving this conjecture belongs to Hatami and Shor [14] who showed that in any such case a partial rainbow matching of size $n - 11 \log_2^2 n$ exists.

The Ryser-Brualdi conjecture can be generalized in different ways. We may ease the requirement that the matchings F_1, F_2, \dots, F_n form a partition of $K_{n,n}$ [1]:

Conjecture 1.1 *Any n matchings of size n in a bipartite multigraph have a partial rainbow matching of size $n - 1$.*

The best result in this direction is due to Woolbright [18] and Brouwer, de Vries and Wieringa [9] who showed (essentially) that a rainbow matching of size $n - \lfloor \sqrt{n} \rfloor$ exists.

Given sets F_1, F_2, \dots, F_n , if a matching of size n in their multiset union intersects each F_i at most twice, we call it a *half-rainbow matching*. Cameron and Wanless [11] showed that in the Ryser-Brualdi setup (that is, when the matchings F_1, F_2, \dots, F_n form a partition of $K_{n,n}$) a half-rainbow matching of size n exists. We generalize the Cameron-Wanless result to the case of any n matchings, namely,

¹ Email: raharoni@gmail.com

² Email: berger.haifa@gmail.com

³ Email: dannykotlar@gmail.com

⁴ Email: ranzivziv@gmail.com

Download English Version:

<https://daneshyari.com/en/article/6423637>

Download Persian Version:

<https://daneshyari.com/article/6423637>

[Daneshyari.com](https://daneshyari.com)