


# Accepted Manuscript

Directed cell growth in multi-zonal scaffolds for cartilage tissue engineering

Sandra Camarero-Espinosa, Barbara Rothen-Rutishauser, Christoph Weder, E.Johan Foster



PII: S0142-9612(15)00784-X

DOI: [10.1016/j.biomaterials.2015.09.033](https://doi.org/10.1016/j.biomaterials.2015.09.033)

Reference: JBMT 17083

To appear in: *Biomaterials*

Received Date: 9 June 2015

Revised Date: 22 September 2015

Accepted Date: 23 September 2015

Please cite this article as: Camarero-Espinosa S, Rothen-Rutishauser B, Weder C, Foster EJ, Directed cell growth in multi-zonal scaffolds for cartilage tissue engineering, *Biomaterials* (2015), doi: 10.1016/j.biomaterials.2015.09.033.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## **Directed cell growth in multi-zonal scaffolds for cartilage tissue engineering**

Sandra Camarero-Espinosa<sup>1\*</sup>, Barbara Rothen-Rutishauser<sup>1</sup>, Christoph Weder<sup>1</sup> and E. Johan Foster<sup>1,2\*</sup>

<sup>1</sup>*Adolphe Merkle Institute, University of Fribourg, Chemin des Verdiers 4, 1700 Fribourg, Switzerland.*

<sup>2</sup>*Virginia Tech, Macromolecules and Interfaces Institute, Virginia Tech Center for Sustainable Nanotechnology, Department of Materials Science & Engineering, 445 Old Turner Street, 213 Holden Hall, Blacksburg VA 24061, USA.*

### **ABSTRACT**

Articular cartilage serves as a low-friction cushion in synovial joints and is vital for mammalian skeletal movements. Due to its avascular nature and the low cell density, the tissue has a limited ability to regenerate, and damage due to injury, wear and tear, or disease usually requires surgical intervention. While articular cartilage had been predicted to be one of the first tissues to be successfully engineered, it proved to be challenging to reproduce the complex architecture and biomechanical properties of the native tissue. Here we report the fabrication of multi-layer polymer nanocomposite scaffolds that mimic the structural design, chemical cues, and mechanical characteristics of mature articular cartilage. These scaffolds guide the morphology, orientation, and phenotypic state of cultured chondrocytes in a spatially controlled manner, support the growth of tissue with features that are reminiscent of the natural analogue, and promote localized hydroxyapatite formation to permit integration with the subchondral bone.

**KEYWORDS:** cartilage, scaffold, tissue-engineering, multi-layer, nanocomposite

Download English Version:

<https://daneshyari.com/en/article/6485350>

Download Persian Version:

<https://daneshyari.com/article/6485350>

[Daneshyari.com](https://daneshyari.com)