## Accepted Manuscript

Title: The Carotenoid Bixin: Optical Studies of Aggregation in

Polar/Water Solvents

Authors: Sefadzi Tay-Agbozo, Shane Street, Lowell D. Kispert

PII: \$1010-6030(18)30453-2

DOI: https://doi.org/10.1016/j.jphotochem.2018.05.008


Reference: JPC 11278

To appear in: Journal of Photochemistry and Photobiology A: Chemistry

Received date: 7-4-2018 Revised date: 4-5-2018 Accepted date: 7-5-2018

Please cite this article as: Tay-Agbozo S, Street S, Kispert LD, The Carotenoid Bixin: Optical Studies of Aggregation in Polar/Water Solvents, *Journal of Photochemistry and Photobiology, A: Chemistry* (2018), https://doi.org/10.1016/j.jphotochem.2018.05.008

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



### ACCEPTED MANUSCRIPT


# The Carotenoid Bixin: Optical Studies of Aggregation in Polar/Water Solvents

Sefadzi Tay-Agbozo, Shane Street, Lowell D. Kispert\*

Department of Chemistry, BOX 870336, University of Alabama, Tuscaloosa, AL, 35487, United States

\*lkispert@ua.edu

#### **Graphical Abstract**



#### Highlights:

- Bixin exhibits a 4 nm red shift in a 50% water-methanol solution, suggesting possible J-aggregation.
- The lack of H aggregation can be traced to the fact that the ground state geometry for bixin is *cis* and not *trans*, resulting in an unfavorable geometry for H aggregate formation.

#### Download English Version:

# https://daneshyari.com/en/article/6492468

Download Persian Version:

https://daneshyari.com/article/6492468

Daneshyari.com