


Accepted Manuscript

Antiphlogistic effect by zeolite as determined by a murine inflammation model

Javiera Cervini-Silva, Antonio Nieto-Camacho, Stephan Kaufhold, Kristian Ufer, Eduardo Palacios, Ascención Montoya, Wilfried Dathe



PII: S1387-1811(16)30069-5

DOI: [10.1016/j.micromeso.2016.03.043](https://doi.org/10.1016/j.micromeso.2016.03.043)

Reference: MICMAT 7657

To appear in: *Microporous and Mesoporous Materials*


Received Date: 20 February 2016

Revised Date: 24 March 2016

Accepted Date: 29 March 2016

Please cite this article as: J. Cervini-Silva, A. Nieto-Camacho, S. Kaufhold, K. Ufer, E. Palacios, A. Montoya, W. Dathe, Antiphlogistic effect by zeolite as determined by a murine inflammation model, *Microporous and Mesoporous Materials* (2016), doi: 10.1016/j.micromeso.2016.03.043.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



Zeolite (65% clinoptilolite, 30% mordenite, and 5% smectite) shows antiphlogistic properties shortly after exposure. Representative histological sections from mice ears exposed to TPA (4 h) and stained with hematoxylin-eosin treated with 1 mg ear^{-1} zeolite for 24 h. (20 X Scale = $100 \mu\text{m}$). Arrows indicate the infiltrated leukocytes.

Download English Version:

<https://daneshyari.com/en/article/6532772>

Download Persian Version:

<https://daneshyari.com/article/6532772>

[Daneshyari.com](https://daneshyari.com)