



Original Article

Effects of phosphorus addition on nitrogen cycle and fluxes of N₂O and CH₄ in tropical tree plantation soils in Thailand



Taiki Mori,^{a, c, *} Chongrak Wachrinrat,^b Duriya Staporn,^b Ponthep Meunpong,^b Warawich Suebsai,^b Kazuki Matsubara,^c Khitja Boonsri,^b Warisa Lumban,^b Manassawee Kuawong,^b Thanida Phukdee,^b Juruwan Srifai,^b Kannika Boonman^b

^a Forest Ecology Laboratory, Graduate School of Agriculture, Kyoto University, Kitashirakawa Oiwake-cho, Sakyo-ku, Kyoto 606-8502, Japan

^b Faculty of Forestry, Kasetsart University, 50 Ngamwongwan Road, Chatuchak, Bangkok 10900, Thailand

^c Key Laboratory of Vegetation Restoration and Management of Degraded Ecosystems, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou, Guangdong 510650, China

ARTICLE INFO

Article history:

Received 3 August 2015

Accepted 15 March 2016

Available online 10 June 2017

Keywords:

Soil respiration

Nitrous oxide

Methane

Tropics

Phosphorus

Tree plantation

ABSTRACT

An incubation experiment was conducted to test the effects of phosphorus (P) addition on nitrous oxide (N₂O) emissions and methane (CH₄) uptakes, using tropical tree plantation soils in Thailand. Soil samples were taken from five forest stands—*Acacia auriculiformis*, *Acacia mangium*, *Eucalyptus camaldulensis*, *Hopea odorata*, and *Xylia xylocarpa*—and incubated at 80% water holding capacity. P addition stimulated N₂O emissions only in *Xylia xylocarpa* soils. Since P addition tended to increase net ammonification rates in *Xylia xylocarpa* soils, the stimulated N₂O emissions were suggested to be due to the stimulated nitrogen (N) cycle by P addition and the higher N supply for nitrification and denitrification. In other soils, P addition had no effects on N₂O emissions or soil N properties, except that P addition tended to increase the soil microbial biomass N in *Acacia auriculiformis* soils. No effects of P addition were observed on CH₄ uptakes in any soil. It is suggested that P addition on N₂O and CH₄ fluxes at the study site were not significant, at least under laboratory conditions.

Copyright © 2017, Kasetsart University. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Since the primary source of phosphorus (P) for terrestrial ecosystems is rock weathering, P has been considered to be the ultimate limiting soil nutrient in terrestrial ecosystems, and ecosystems with old soils can become depleted in P (Walker and Syers, 1976). Thus, highly-weathered lowland tropical forest soils have been considered to have low concentrations of biologically available P (Vitousek and Farrington, 1997; Elser et al., 2007; Vitousek et al., 2010), and it is widely believed that P limits the ecosystem process in tropical forests. Several studies have reported that P availability also limits soil microbial activities in tropical soils (Cleveland et al., 2002; Ilstedt et al., 2003, 2006; Cleveland and Townsend, 2006; Mori et al., 2010; Liu et al., 2012). Thus, changes in P availability may change soil microbial activities, metabolisms

and community composition, which could be associated with altered nitrous oxide (N₂O) (Mori et al., 2013a, 2013b, 2014) and methane (CH₄) (Mori et al., 2013c) exchange between terrestrial ecosystems and the atmosphere (Mori et al., 2013d).

Soil of tropical forests is an important source of N₂O, which is a by-product or intermediate product of microbial nitrification and denitrification, respectively (Wrage et al., 2001). Tropical forest soils have also the potential to function as substantial sinks of CH₄ (Potter et al., 1996). CH₄ fluxes are the net result of the simultaneous microbial oxidation of CH₄ by methanotrophs in predominantly aerobic soil zones and microbial production of CH₄ by methanogenesis in predominantly anaerobic soil zones (Le Mer and Roger, 2001).

Recently several studies have reported that P addition reduced N₂O emissions through stimulated plant N uptake (Mori et al., 2013d; Baral et al., 2014; Zhang et al., 2014). Mori et al. (2014) experimentally confirmed this suggestion by showing that P addition reduced N₂O emissions from an *Acacia mangium* plantation, but did not if plant roots were excluded by the trenching method. The authors' understanding is still lacking on the effects of P

* Corresponding author. Forest Ecology Laboratory, Graduate School of Agriculture, Kyoto University, Kitashirakawa Oiwake-cho, Sakyo-ku, Kyoto 606-8502, Japan.

E-mail address: taikimori7@gmail.com (T. Mori).

addition on microbial activity, without plant interaction, and accompanying N₂O emissions. Hall and Matson (1999) suggested that a P shortage in tropical soils limits microbial N immobilization, resulting in an increase in N resources for nitrification or denitrification or both. Their report implied that P addition alleviates a P shortage and increases microbial N immobilization, resulting in a reduction in N resources for nitrification or denitrification or both and a reduction in N₂O emissions. However, previous papers have reported completely opposite results, with higher N₂O emission rates in P-added soils than in the non-added control (Mori et al., 2010, 2013a). Alleviation of P-limitation on nitrifying or denitrifying bacteria or both, stimulated whole N cycling in the soil, and improved reductive conditions with stimulated denitrifying activity being the suggested mechanism. However only one soil was used (soil taken from an *Acacia mangium* plantation in South Sumatra, Indonesia), and there are still very few reports testing the effects of P addition on N₂O emissions.

It has been shown that P addition increased (Aerts and Toet, 1997), decreased (Conrad et al., 2000), or had no effects (Lund et al., 2009; Keller et al., 2005) on CH₄ fluxes. It was also reported that P application lowered CH₄ fluxes indirectly by stimulating root water uptake (Zhang et al., 2011). Mori et al. (2013b) reported that P addition had no effects on CH₄ fluxes in an *Acacia mangium* plantation, but significantly increased CH₄ uptake when there were no plant interactions. Thus, there are several reports examining the effects of P addition on CH₄ fluxes, but few studies have reported the effects in tropical forest soils, especially tropical tree plantation soils.

The current study conducted another incubation experiment using soils taken from five different tropical, monoculture, tree plantations. The effects of P addition on N₂O and CH₄ fluxes were examined.

Materials and methods

Soil sampling

Soil samples were collected at plantation sites inside the Sakaerat Environment Research Station (SERS) in Nakhon Ratchasima, Thailand (14°30'N, 101°55'E). The climate of the region is classified as tropical savannah (Yamashita et al., 2010). The mean annual temperature was 25.5 °C and annual precipitation was 1407 mm for 2000–2008 (Yamashita et al., 2011). The main soil type is Acrisols (Yamashita et al., 2011). Five forest plantation stands were chosen—*Acacia auriculiformis* (Leguminosae), *Acacia mangium* (Leguminosae), *Eucalyptus camaldulensis* (Myrtaceae), *Hopea odorata* (Dipterocarpaceae), and *Xylia xylocarpa* (Leguminosae). At the beginning of the experiment, each stand was aged 9 yr. Soil samples (0–5 cm depth) were collected from six randomly selected points in each forest stand using 100 mL soil cores. Litter layers were removed before soil sampling. After collection, each soil

sample was sieved through a 2 mm sieve. The general physicochemical characteristics of the 0–5 cm soil samples are shown in Table 1. The particle size distribution was determined using the pipette method (Gee and Bauder, 1986). The pH (H₂O) was determined for 1:2.5 water suspensions using a glass electrode (Horiba; Kyoto, Japan). The total C and total N contents were determined using an NC analyzer (JM 1000CN; J-Science Lab Co. Ltd.; Kyoto Japan). The available P contents were determined using the Bray-1 method (Kuo, 1996).

Incubation

Fresh soil samples of 30 g were placed in 223 mL wide-mouth jars for gas sampling, and 5 g samples were placed in 50 mL bottles for chemical analyses (each forest stand had six replications). For each analysis, two subsamples were prepared—one for P addition and the other for the non-P-added control. P was added as KH₂PO₄ solution (100 µg P/g soil, dissolved in distilled water). Non-P-added controls were prepared without P addition in the same manner. After the soil water condition was adjusted to 80% water holding capacity, the samples were incubated at 25 °C in the dark for 48 h. The wide-mouth jars were closed with butyl rubber stoppers equipped with sampling ports, and gas samples were taken at 0 h and 48 h after the closure of the stoppers. Gas concentrations were analyzed using a gas chromatograph (GC-14B; Shimadzu; Kyoto, Japan) equipped with an electron capture detector for N₂O and a flame ionization detector for CH₄. The gas fluxes were calculated from the differences between the gas concentrations at 0 h and 48 h.

Inorganic N and dissolved N were extracted by shaking 5 g of fresh soil with 25 mL of 0.5 M K₂SO₄ for 30 min. The NH₄ and NO₃⁻ contents were determined using a flow-injection analyzer (AQLA-700-NO; Aqualab; Tokyo, Japan). The dissolved N concentration was analyzed using a total organic carbon analyzer with a total organic nitrogen measurement unit (TOC-V_E/TNM-1; Shimadzu; Kyoto, Japan). The soil microbial biomass N was determined using a chloroform fumigation extraction method (Jenkinson et al., 2004). Fresh soil samples of 5 g were exposed to CHCl₃ vapor for 24 h in a vacuum desiccator at 25 °C. After the residual CHCl₃ had been removed, the fumigated soils were shaken with 50 mL of 0.5 M K₂SO₄ extractant for 30 min and the dissolved N was extracted. The soil microbial biomass element contents were calculated from the differences of the dissolved N contents between the fumigated and unfumigated samples using a conversion factor of 0.45 (Jenkinson et al., 2004). The soil pH (H₂O) was measured at the end of the incubation period.

Statistical analysis

Statistical analyses were performed using the Excel software (version 2013; Microsoft Corp; Redmond, WA, USA). The level of

Table 1
Physicochemical properties of soils at different plantation sites (data from Mori et al., 2016).

Site ^a	pH (H ₂ O)	Total C ^b (mg C/g soil)	Total N ^b (mg N/g soil)	Available P ^c (µg P/g soil)	Clay ^d (%)	Silt ^d (%)	Sand ^d (%)
AA	4.9	21.6	2.1	12.1	7.3	21.3	71.5
AM	5.1	12.0	1.4	7.0	3.8	16.0	80.2
EC	5.3	13.5	1.4	12.2	6.4	15.6	78.0
HO	5.0	10.7	1.5	9.9	5.1	17.3	77.5
XX	5.2	11.7	1.4	10.9	3.1	16.5	80.4

^a AA, *Acacia auriculiformis*; AM, *Acacia mangium*; EC, *Eucalyptus camaldulensis*; and HO, *Hopea odorata*; XX, *Xylia xylocarpa*.

^b Average of three replications.

^c Average of six replications.

^d Average of two replications.

Download English Version:

<https://daneshyari.com/en/article/6538099>

Download Persian Version:

<https://daneshyari.com/article/6538099>

[Daneshyari.com](https://daneshyari.com)