## Accepted Manuscript

Title: Establishment of a prediction method for the mid-facial region of unknown human Mongoloid skeletal remains

Authors: Hajime Utsuno, Toru Kageyama, Keiicchi Uchida, Namiko Ishii, Saki Minegishi, Koichi Uemura, Koichi Sakurada



PII:	S0379-0738(18)30219-6
DOI:	https://doi.org/10.1016/j.forsciint.2018.04.052
Reference:	FSI 9291
To appear in:	FSI
Received date:	12-11-2017
Revised date:	19-4-2018
Accepted date:	26-4-2018

Please cite this article as: Hajime Utsuno, Toru Kageyama, Keiicchi Uchida, Namiko Ishii, Saki Minegishi, Koichi Uemura, Koichi Sakurada, Establishment of a prediction method for the mid-facial region of unknown human Mongoloid skeletal remains, Forensic Science International https://doi.org/10.1016/j.forsciint.2018.04.052

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## ACCEPTED MANUSCRIPT

## Establishment of a prediction method for the mid-facial region of unknown human Mongoloid skeletal remains

Hajime Utsuno<sup>a</sup>, Toru Kageyama<sup>b</sup>, Keiicchi Uchida <sup>c</sup>, Namiko Ishii <sup>a</sup>, Saki Minegishi <sup>a</sup>, Koichi Uemura<sup>d</sup>, Koichi Sakurada <sup>a</sup>

<sup>a</sup> Department of Forensic Dentistry, Graduate School of Medical and Dental Sciences, Tokyo Medical and Dental University, 1-5-45 Yushima, Bunkyou-ku, Tokyo 113-8510, Japan. Ph:+81 3 5803 4164; Email: hazimeu.fde@tmd.ac.jp

<sup>b</sup> Department of Orthodontics, Matsumoto Dental University, School of Dentistry, 1780 GobaraHirooka, Shiojiri, Nagano 399-0781, Japan. Ph: +81 263 52 2100; Email: toru@po.mdu.ac.jp

<sup>c</sup> Department of Oral Radiology, Matsumoto Dental University, School of Dentistry, 1780 GobaraHirooka, Shiojiri, Nagano 399-0781, Japan. Ph: +81 263 52 2100; Email: keiboba@po.mdu.ac.jp

<sup>d</sup> Department of Legal Medicine, Graduate School of Medical and Dental Sciences, Tokyo Medical and Dental University, 1-5-45 Yushima, Bunkyou-ku, Tokyo 113-8510, Japan. Ph:+81 3 5803 5199; Email: kuemura.legm@tmd.ac.jp

Corresponding author: Hajime Utsuno, Department of Forensic Dentistry, Graduate School of Medical and Dental Sciences, Tokyo Medical and Dental University. 1-5-45 Yushima Bunkyou-ku, Tokyo 113-8510, Japan Ph:+81 3 5803 4164; Email: hazimeu.fde@tmd.ac.jp

## Highlights

- Sex differences were observed in the measurements of mid-lower part of the maxilla.
- Proportional differences in the skeletal classes were not observed in the same sex.
- The calculations to predict the *Sn* were generated from measurements.
- The prediction demonstrated  $\leq 5$  mm error with high correlations.

Download English Version:

https://daneshyari.com/en/article/6551015

Download Persian Version:

https://daneshyari.com/article/6551015

Daneshyari.com