

Accepted Manuscript

Title: Three-dimensionally derived interlandmark distances for sex estimation in intact and fragmentary crania

Authors: Candice Small, Lynne Schepartz, Jason Hemingway, Desiré Brits



PII: S0379-0738(18)30064-1
DOI: <https://doi.org/10.1016/j.forsciint.2018.02.012>
Reference: FSI 9169

To appear in: *FSI*

Received date: 23-11-2017
Revised date: 29-1-2018
Accepted date: 14-2-2018

Please cite this article as: Candice Small, Lynne Schepartz, Jason Hemingway, Desiré Brits, Three-dimensionally derived interlandmark distances for sex estimation in intact and fragmentary crania, Forensic Science International <https://doi.org/10.1016/j.forsciint.2018.02.012>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Three-dimensionally derived interlandmark distances for sex estimation in intact and fragmentary crania

Candice Small*, Lynne Schepartz, Jason Hemingway, Desiré Brits

School of Anatomical Sciences, Faculty of Health Sciences, University of the Witwatersrand,
Human Variation and Identification Research Unit (HVIRU)

Johannesburg 2193, Republic of South Africa

Candice.small@wits.ac.za

Lynne.schepartz@wits.ac.za

Jason.hemingway@wits.ac.za

Desire.brits@wits.ac.za

*Corresponding author:

Candice Small

University of the Witwatersrand,

Faculty of Health Sciences, School of Anatomical Sciences

7 York Road

Parktown

Johannesburg

2193

Republic of South Africa

Office: +2711 717 2077

E-mail: Candice.small@wits.ac.za

Highlights:

- 990 interlandmark distances were mathematically derived using Pythagorean geometry
- Distances include non-traditional measures used to derive discriminant functions
- The functions represent five regions and are applicable to fragmentary remains
- Functions for fragmentary remains have accuracies ranging from 74.0% to 83.2%

Download English Version:

<https://daneshyari.com/en/article/6551047>

Download Persian Version:

<https://daneshyari.com/article/6551047>

[Daneshyari.com](https://daneshyari.com)