

Accepted Manuscript

Title: Amphetamines and Cannabinoids testing in hair:
evaluation of results from a two-year period

Author: María José Burgueño Amaya Alonso Sergio Sánchez



PII: S0379-0738(16)00011-6
DOI: <http://dx.doi.org/doi:10.1016/j.forsciint.2016.01.003>
Reference: FSI 8271

To appear in: *FSI*

Received date: 26-9-2015
Revised date: 3-1-2016
Accepted date: 4-1-2016

Please cite this article as: M.J. Burgueño, A. Alonso, S. Sánchez, Amphetamines and Cannabinoids testing in hair: evaluation of results from a two-year period., *Forensic Science International* (2016), <http://dx.doi.org/10.1016/j.forsciint.2016.01.003>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Amphetamines and Cannabinoids testing in hair: evaluation of results from a two-year period.

María José Burqueño^{*}, Amaya Alonso, Sergio Sánchez
National Institute of Toxicology and Forensic Sciences, Drug of Abuse Department. C/ José Echegaray, 4, 28232 - Madrid, Spain

mjose.burgueno@justicia.es

Highlights

- Amphetamines and cannabinoids tests performed on 3,000 hair samples were studied.
- While THC and MDMA use were conditioned by age and gender, AP use was not.
- Positivity rate for AP was significant higher in dyed vs. natural hair.
- Hair length between 3.0 and 5.5 cm should be agreed for THC testing.

Download English Version:

<https://daneshyari.com/en/article/6551738>

Download Persian Version:

<https://daneshyari.com/article/6551738>

[Daneshyari.com](https://daneshyari.com)