

Accepted Manuscript

Title: Defining mtDNA origins and population stratification in Rio de Janeiro

Authors: Filipa Simão, Ana Paula Ferreira, Elizeu Fagundes de Carvalho, Walther Parson, Leonor Gusmão



PII: S1872-4973(18)30072-3
DOI: <https://doi.org/10.1016/j.fsigen.2018.02.003>
Reference: FSIGEN 1844

To appear in: *Forensic Science International: Genetics*

Received date: 5-12-2017
Revised date: 29-1-2018
Accepted date: 2-2-2018

Please cite this article as: Filipa Simão, Ana Paula Ferreira, Elizeu Fagundes de Carvalho, Walther Parson, Leonor Gusmão, Defining mtDNA origins and population stratification in Rio de Janeiro, *Forensic Science International: Genetics* <https://doi.org/10.1016/j.fsigen.2018.02.003>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1 **Defining mtDNA origins and population stratification in Rio de Janeiro**

2 Filipa Simão¹, Ana Paula Ferreira¹, Elizeu Fagundes de Carvalho¹, Walther Parson^{2,3,*},
3 Leonor Gusmão¹

4 ¹DNA Diagnostic Laboratory (LDD), State University of Rio de Janeiro (UERJ), Rio de
5 Janeiro, Brazil

6 ²Institute of Legal Medicine, Medical University of Innsbruck, Innsbruck, Austria

7 ³Forensic Science Program, The Pennsylvania State University, University Park, PA,
8 USA.

9

10 * Corresponding author at: Medical University of Innsbruck, Muellerstr. 44, Innsbruck
11 6020, Austria. E-mail address: walther.parson@i-med.ac.at (W. Parson).

12

13

14 **Highlights**

15 Population data (n = 205) are reported for the mtDNA control region in samples from Rio
16 de Janeiro (Brazil).

17 High haplotype diversity is present in Rio de Janeiro as a result of significant African,
18 European and Native American contributions.

19 The main maternal gene pool can be traced to Africa, arriving mainly from the Atlantic
20 Coast region.

21 A comparison between mtDNA and autosomal data revealed population substructure,
22 which has to be considered in forensic analysis.

23

24

25

26

27

28 **Introduction**

29 Genetic profiling is commonly used to assist criminal investigations. The study of genetic
30 markers in forensic casework requires the implementation of DNA databases
31 representative of the composition of reference populations. The accuracy of these
32 databases is crucial to correctly estimate the frequency of a genetic profile, which is the
33 only way to attribute statistical weight to the evidence. Additionally, to avoid erroneous
34 estimates of the probability of a certain observation, it is also important to evaluate if

Download English Version:

<https://daneshyari.com/en/article/6553265>

Download Persian Version:

<https://daneshyari.com/article/6553265>

[Daneshyari.com](https://daneshyari.com)