



ELSEVIER

Contents lists available at ScienceDirect

Data in Brief

journal homepage: www.elsevier.com/locate/dib

Data article

Dataset of plasma and aqueous humor cytokine profiles in patients with exudative age related macular degeneration and polypoidal choroidal vasculopathy

Praveen Kumar Balne^{a,b}, Rupesh Agrawal^{a,b,*}, Veonice Bijin AU^c,
Bennett Lee^d, Arkasubhra Ghosh^e, Swaminathan Sethu^e,
Mukesh Agrawal^f, Raja Narayanan^g, John Connolly^c

^a National Healthcare Group Eye Institute, Tan Tock Seng Hospital, 11 Jalan Tan Tock Seng, Singapore 308433, Singapore

^b Singapore Eye Research Institute, Singapore 169856, Singapore

^c Institute of Molecular and Cell Biology, Singapore 138673, Singapore

^d Singapore Immunology Network, Singapore 138648, Singapore

^e GROW Research Laboratory, Narayana Nethralaya Foundation, 560099, India

^f Vimta Labs Limited, Hyderabad, Telangana 500051, India

^g L. V. Prasad Eye Institute, Hyderabad, Telangana 500034, India

ARTICLE INFO

Article history:

Received 16 January 2018

Received in revised form

15 May 2018

Accepted 18 May 2018

Available online 24 May 2018

Keywords:

Exudative age related macular degeneration

Choroidal neovascularization, Polypoidal

choroidal vasculopathy

Cytokine profiles

Aqueous humor

Plasma

ABSTRACT

In this report the data was obtained from a prospective case-control study with a sample size of sixteen patients with exudative age related macular degeneration (AMD) due to choroidal neovascularization (CNV) and eighteen patients with polypoidal choroidal vasculopathy (PCV) and fifty controls (cataract patients without any other ocular diseases). Luminex bead based multiplex assay with a panel of 41 analytes was used to study the cytokine levels in plasma and aqueous humor.

© 2018 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

* Corresponding author at: National Healthcare Group Eye Institute, Tan Tock Seng Hospital, Singapore 308433, Singapore.

E-mail addresses: praveenbalne@gmail.com (rupesh_agrawal@ttsh.com.sg (R. Agrawal), bjau@imcb.a-star.edu.sg (V.B. AU), bennett_lee@immunol.a-star.edu.sg (B. Lee), arkasubhra@narayananeethralaya.com (A. Ghosh), swaminathansethu@narayananeethralaya.com (S. Sethu), mukesh.agrawal@vimta.com (M. Agrawal), narayanan@lvpei.org (R. Narayanan), jeconnolly@imcb.a-star.edu.sg (J. Connolly).

<https://doi.org/10.1016/j.dib.2018.05.085>

2352-3409/© 2018 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Specifications Table

Subject area	<i>Biology</i>
More specific subject area	<i>Plasma and aqueous humor cytokine levels in exudative age related macular degeneration (AMD)</i>
Type of data	<i>Table</i>
How data was acquired	<i>Bead based multiplex assay using the Luminex xMAP technology</i>
Data format	<i>Raw</i>
Experimental factors	<i>Plasma and aqueous humor samples were collected from study subjects and each sample was analyzed for 41 cytokine levels</i>
Experimental features	<i>Milliplex[®] MAP human cytokine / chemokine magnetic bead panel -1 kit (Millipore, USA) with FlexMAP 3D (Luminex[®]) platform was used for cytokine profiling in each sample</i>
Data source location	<i>Singapore</i>
Data accessibility	<i>Data is with this article</i>

Value of the data

- Here we report the plasma and aqueous humor cytokine levels in patients with exudative AMD, PCV and controls (cataract patients without any other ocular diseases).
- In exudative AMD patients, aqueous humor cytokines GRO, MDC and MIP-1 α were significantly higher than controls ($p < 0.04$).
- In PCV patients, aqueous humor cytokines GRO, MDC, MIP-1 α , MIP-1 β , IL-8, IP-10 and MCP levels were significantly higher than controls ($p < 0.03$).
- No significant differences were seen in plasma cytokine levels between cases and controls.
- None of the plasma and aqueous humor cytokines showed significant differences between exudative AMD and PCV.
- This data will help researchers to understand the differences in cytokine expression levels between patients with Exudative AMD, PCV and controls.
- This data will also help researchers to understand the differences between plasma and aqueous humor in patients with Exudative AMD, PCV and healthy controls.

1. Data

The data presented in this article was obtained from a prospective case-control study and shows the concentration of 41 cytokines in plasma and aqueous humor from patients with exudative age related macular degeneration (AMD), polypoidal choroidal vasculopathy (PCV) and healthy controls (Table 1).

2. Experimental design, materials and methods

In a prospective case-control study, sixteen patients who were diagnosed with exudative age related macular degeneration (AMD) due to choroidal neovascularization (CNV) and eighteen patients with polypoidal choroidal vasculopathy (PCV) and fifty age and gender matched cataract patients without any other ocular complications (controls) were recruited. The study was approved by the Institutional review board of Tan Tock Seng Hospital, Singapore and adhered to the tenets of the Declaration of Helsinki. Prior informed consent was obtained from all the subjects. Five milliliters of peripheral venous blood collected aseptically from 14 AMD patients, 18 PCV patients and fifty controls and approximately 200 μ L of aqueous humor collected from 16 AMD patients, 16 PCV patients and 41 controls using standard protocols were used for cytokine profiling [1]. Clinical samples were

Download English Version:

<https://daneshyari.com/en/article/6596678>

Download Persian Version:

<https://daneshyari.com/article/6596678>

[Daneshyari.com](https://daneshyari.com)