



ELSEVIER

Contents lists available at ScienceDirect

Data in Brief

journal homepage: www.elsevier.com/locate/dib

Data Article

Q1 Data on phosphorous concentration of rivers feeding into Taham dam in Zanjan, Iran

Mohammadreza Masoudinejad ^a, Mansour Ghaderpoori ^{b,c,*},
 Ahmad Zarei ^d, Jamal Nasehifar ^e, Alireza Malekzadeh ^{e,*},
 Jalil Nasiri ^e, Afshin Ghaderpoury ^f

Q2 ^a Safety Promotion and Injury Prevention Research Center (SPIPRC), Shahid Beheshti University of Medical Sciences, Department of Environmental Health Engineering, Tehran, Iran

^b Nutritional Health Research Center, Lorestan University of Medical Sciences, Khorramabad, Iran

^c Department of Environmental Health Engineering, School of Health and Nutrition, Lorestan University of Medical Sciences, Khorramabad, Iran

^d Department of Environmental Health Engineering, School of Public Health, Gonabad University of Medical Sciences, Gonabad, Iran

^e Department of Environmental Health Engineering, School of Public Health, Zanjan University of Medical Sciences, Zanjan, Iran

^f Shahid Beheshti University of Medical Sciences, Tehran, Iran

ARTICLE INFO

Article history:

Received 5 December 2017

Received in revised form

19 January 2018

Accepted 22 January 2018

Keywords:

Eutrophication

Phosphorous

Rivers

Sewage

Seasons

ABSTRACT

Due to the great possibility of water contamination of many rivers by human activities in Iran, the study of water quality is crucial for water resource protection and human health. High level of phosphorous is the main reason for eutrophication of freshwater systems. The main aim of this study was to investigate the concentration of phosphorus in the rivers feeding into Taham dam in Zanjan, using GIS software. 40 sampling stations were selected along Taham and Ghalharod Rivers with respect to sewage discharge points and feeding characteristics of water entering to Taham dam. In total, 160 water samples were taken from rivers with regard to precipitation season in two different periods from winter 2014 to spring 2015. The obtained data were analyzed using SPSS and ArcView GIS. The findings showed that 15% of the studied stations had phosphorous levels higher than acceptable levels set by EPA. The highest levels of phosphorous contamination were observed in stations No. 145, 154,

* Corresponding author at: Nutritional Health Research Center, Lorestan University of Medical Sciences, Khorramabad, Iran.

** Corresponding author.

E-mail addresses: massoudi2010@yahoo.com (M. Masoudinejad), mghaderpoori@gmail.com (M. Ghaderpoori), a.zarei.tums@gmail.com (A. Zarei), jamal_46@yahoo.com (J. Nasehifar), malekzadea@yahoo.com (A. Malekzadeh), jalil.nassiri@gmail.com (J. Nasiri), ghaderpoury_a@yahoo.com (A. Ghaderpoury).

<https://doi.org/10.1016/j.dib.2018.01.068>

2352-3409/© 2018 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

155, 161, 166 and 168. The elevated concentrations of phosphorous in the rivers can be responsible for the eutrophication of Taham dam reservoir.

© 2018 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Specifications Table

Subject area	<i>Chemistry, biology</i>
More specific subject area	<i>Water monitoring and quality</i>
Type of data	<i>Figure</i>
How data was acquired	<i>The levels of phosphorus in water samples were determined using a DR-5000 UV-vis spectrophotometer at wavelength 470nm (HACH, Germany).</i>
Data format	<i>Raw, analyzed,</i>
Experimental factors	<i>After selection of sampling stations, the coordinates of stations were recorded using a GPS system (Garmin GPSMAP 76CSX) and subsequently a map of the GPS points were created. The final transfer of data was done by Arc GIS.</i>
Experimental features	<i>Measuring the concentration of phosphorus in the feeding rivers of Taham dam which provides the water demands of Zanjan city</i>
Data source location	<i>Zanjan city, Zanjan province, west of Tehran, Iran</i>
Data accessibility	<i>Data are included in this study and supplemented excel file</i>

Value of the data

- Contamination of water in rivers and streams can be considered as an index of environment pollution by human activities.
- Because rivers are the only water resources which pass long distances among cities, towns, industrial and agricultural lands and therefore, there is a great possibility of their contamination by various contaminants from anthropogenic activities.
- High level of phosphorous is the main reason for eutrophication of freshwater systems (9). Eutrophication can be occurred due to the existence of nitrogen and phosphorous compounds in water which changes the water ecosystem and causes many problems in water bodies.
- High levels of phosphorous in water are not poisonous directly, and there are no reports in the literature regarding direct poisonous or harmful effects associated with elevated concentrations of phosphorous in the human body but in general decrease the concentrations of calcium and magnesium in human blood.

1. Data

Zanjan city is located in 330 km from west of Tehran with a total population of 411,001 inhabitants. The study area lies in Longitudes 48°29'E and Latitudes 36°40'N with an elevation of 1650 m above sea level. The location of the Zanjan city shows in Fig. 1. Taham dam is located in 15 km in the north-west of Zanjan city and 8 km downward of Taham town. This dam provides distribution water of Zanjan city.

Download English Version:

<https://daneshyari.com/en/article/6597221>

Download Persian Version:

<https://daneshyari.com/article/6597221>

[Daneshyari.com](https://daneshyari.com)