



ELSEVIER

Contents lists available at ScienceDirect

Data in Brief

journal homepage: www.elsevier.com/locate/dib

Data Article

Adsorption of diclofenac onto different biochar microparticles: Dataset – Characterization and dosage of biochar

Linson Lonappan^a, Tarek Rouissi^a, Satinder Kaur Brar^{a,*}, Mausam Verma^b, Rao Y. Surampalli^c^a INRS-ETE, Université du Québec, 490, Rue de la Couronne, Québec, Canada G1K 9A9^b CO2 Solutions Inc., 2300, Rue Jean-Perrin, Québec, Canada G2C 1T9^c Department of Civil Engineering, University of Nebraska-Lincoln, N104 SEC P.O. Box 886105, Lincoln, NE 68588-6105, United States

ARTICLE INFO

Article history:

Received 11 October 2017

Accepted 17 October 2017

Keywords:

Adsorption

Diclofenac

Biochar

Characterization

ABSTRACT

Due to its wide occurrence in water resources and toxicity, pharmaceuticals and personal care products are becoming an emerging concern throughout the world. Application of residual/waste materials for water remediation can be a good strategy in waste management as well as in waste valorization. Herein, this dataset provides information on biochar application for the removal of emerging contaminant, diclofenac from water matrices. The data presented here is an extension of the research article explaining the mechanisms of adsorption diclofenac on biochars (Lonappan et al., 2017 [1]). This data article provides general information on the surface features of pine wood and pig manure biochar with the help of SEM and FTIR data. This dataset also provides information on XRD profiles of pine wood and pig manure biochars. In addition, different amounts of biochars were used to study the removal of a fixed concentration of diclofenac and the data is provided with this data set.

© 2017 Published by Elsevier Inc. This is an open access article under the CC BY license

(<http://creativecommons.org/licenses/by/4.0/>).

DOI of original article: <http://dx.doi.org/10.1016/j.biortech.2017.10.039>

* Corresponding author.

E-mail address: satinder.brar@ete.inrs.ca (S.K. Brar).

<http://dx.doi.org/10.1016/j.dib.2017.10.041>

2352-3409/© 2017 Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Specifications Table

Subject area	Chemistry/Chemical engineering
More specific subject area	Adsorption, Surface Chemistry, Environmental Engineering
Type of data	Table, image (XRD, SEM)), text file, figure (FTIR)
How data was acquired	SEM: Zeiss Evo@50 Smart SEM FTIR: Perkin Elmer, Spectrum RXI, FT-IR instrument fitted with lithium tantalate (LiTaO ₃) detector XRD: Panalytical Empyrean XRD with monochromatized CuK alfa radiation (1.5418Å). LDTD-MS/MS: Concentrations of diclofenac was measured using LDTD-APCI (atmospheric pressure chemical ionization) source (LDTD T-960, Phytronix Technologies, Quebec, Canada) mounted on a TSQ Quantum access triple quadruple mass spectrometer (Thermo Scientific, Mississauga, Ontario, Canada)
Data format	Pre-processed and analyzed
Experimental factors	Biochar samples (from pinewood and pig manure) were grounded to obtain microparticles and the data here is given is for characterization of biochar. Moreover, data for dosage effect of biochar on adsorption for diclofenac is given.
Experimental features	Characterization data of biochar microparticles obtained from SEM, XRD, and FTIR are given. Adsorption studies were carried out for the removal of diclofenac using biochar microparticles. Various biochar dosages ranging from 1 g L ⁻¹ to 20 g L ⁻¹ were tested.
Data source location	Bioprocessing and NanoEnzyme Formulation Facility (BANEFF), INRS-ETE, Université du Québec, 490, Rue de la Couronne, Québec, Canada G1K 9A9
Data accessibility	Data presented in this article
Related research article	The associated research article related to this data set is [1]

Value of the data

- Characterization data for biochar derived from two different feedstock (pine wood and pig manure) are given.
- Dataset provides an insight to the surface features of biochar.
- Dataset gives information on the adsorption capacity of biochar for emerging contaminant diclofenac.
- Dataset would be useful to identify the dosage effect of biochar on the adsorption of diclofenac.

1. Data

The dataset comprises characterization as well as experimental data. Fig. 1 presents the scanning electron micrographs (SEM) of pine wood and pig manure biochar microparticles. Fig. 2. presents Fourier-transform infrared spectroscopy (FTIR) images of biochar microparticles. Fig. 3 presents X-ray Diffraction (XRD) images of biochar microparticles. Table 1 shows the effect of adsorbent dosage on the removal of diclofenac and removal efficiency.

Download English Version:

<https://daneshyari.com/en/article/6597253>

Download Persian Version:

<https://daneshyari.com/article/6597253>

[Daneshyari.com](https://daneshyari.com)