

Technical Note

Duhamel integral form for the interface heat flux between bubble and liquid

Nail S. Khabeev *

Department of Mathematics, University of Bahrain, P.O. Box 32038, Bahrain

Received 27 March 2006; received in revised form 4 June 2007

Available online 24 September 2007

Abstract

The solution of heat equation inside oscillating gas bubble with moving boundary was obtained by Fourier's method. The integral formula for interface heat flux, containing theta-function in the integrand was derived. The kernel of the integral is represented by a series of exponential functions, and a simple analytic approximation obtained earlier is used for it with high accuracy. The asymptotic expression for the interface heat flux in the Duhamel integral form with rooted kernel was derived.

The vapor bubbles were also considered. In this case the major problem is external heat problem in liquid. It is shown that the asymptotic expression for the heat flux at the interface in the case of gas bubbles has the similar structure as the heat flux from the vapor bubble surface to the liquid. In both cases it is Duhamel integral with rooted kernel.

© 2007 Elsevier Ltd. All rights reserved.

Keywords: Heat flux; Duhamel integral; Gas bubble; Vapor bubble

1. Introduction

Mass, force, and energy interactions proceeding at the interfaces in gas–liquid flows result in substantial variations in the flow velocity, pressure, and temperature field. To describe the processes of interface heat-exchange in gas–liquid bubble flows correctly, one has to analyze the interaction between a bubble and the carrier phase.

Oscillations of gas and vapor bubbles in liquid have been analyzed, both theoretically and experimentally, in a number of works which are discussed in detail in [1–3]. However, of both methodological and practical importance is the derivation of simple analytical dependencies for interface heat-exchange.

2. Formulation of the problem for the gas bubble

Let us consider the behavior of a gas bubble in a liquid. The surrounding liquid is incompressible and ideal. The

processes occurring within a bubble are assumed to be spherically symmetric. Phase transitions are ignored due to the low temperature of the gas and the liquid.

The pressure inside the bubble is assumed to be uniform [4–6]. It takes place when the length of sound wave in gas is much greater than the bubble radius. In the absence of phase transitions the temperature of a liquid remains practically unchanged, and the heat flux across the interface, q , is fully defined by the thermal resistance of a gas. Since no phase transitions take place, the heat flux across the boundary is continuous. Hence, q may be found by solving the internal heat-exchange problem for a bubble [4–7]

$$\frac{\partial T}{\partial t} + v \frac{\partial T}{\partial r} = \frac{a}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial T}{\partial r} \right) + \frac{\gamma - 1}{\gamma} \frac{T}{p} \frac{dp}{dt} \quad (2.1)$$

$$T(r, 0) = T_0, T(R, t) = T_0, \frac{\partial T}{\partial r} = 0(r = 0), p(0) = p_0 \quad (2.2)$$

The temperature of the bubble surface stays practically constant since the liquid has much higher thermal conductivity and much smaller thermal diffusivity than the gas [8].

* Tel.: +973 17437562; fax: +973 17449145.

E-mail address: nail@sci.uob.bh

Nomenclature

R	bubble radius	B	gas constant
\dot{R}	time derivative of the radius	γ	specific heat ratio
r	radial Euler coordinate	C	specific heat
x	longitudinal coordinate	C_p	specific heat of the gas at constant pressure
t	time	$P = \frac{p}{p_0}, \xi = \frac{r}{R}, \theta = \frac{T}{T_0}, \tau = \frac{t}{t_*}$ nondimensional parameters	
j	the phase transition rate	$t_* = \frac{R_0^2}{a_g}, u = \theta\xi, \tau_0 = x_1/\pi^2 = 0.1548, x_1 = 1.526$	
ℓ	the latent heat of evaporation	<i>Subscripts</i>	
$Pe = 2R_0(3\gamma p_0/\rho_e)^{1/2}a_g$	Peclet number	l	liquid
T	temperature	g	gas
z	dummy variable (nondimensional)	v	vapor
ρ	density	O	at equilibrium
p	pressure	R	at the bubble surface
v	radial velocity	S	at the saturation
V	longitudinal velocity		
λ	thermal conductivity		
a	thermal diffusivity		

The continuity equation for the gas, with using the uniformity of the pressure and the boundary condition $v(0, t) = 0$, yields the velocity profile in the bubble:

$$v(r, t) = \frac{r}{R}\dot{R} + \frac{\gamma - 1}{\gamma} \left[\lambda \frac{\partial T}{\partial r} - \frac{r}{R} \left(\lambda \frac{\partial T}{\partial r} \right)_R \right] \quad (2.3)$$

3. Analytic solution

The problem was solved by using the variable $\xi = r/R(t)$ which “freezes” the moving boundary of the bubble. By using formulas for the change of variables:

$$\begin{aligned} \frac{\partial}{\partial t} \Big|_r &= \frac{\partial}{\partial t} \Big|_\xi + \frac{\partial}{\partial \xi} \frac{\partial \xi}{\partial t} = \frac{\partial}{\partial t} \Big|_\xi - \frac{r}{R^2} \dot{R} \\ \frac{\partial}{\partial r} &= \frac{1}{R(t)} \frac{\partial}{\partial \xi} \end{aligned} \quad (3.1)$$

we obtain

$$\frac{\partial T}{\partial t} + \left(\frac{v - \xi \dot{R}}{R} \right) \frac{\partial T}{\partial \xi} = \frac{a}{R^2} \frac{\partial}{\partial \xi} \left(\xi^2 \frac{\partial T}{\partial \xi} \right) + \frac{\gamma - 1}{\gamma} \frac{T}{p} \frac{dp}{dt} \quad (3.2)$$

The estimates show that deviation of velocity profile inside the bubble from linear dependence due to the temperature gradient (2.3) is usually not exceed 20%. For this reason the convective term of heat Eq. (3.2) in new variables (ξ, t) can be neglected.

Let us consider the case when the deviation of bubble’s radius from equilibrium position is small enough. Then Eq. (3.2) can be simplified

$$\frac{\partial T}{\partial t} = \frac{a}{R_0^2} \frac{\partial}{\partial \xi} \left(\xi^2 \frac{\partial T}{\partial \xi} \right) + \frac{\gamma - 1}{\gamma} \frac{T}{p} \frac{dp}{dt} \quad (3.3)$$

Let us use nondimensional variables

$$P = \frac{p}{p_0}, \theta = \frac{T}{T_0}, \tau = \frac{t}{t_*}, t_* = \frac{R_0^2}{a_g} \quad (3.4)$$

Eq. (3.3) in nondimensional variables will have a form:

$$\frac{\partial \theta}{\partial \tau} = \frac{\partial}{\partial \xi} \left(\xi^2 \frac{\partial \theta}{\partial \xi} \right) + \frac{\gamma - 1}{\gamma} \frac{d(\ln P)}{d\tau} \theta \quad (3.5)$$

$$\theta(\xi, 0) = \theta_0, \theta(1, \tau) = \theta_0, \frac{\partial \theta}{\partial \xi}(0, \tau) = 0$$

$$P(0) = 1$$

Let us use a new variable $u = \theta\xi$. In new variables Eq. (3.5) will have a form:

$$\frac{\partial u}{\partial \tau} = \frac{\partial^2 u}{\partial \xi^2} + \frac{\gamma - 1}{\gamma} \frac{d(\ln P)}{d\tau} u \quad (3.6)$$

Solving this nonhomogeneous equation by Fourier’s method we will obtain the solution

$$\begin{aligned} u &= \xi \theta_0 + \sum_{n=1}^{\infty} a_n(\tau) \sin \pi n \xi \\ a_n(\tau) &= \int_0^\tau q_n(z) \exp[-(\pi n)^2(\tau - z)] dz \\ q_n(z) &= 2 \frac{\gamma - 1}{\gamma} \frac{d \ln P}{dz} \int_0^1 \xi \sin(\pi n \xi) d\xi \\ &= 2 \frac{\gamma - 1}{\gamma} \frac{d \ln P}{dz} \frac{(-1)^{n+1}}{\pi n} \\ u &= \xi \theta_0 + \frac{2(\gamma - 1)}{\gamma} \sum_{n=1}^{\infty} \int_0^\tau \frac{d(\ln P)}{dz} \frac{(-1)^{n+1}}{\pi n} \\ &\quad \times \exp[-(\pi n)^2(\tau - z)] dz \sin(\pi n \xi) \end{aligned} \quad (3.7)$$

Download English Version:

<https://daneshyari.com/en/article/661382>

Download Persian Version:

<https://daneshyari.com/article/661382>

[Daneshyari.com](https://daneshyari.com)